

SPECIFICATION

PHENOXYPROPYLAMINE COMPOUNDS

This is a continuation in part of PCT/JP00/03279 filed on May 22, 2000.

Technical Field

5 The present invention relates to a compound that acts on 5-hydroxytryptamine (5-HT) neurotransmission. More particularly, the present invention relates to a novel phenoxypropylamine compound having selective affinity for and
10 simultaneous antagonistic activity against a 5-hydroxytryptamine 1A (5-HT_{1A}) receptor in the central nervous system, as well as a 5-HT reuptake inhibitory activity, which is useful as a pharmaceutical agent, and to a therapeutic agent for depression and the like, which contains this compound. 5-
15 Hydroxytryptamine (5-HT) is also known as "serotonin".

Background Art

As a compound having an antagonistic activity against 5-HT_{1A} receptor as well as an inhibitory activity on the reuptake of 5-HT, there are known, for example, 1-(4-indolyloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol derivative (EP
20 0722941), 4-(4-fluorophenyl)-1-((6-(methylamino)indan-1-yl)methyl)piperidine derivative (WO 95/33721), 3,6-dihydro-N-methyl-N-(5-chloro-2-pyridyl)-4-(1-naphthalenyl)-1-(2H)pyridine propanamine derivative (US Patent No. 5472966), 3-(5-
25 chlorobenzo[b]thiophen-3-yl)-5,6-dihydroimidazo[2,1-b]thiazol derivative (WO 97/02269), S-(-)-N-(2-(3-(2-naphthyl)-pyrrolidino)ethyl)-N-(2-pyridyl)cyclohexanecarboxamide derivative (WO 97/40038), (R)-3-(N-cyclopentyl-N-n-propylamino)-8-fluoro-5-(N-methylcarbamoyl)-3,4-dihydro-2H-1-
30 benzopyran derivative (WO 96/33710), 3-(2-(4-methylpiperazin-1-yl)benzylidene)-1,3-dihydroindol-2-one derivative (WO 97/36867), (S)-1-(4-indolyloxy)-3-[4-hydroxy-4-(2-naphthyl)piperidino]-propan-2-ol derivative (WO 97/48698) and the like.

JP-A-62-116557 discloses substituted benzylactams, such

as 2-hydroxy-1-[2-((2-oxo-4-pyrrolidinyl)methyl)phenoxy]-3-(4-diphenylmethyl-piperazin-1-yl)propane and the like, which have a binding ability with a serotonin receptor and a muscarinic acetylcholine receptor, and which can be used for the treatment
5 of senile dementia, Alzheimer's disease, cerebrovascular dementia and the like.

Various diseases of the central nervous system (e.g., depression, anxiety) are considered to be caused by disorders of noradrenalin (NA) and 5-hydroxytryptamine (5-HT), which are
10 neurotransmitters. Accordingly, augmentation of 5-HTergic neurotransmission is considered to mainly influence depressive mood and anxious, whereas augmentation of noradrenergic neurotransmission is considered to influence retardation in depressive patients. The pharmaceutical agents, such as
15 imipramine, desipramine and the like, which are most frequently used for the treatment of depression, are considered to act on depressive patients by improving neurotransmission of one or both of these NA and 5-HT.

The activity of 5-HT is considered to relate to a number
20 of various types of psychiatric disorders. In addition, 5-HT has been considered to be responsible for various conditions (e.g., eating disorder, gastrointestinal injury, control of cardiovascular system and sexual behavior). However, conventional antidepressants, such as imipramine, desipramine
25 and the like, are defective in that they require 3 - 4 weeks or even longer time for the expression of an anti-depressive effect, which poses clinical problems.

A combined use of various pharmaceutical agents has been considered in an attempt to accelerate expression of effects of
30 antidepressants or to increase their efficacy (Journal of Clinical Psychiatry, Vol. 57; Supplement 7; pp 25-31). Therein, a noticeably shortened time for clinical expression of the effect by concurrent use of a selective serotonin (5-HT) reuptake inhibitor (SSRI) and a 5-HT_{1A} antagonist, pindolol,

has been reported (Journal of Clinical Psychopharmacology, Vol. 17, No. 6, pp. 446-450). It is known that the amount of 5-HT release in the brain does not increase much by SSRI alone, but when combined with a 5-HT_{1A} antagonist, the amount increases
5 markedly (Neurochemical Research, Vol. 21, No. 5, 1996, pp. 557-562). Under such circumstances, the "5-HT enhancement hypothesis" was proposed with regard to the expression of the action of antidepressants by Blier and de Montigny (Trends in Pharmacological Sciences, 1994, vol. 15, pp. 220-226). The 5-
10 HT enhancement hypothesis means that the effector mechanism of antidepressant rests in the enhancement of 5-HT release at a terminal. It is based on the understanding that the conventional antidepressants decrease the 5-HT release by single administration, but increase the 5-HT release and
15 express an anti-depressive effect only when they are administered consecutively. From those mentioned above, it is expected that a drug that promotes 5-HT release in the brain from the first can be a rapid onset antidepressant. In other words, a compound concurrently having a serotonin reuptake
20 inhibitory action and a 5-HT_{1A} antagonistic action is considered to be an antidepressant showing quick expression of an anti-depressive effect, namely, a rapid onset antidepressant.

It is an object of the present invention to find a
25 subgroup of 5-hydroxytryptamine (5-HT) receptor, namely, a compound simultaneously having selective affinity for and antagonistic activity against 5-HT_{1A} receptor in the central nervous system in mammals inclusive of human, which compound also having a 5-HT reuptake inhibitory activity.

30 It is therefore an object of the present invention to provide a compound that expresses an anti-depressive effect quickly, which is a so-called rapid onset antidepressant, and a compound useful for the treatment of 5-HT mediated diseases in the central nervous system, such as schizophrenia, anxiety


neurosis, obsessive-compulsive disorder (OCD), panic disorder, social anxiety disorder, seasonal emotional disorder, Anorexia Nervosa, Bulimia Nervosa, nocturnal enuresis, children's hyperlocomotion, post-traumatic stress disorder (PTSD), senile dementia, hemicrania, stroke, Alzheimer's disease, recognition disorder, hypertension, gastrointestinal injury, feeding disorders, premenstrual syndrome (PMS), abnormal body temperature regulation, sexual disorder and pain, as well as for the treatment of abnormality in the cardiovascular system, treatment of drug abuse and the like.

Summary of the Invention

The present inventors have conducted intensive studies, and as a result, found that a novel phenoxypropylamine compound of the formula (I), an optical isomer thereof and a pharmaceutically acceptable salt thereof have selective affinity for and simultaneous antagonistic activity against a 5-hydroxytryptamine 1A (5-HT_{1A}) receptor, as well as 5-HT reuptake inhibitory activity, and can be a useful pharmaceutical agent that meets the above-mentioned objects, which resulted in the completion of the present invention. Moreover, the present inventors have also found novel compounds of the formulas (II) and (III) to be mentioned below, which are the synthetic intermediates for the phenoxypropylamine compound.

Accordingly, the present invention provides the following.

1. A phenoxypropylamine compound of the formula (I)




wherein each symbol in the formula means as follows:

a bond represented by a solid line and a dotted line shows a double bond or a single bond;

X is a hydrogen atom, a hydroxy group, a C₁-C₈ alkoxy

group, an acyloxy group or an oxo group;
provided that when R^1 is a group of the following formula (2),
X should not be a hydrogen atom;

R^1 is a group of the following formula



wherein

Y is O or S,

Ar is optionally substituted aromatic hydrocarbon,

R^2 is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

R^5 is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

Z is void or $-CH_2-$, and

R^6 is hydrogen atom, hydroxy group, acetamido group, carboxyl group, alkoxycarbonyl group, cyano group or C_1-C_8 alkoxy group;

R^3 is a hydrogen atom, a C_1-C_{18} alkyl group or a halogen atom;

V is $-CH_2-$, $-O-$, $-S-$ or the formula $-N(R^4)-$


wherein R^4 is hydrogen atom, C_1-C_{18} alkyl group or optionally substituted aralkyl group;

W is void or $-CH_2-$ or $-C(=O)-$;

R^7 is a C_1-C_4 hydroxyalkyl group, an acyl group, an optionally substituted saturated or unsaturated heterocyclic group, an optionally substituted fused heterocyclic group, a C_1-C_4 alkylsulfonyl group or the formula $-Q-R^9$

wherein

Q is $-C(=O)-$, $-C(=S)-$, $-CH_2-$ or $-S(=O)_2-$, and
 R^9 is a group of the following formula



or $-NH-NH-R^{15}$

5 wherein R^{10} and R^{11} are each independently hydrogen atom, C_1-C_{18} alkyl group, optionally substituted aryl group, optionally substituted aralkyl group or alkoxy group, R^{12} is hydrogen atom, optionally substituted aryl group, C_1-C_{18} alkyl group, C_1-C_8 alkoxy group or acyl group, and R^{15} is hydrogen atom, phenyl group, C_1-C_4 alkyl group, C_1-C_2 halogenated alkyl group, halogen atom, C_2-C_4 alkenyl group, C_1-C_4 hydroxyalkyl group, alkoxyalkyl group, alkyloxycarbonyl group, optionally substituted amino group, acetamido group, carboxyl group, acyl group, optionally substituted alkyloxy group, alkylthio group or cyano group;

15 provided that when R^1 is a group of the above formula (2), R^7 should not be C_1-C_4 hydroxyalkyl group or acyl group, and R^{10} and R^{11} are not each hydrogen atom at the same time; or

20

R⁷ and W in combination may form a ring of the following formula



wherein


- 5 E is oxygen atom or sulfur atom, and
 Q' is an optionally substituted 4 to 7-membered
 heterocycle having 1 or 2 hetero atom(s) selected
 from the group consisting of nitrogen atom and
 oxygen atom in the ring, in which case V is
 10 hydrogen atom; and

Ra, Rb and Rc are each independently a hydrogen atom, a C₁-C₁₈
 alkyl group, a hydroxy group, a C₁-C₈ alkoxy group,
 a halogen atom, an acyl group, a nitro group or an
 amino group;

- 15 provided that when R⁷ and W are bonded to form a ring of the
 above formula (14), Ra, Rb and Rc are not each hydroxy group or
 C₁-C₈ alkoxy group;

an optically active compound thereof, a pharmaceutically
 acceptable salt thereof or a hydrate thereof.

- 20 2. The compound of the aforementioned 1, which is represented
 by the formula (I)




wherein each symbol in the formula means as follows:

a bond represented by a solid line and a dotted line shows a
 25 double bond;

X is a hydrogen atom, a hydroxy group, a C₁-C₈ alkoxy
 group, an acyloxy group or an oxo group;

R¹ is a group of the following formula



wherein

Y is O or S,

Ar is optionally substituted benzene or naphthalene,

5 R^2 is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

R^5 is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

Z is void or $-\text{CH}_2-$, and

10 R^6 is hydrogen atom, hydroxy group, acetamido group, carboxyl group, alkoxycarbonyl group, cyano group or C_1 - C_8 alkoxy group;

R^3 is a hydrogen atom, a C_1 - C_{18} alkyl group or a halogen atom;

15 V is $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$ or the formula $-\text{N}(\text{R}^4)-$

wherein R^4 is hydrogen atom, C_1 - C_{18} alkyl group or optionally substituted aralkyl group;


W is void or $-\text{CH}_2-$ or $-\text{C}(=\text{O})-$;

20 R^7 is a C_1 - C_4 hydroxyalkyl group, an acyl group, an optionally substituted saturated or unsaturated heterocyclic group, an optionally substituted fused heterocyclic group, a C_1 - C_4 alkylsulfonyl group or the formula $-\text{Q}-\text{R}^9$

wherein

25 Q is $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{CH}_2-$ or $-\text{S}(=\text{O})_2-$, and

R^9 is a group of the following formula



or —NH—NH—R^{15}

wherein R^{10} and R^{11} are each independently hydrogen atom, $\text{C}_1\text{—C}_{18}$ alkyl group, optionally substituted aryl group, optionally substituted aralkyl group or alkoxy group, R^{12} is hydrogen atom, optionally substituted aryl group, $\text{C}_1\text{—C}_{18}$ alkyl group, $\text{C}_1\text{—C}_8$ alkoxy group or acyl group, and R^{15} is hydrogen atom, phenyl group, $\text{C}_1\text{—C}_4$ alkyl group, $\text{C}_1\text{—C}_2$ halogenated alkyl group, halogen atom, $\text{C}_2\text{—C}_4$ alkenyl group, $\text{C}_1\text{—C}_4$ hydroxyalkyl group, alkoxyalkyl group, alkyloxycarbonyl group, optionally substituted amino group, acetamido group, carboxyl group, acyl group, optionally substituted alkyloxy group, alkylthio group or cyano group; and

Ra , Rb and Rc are each independently a hydrogen atom, a $\text{C}_1\text{—C}_{18}$ alkyl group, a hydroxy group, a $\text{C}_1\text{—C}_8$ alkoxy group, a halogen atom, an acyl group, a nitro group or an amino group;

provided that when R^1 is a group of the above formula (2), R^7


should not be C₁-C₄ hydroxyalkyl group or acyl group, and R¹⁰ and R¹¹ are not each hydrogen atom at the same time; an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

5 3. The compound of the aforementioned 2, which is represented by the formula (I) wherein each symbol in the formula means as follows:

a bond represented by a solid line and a dotted line shows a double bond;

10 X is a hydroxy group;

R¹ is a group of the following formula



wherein

R⁵ is optionally substituted phenyl group or naphthyl group,

Z is void, and

R⁶ is hydrogen atom;


R³ is a hydrogen atom or a C₁-C₄ alkyl group;

V is -CH₂-, -O-, -S- or -N(R⁴)-

20 wherein R⁴ is hydrogen atom, C₁-C₆ alkyl group or optionally substituted aralkyl group;

W is void;

R⁷ is a group of the following formula








25 or the formula -CO-R⁹


wherein

R⁸ is hydrogen atom, phenyl group, C₁-C₄ alkyl group, C₁-C₂ halogenated alkyl group, halogen atom, C₂-C₄ alkenyl group, C₁-C₄ hydroxyalkyl group,

alkoxyalkyl group, alkyloxycarbonyl group,
optionally substituted amino group, acetamido group,
carboxyl group, acyl group, optionally substituted
alkyloxy group, alkylthio group or cyano group, and
5 R^9 is a group of the following formula



or




wherein R^{10} and R^{11} are each independently hydrogen
atom, C_1 - C_{18} alkyl group, optionally substituted aryl
group, optionally substituted aralkyl group or
10 alkoxy group, and R^{12} is hydrogen atom, optionally
substituted aryl group, C_1 - C_{18} alkyl group, C_1 - C_8
alkoxy group or acyl group; and

R_a , R_b and R_c are each a hydrogen atom;

an optically active compound thereof, a pharmaceutically

15 acceptable salt thereof or a hydrate thereof.

4. The compound of the aforementioned 2 or 3, which is
represented by the formula (I')



wherein each symbol is as in the aforementioned 2,


20 an optically active compound thereof, a pharmaceutically
acceptable salt thereof or a hydrate thereof.

5. The compound of the aforementioned 2, which is selected from

the group consisting of

- (1) 1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-ylcarbonyl)pyrrolidine,
- (2) 4-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-ylcarbonyl)morpholine,
- 5 (4) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)furan-2-carboxamide,
- (12) 1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)thiophen-2-ylcarbonyl)pyrrolidine,
- 10 (13) 4-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)thiophen-2-ylcarbonyl)morpholine,
- (15) 4-(2-hydroxy-3-(4-(naphthalen-1-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide,
- (17) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide,
- 15 (20) 4-(7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-ylcarbonyl)morpholine,
- (21) 7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)furan-2-carboxamide,
- 20 (27) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethyl-1H-indole-2-carboxamide,
- (30) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethyl-1-methylindole-2-carboxamide,
- (35) 1-(2-(5-methyl-1,2,4-oxadiazol-3-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
- 25 (37) 1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
- (38) 1-(2-(5-trifluoromethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
- 30 (39) 1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
- (42) 1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indole-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,


- (44) 1-(2-(3-methyl-1,2,4-oxadiazol-5-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
 (48) 1-(2-(5-methyloxazol-2-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
 5 (81) 3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol,
 (88) 1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol, and
 (93) 3-(4-(3,4-dimethylphenyl)piperidino)-1-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol,
 10 an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.
 6. The compound of the aforementioned 1, which is represented by the formula (I)



15 wherein each symbol in the formula means as follows:
 a bond represented by a solid line and a dotted line shows a double bond or a single bond;

X is a hydrogen atom, a hydroxy group, a C₁-C₈ alkoxy group or an acyloxy group;

20 R¹ is a group of the following formula



wherein

R² is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

25 R⁵ is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

Z is void or -CH₂-, and

R⁶ is hydrogen atom, hydroxy group or C₁-C₈ alkoxy

group;

R^3 is a hydrogen atom, a C_1-C_{18} alkyl group or a halogen atom;

R^7 and W are bonded to form a ring of the following formula



5

wherein

E is an oxygen atom or a sulfur atom, and

Q' is an optionally substituted 4 to 7-membered heterocycle having 1 or 2 hetero atom(s) selected from the group consisting of nitrogen atom and oxygen atom in the ring,

10

and V is hydrogen atom; and

R_a , R_b and R_c are each independently a hydrogen atom, a C_1-C_{18} alkyl group, a halogen atom, an acyl group, a nitro

15

group or an amino group;

an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.




7. The compound of the aforementioned 6, which is represented by the formula (I) wherein each symbol in the formula means as

20 follows:


a group of the following formula



is a group of the following formula



or



25 wherein


E is an oxygen atom or a sulfur atom,
 q is 0, 1, 2 or 3,
 R^{4'}, R^{7'} and R^{8'} are each independently a hydrogen atom, a C₁-C₁₈
 alkyl group, an optionally substituted aryl group or
 5 an optionally substituted aralkyl group, and
 other symbols are as defined in the aforementioned 6,
 an optically active compound thereof, a pharmaceutically
 acceptable salt thereof or a hydrate thereof.

8. The compound of the aforementioned 6, which is represented
 10 by the formula (I) wherein each symbol in the formula means as
 follows:

a bond represented by a solid line and a dotted line shows a
 double bond;

X is a hydroxy group;

15 R¹ is a group of the following formula



wherein

R⁵ is optionally substituted phenyl group or naphthyl
 group,


20 Z is void, and

R⁶ is hydrogen atom;

R³ is a hydrogen atom or a C₁-C₄ alkyl group;
 a group of the following formula



25 is a group of the following formula




wherein q is 1 and R^{4'} is hydrogen atom or C₁-C₄ alkyl

group; and

Ra, Rb and Rc are each a hydrogen atom;

an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

- 5 9. The compound of the aforementioned 6, which is represented by the formula (I'')



wherein each symbol is as defined in the aforementioned 6, an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

10 10. The compound of the aforementioned 6, which is selected from the group consisting of

(306) 5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-1,3-dimethylimidazolidine-2,4-dione,

15 (307) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone,

(308) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone,

20 (309) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone,

(310) α -(2'-(3-(4-(3-fluoro-4-methylphenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,

(311) α -(2'-(3-(4-(3,4-dimethylphenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,

25 (312) α -(2'-(3-(4-(4-chloro-3-fluorophenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,

(313) α -(2'-(3-(4-(4-chloro-3-trifluoromethylphenyl)-piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,

(314) α -(2'-(2-hydroxy-3-(4-(naphthalen-1-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone,

(315) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- δ -valerolactone,

5 (316) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -valerolactone,

(319) 3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-2-pyrrolidone,

(322) 3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-1-methyl-2-pyrrolidone, and

(325) α -(2'-(2-hydroxy-3-(4-(6-methoxynaphthalen-2-yl)piperidino)propyloxy)benzylidene)- γ -butyrolactone,

an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

15 11. A pharmaceutical agent comprising a compound of the aforementioned 1, an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

12. The pharmaceutical agent of the aforementioned 11, which is an agent for the treatment of depression.


20 13. A pharmaceutical composition comprising at least one member selected from the group consisting of a compound of the aforementioned 1, an optically active compound thereof, a pharmaceutically acceptable salt thereof and a hydrate thereof, and a pharmaceutically acceptable carrier.

25 14. The pharmaceutical composition of the aforementioned 13, which is an agent for the treatment of depression.

15. A 5HT_{1A} antagonist having a selective serotonin reuptake inhibitory action, which comprises a compound of the aforementioned 1, an optically active compound thereof, a
30 pharmaceutically acceptable salt thereof or a hydrate thereof.


16. A selective serotonin reuptake inhibitor having a 5HT_{1A} antagonistic action, which comprises a compound of the aforementioned 1, an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

17. A compound of the formula (II)



wherein each symbol in the formula means as follows:

X is a hydrogen atom, a hydroxy group, a C₁-C₈ alkoxy group or an acyloxy group or an oxo group;
 R¹ is a group of the following formula



wherein

Y is O or S,

Ar is optionally substituted benzene or naphthalene,

R² is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

R⁵ is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

Z is void or -CH₂-, and

R⁶ is hydrogen atom, hydroxy group, acetamido group, carboxyl group, alkoxycarbonyl group, cyano group or C₁-C₈ alkoxy group,

provided that when V is -N(R⁴)-, R⁶ should not be hydroxy group;

R³ is a hydrogen atom, a C₁-C₁₈ alkyl group or a halogen atom;

V is -CH₂-, -O-, -S- or the formula -N(R⁴)-

wherein

R⁴ is hydrogen atom, C₁-C₁₈ alkyl group or optionally substituted aralkyl group;


W is void, -CH₂- or -C(=O)-;

R¹⁴ is a hydrogen atom or a C₁-C₄ alkyl; and

5 Ra, Rb and Rc are each independently a hydrogen atom, a C₁-C₁₈ alkyl group, a hydroxy group, a C₁-C₈ alkoxy group, a halogen atom, an acyl group, a nitro group or an amino group;

an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

18. A compound of the formula (III)



wherein each symbol is as follows:

R is an allyl group or a 2,3-epoxypropan-1-yl group;

15 a bond represented by a solid line and a dotted line shows a double bond or a single bond;

E is an oxygen atom or a sulfur atom;

R³ is a hydrogen atom, a C₁-C₁₈ alkyl group or a halogen atom;

20 Q' is an optionally substituted 4 to 7-membered heterocycle having 1 or 2 hetero atom(s) selected from the group consisting of nitrogen atom and oxygen atom in the ring; and


Ra, Rb and Rc are each independently a hydrogen atom, a C₁-C₁₈ alkyl group, a hydroxy group, a C₁-C₈ alkoxy group, a halogen atom, an acyl group, a nitro group or an amino group;

an optically active compound thereof, a pharmaceutically acceptable salt thereof or a hydrate thereof.

19. The compound of the aforementioned 18, wherein, in the formula (III), each symbol is as follows:
the group of the following formula



5 is a group of the following formula



wherein

E is oxygen atom or sulfur atom,

q is 0, 1, 2 or 3,

10 R^{4'}, R^{7'} and R^{8'} are each independently hydrogen atom, C₁-C₁₈ alkyl group, optionally substituted aryl group or optionally substituted aralkyl group, and

other symbols are as defined in the aforementioned 18,
an optically active compound thereof, a pharmaceutically

15 acceptable salt thereof or a hydrate thereof.

20. A compound selected from the group consisting of

2-(4-methoxybenzo(b) furan-2-yl)-5-methyl-1,3,4-oxadiazole,

2-(4-hydroxybenzo(b) furan-2-yl)-5-methyl-1,3,4-oxadiazole,

(S)-2-(4-glycidyloxybenzo(b) furan-2-yl)-5-methyl-1,3,4-

20 oxadiazole,

2-(7-methoxybenzo(b) furan-2-yl)-5-methyl-1,3,4-oxadiazole,

2-(4-(methoxymethyloxy) benzo(b) thiophen-2-yl)-5-methyl-1,3,4-oxadiazole,

2-(4-hydroxybenzo(b) thiophen-2-yl)-5-methyl-1,3,4-oxadiazole,

25 4-benzyloxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indole,

2-(7-methoxybenzo(b) furan-2-yl)-5-phenyl-1,3,4-oxadiazole,

2-(4-methoxybenzo(b) furan-2-yl)-5-trifluoromethyl-1,3,4-

oxadiazole,
 2-(4-hydroxybenzo (b) furan-2-yl)-5-trifluoromethyl-1,3,4-
 oxadiazole,
 (S)-2-(4-glycidyloxybenzo (b) furan-2-yl)-5-trifluoromethyl-
 5 1,3,4-oxadiazole,
 2-(7-methoxybenzo (b) furan-2-yl)-5-trifluoromethyl-1,3,4-
 oxadiazole,
 2-(7-hydroxybenzo (b) furan-2-yl)-5-trifluoromethyl-1,3,4-
 oxadiazole,
 10 (S)-2-(7-glycidyloxybenzo (b) furan-2-yl)-5-trifluoromethyl-
 1,3,4-oxadiazole,
 N'-(4-methoxybenzo (b) furan-2-ylcarbonyl)propionohydrazide,
 2-(4-methoxybenzo (b) furan-2-yl)-5-ethyl-1,3,4-oxadiazole,
 2-(4-hydroxybenzo (b) furan-2-yl)-5-ethyl-1,3,4-oxadiazole,
 15 (S)-2-(4-glycidyloxybenzo (b) furan-2-yl)-5-ethyl-1,3,4-
 oxadiazole,
 2-(4-methoxybenzo (b) furan-2-yl)-5-methyl-1,3,4-thiadiazole,
 2-(4-hydroxybenzo (b) furan-2-yl)-5-methyl-1,3,4-thiadiazole,
 (S)-2-(4-glycidyloxybenzo (b) furan-2-yl)-5-methyl-1,3,4-
 20 thiadiazole,
 5-ethoxycarbonyl-2-(4-methoxybenzo (b) furan-2-yl)-1,3,4-
 oxadiazole,
 5-ethoxycarbonyl-2-(4-hydroxybenzo (b) furan-2-yl)-1,3,4-
 oxadiazole,
 25 5-(4-(methoxymethyloxy) benzo (b) furan-2-yl)-2,3-dihydro-1,3,4-
 oxadiazole-2-thione,
 5-(4-(methoxymethyloxy) benzo (b) furan-2-yl)-2-methylthio-1,3,4-
 oxadiazole,
 5-(4-hydroxybenzo (b) furan-2-yl)-2-methylthio-1,3,4-oxadiazole,
 30 5-(4-(methoxymethyloxy) benzo (b) furan-2-yl)-2,3-dihydro-1,3,4-
 oxadiazol-2-one,
 5-(4-(methoxymethyloxy) benzo (b) furan-2-yl)-2-methoxy-1,3,4-
 oxadiazole,
 (S)-5-(4-glycidyloxybenzo (b) furan-2-yl)-2-methoxy-1,3,4-

oxadiazole,

2-ethoxy-5-(4-(methoxymethyloxy)benzo(b) furan-2-yl)-1,3,4-

oxadiazole,

(S)-2-ethoxy-5-(4-glycidyloxybenzo(b) furan-2-yl)-1,3,4-

5 oxadiazole,

2-(1-methylethyloxy)-5-(4-(methoxymethyloxy)benzo(b) furan-2-yl)-1,3,4-oxadiazole and

(S)-2-(1-methylethyloxy)-5-(4-glycidyloxybenzo(b) furan-2-yl)-1,3,4-oxadiazole.

10 Detailed Description of the Invention

The definitions and examples of each group in the formulas (I), (I'), (I''), (II) and (III) are shown in the following.

The acyloxy group at X is, for example, acetoxy,
15 propionyloxy, butyryloxy, benzoyloxy and the like, preferably acetoxy.

The "aryl group" of the optionally substituted aryl group at Ar, R², R⁵, R¹⁰, R¹¹, R¹², R^{4'}, R^{7'} and R^{8'} is, for example, phenyl, naphthyl, tetrahydronaphthyl (e.g., 1,2,3,4-
20 tetrahydronaphthalene-6-yl, 5,6,7,8-tetrahydronaphthalen-2-yl etc.), indanyl (e.g., indan-5-yl etc.), indenyl (e.g., inden-5-yl etc.) and the like, with preference given to phenyl and naphthyl. These may be substituted by one or more, the same or different substituents mentioned below. A hydrogen atom may be
25 added to the double bond of these aryl groups. Examples of the "substituent" include halogen atom (e.g., chlorine atom, fluorine atom etc.), trifluoromethyl, C₁-C₄ alkyl group (linear or branched chain, such as methyl, ethyl, propyl, isopropyl, butyl etc.), C₁-C₄ alkoxy group (linear or branched chain, such
30 as methoxy, ethoxy, propoxy, isopropoxy, butoxy etc.), aryl group (e.g., phenyl etc.), aralkyl group (e.g., benzyl etc.), oxo group, alkoxyalkyl group (e.g., methoxyethyl etc.) and the like, with preference given to chlorine atom, fluorine atom, trifluoromethyl, methyl, methoxy, phenyl, benzyl, oxo,

methoxyethyl and the like.

Preferable examples of the optionally substituted aryl group at R⁵ include naphthyl (1-naphthyl, 2-naphthyl), 4-chloro-3-fluorophenyl, 3-chloro-4-trifluoromethylphenyl, 3,4-dimethylphenyl, 3,4-dichlorophenyl, 2,4- or 3,4-dimethylphenyl, 4-methylphenyl, 4-fluorophenyl, 3-chloro-4-methylphenyl, 4-chloro-3-trifluoromethylphenyl, 6-methoxynaphthyl-2-yl, 4-chlorophenyl, 3,4-difluorophenyl, 3,4-dimethoxyphenyl, 3-chlorophenyl, 4-chloro-3-methylphenyl, 4-chloro-3-methoxyphenyl, 2,5-dichlorophenyl, 4-chloro-3-trifluorophenyl and the like.

Examples of the "aromatic hydrocarbon" of the optionally substituted aromatic hydrocarbon at Ar include benzene, naphthalene and the like, which may be substituted by one or more, the same or different substituents mentioned below. The "substituent" is, for example, halogen atom (e.g., chlorine atom, fluorine atom etc.), trifluoromethyl, C₁-C₄ alkyl group (linear or branched chain, such as methyl, ethyl, propyl, isopropyl, butyl etc.), C₁-C₄ alkoxy group (linear or branched chain, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy etc.), aryl group (e.g., phenyl etc.), aralkyl group (e.g., benzyl etc.), oxo group, alkoxyalkyl group (e.g., methoxyethyl etc.) and the like, with preference given to chlorine atom, fluorine atom, trifluoromethyl, methyl, methoxy, phenyl, benzyl, oxo group, methoxyethyl and the like.

The "aromatic heterocyclic group" of the optionally substituted aromatic heterocyclic group at R² and R⁵ is, for example, pyridyl, furyl, thienyl, pyrimidinyl, indolyl (e.g., indol-2-yl, indol-6-yl, indol-5-yl etc.), benzo(b)thienyl (e.g., benzo(b)thiophen-2-yl, benzo(b)thiophen-5-yl, 2,3-dihydrobenzo(b)thiophen-6-yl, 2,3-dihydrobenzo(b)thiophen-5-yl etc.), benzo(b)furyl (e.g., benzo(b)furan-2-yl, benzo(b)furan-5-yl, benzo(b)furan-6-yl, 2,3-dihydrobenzo(b)furan-5-yl, 2,3-dihydrobenzo(b)furan-4-yl, 3,4-dihydro-2H-benzo(b)furan-6-yl, 2,3-dihydrobenzo(b)furan-6-yl etc.), 3,4-methylenedioxyphenyl,

benzimidazolyl (e.g., 2,3-dihydrobenzimidazol-1-yl etc.), 1,4-benzodioxanyl (e.g., 1,4-benzodioxan-6-yl etc.), chromanyl (e.g., chroman-6-yl, chroman-7-yl etc.), indolinyl (e.g., indolin-5-yl etc.), chromenyl (e.g., 2H-chromen-6-yl, 2H-chromen-7-yl etc.), benzo(b)thiiny (e.g., 3,4-dihydro-2H-benzo(b)thiin-7-yl, 3,4-dihydro-2H-benzo(b)thiin-6-yl etc.), benzoisoxazolyl (e.g., benzoisoxazol-5-yl, benzo(d)isoxazol-5-yl etc.), benzo(c)furyl (e.g., 1,3-dihydrobenzo(c)furan-5-yl etc.), isochromanyl (e.g., isochroman-7-yl, isochroman-6-yl etc.), quinolinyl (e.g., quinolin-3-yl, quinolin-6-yl etc.), 3,4-dihydro-2H-benzo(b)oxin-6-yl, 3,4-dihydro-2H-benzo(c)oxin-6-yl, isoindolinyl (e.g., isoindolin-5-yl etc.), isoquinolinyl (e.g., isoquinolin-6-yl etc.) and the like, which may be substituted by one to three the same or different substituents mentioned below.

Examples of the "substituent" include halogen atom (e.g., fluorine atom, chlorine atom, bromine atom etc.), haloalkyl (e.g., fluoromethyl, difluoromethyl, trifluoromethyl etc.), C₁-C₄ alkyl (linear or branched chain, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl etc.), C₁-C₈ alkoxy group (linear or branched chain, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, hexyloxy, heptyloxy, octyloxy etc.), hydroxy, nitro, cyano, amino, mono or dialkylamino wherein each alkyl has 1 to 4 carbon atoms (e.g., methylamino, dimethylamino, diethylamino, dipropylamino etc.), acyl (e.g., acetyl, propionyl, butyryl etc.), C₂-C₆ alkenyl (e.g., vinyl, 1-propenyl, 2-propenyl, 3-propenyl etc.), C₂-C₆ alkynyl (e.g., ethynyl, 1-propynyl, 2-propynyl etc.), phenyl, phenoxy, benzyloxy, R'-S(O)t- wherein R' is C₁-C₄ alkyl and t is 0, 1 or 2, Ph-S(O)t- wherein Ph is phenyl and t is as defined above, carbamoyl, N,N-dialkylcarbamoyl (e.g., N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl, N,N-dipropylcarbamoyl etc.), oxo and the like, with preference given to C₁-C₄ alkyl.

fluoromethyl, difluoromethyl, trifluoromethyl etc.), C₁-C₄ alkyl (linear or branched, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl etc.), C₁-C₈ alkoxy (linear or branched, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, hexyloxy, heptyloxy, octyloxy etc.), hydroxy, nitro, cyano, amino and the like. The optionally substituted aralkyl at R^{4'}, R^{7'} and R^{8'} is, for example, benzyl, 2-phenylethyl, 1-phenylethyl, 1,1-dimethyl-2-phenylethyl, 3-phenylpropyl, 2-phenylpropyl, 1-phenylpropyl, 4-phenylbutyl, 3-phenylbutyl, 2-phenylbutyl, 1-phenylbutyl and the like, with preference given to benzyl.

The C₁-C₂ halogenated alkyl group at R⁸ and R¹⁵ is, for example, chloromethyl, dichloromethyl, trichloromethyl, bromomethyl, dibromomethyl, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl and the like, preferably trichloromethyl and trifluoromethyl.

The halogen atom at R⁸ and R¹⁵ is fluorine atom, chlorine atom, bromine atom or iodine atom, preferably fluorine atom and chlorine atom.

The C₂-C₄ alkenyl group at R⁸ and R¹⁵ is linear or branched chain, such as vinyl, 1-propenyl, allyl, 1-butenyl, 2-butenyl, isopropenyl and the like, with preference given to vinyl, 1-propenyl and isopropenyl.

The C₁-C₄ hydroxyalkyl group at R⁸ and R¹⁵ is, for example, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 3-hydroxypropyl, 4-hydroxybutyl, 1-hydroxy-1-methylethyl, 2-hydroxy-1-methylethyl, 1,2-dihydroxy-1-methylethyl and the like, with preference given to hydroxymethyl.

In the alkoxyalkyl group at R⁸ and R¹⁵, the "alkoxy" moiety is preferably C₁-C₄ linear or branched chain alkoxy and the "alkyl" moiety is preferably C₁-C₄ linear or branched chain alkyl. Examples thereof include methoxymethyl, ethoxymethyl, propyloxymethyl, methoxyethyl, ethoxyethyl and the like, with preference given to methoxymethyl.

In the alkyloxycarbonyl group at R^8 and R^{15} , the "alkyloxy" moiety is preferably C_1 - C_4 linear or branched chain alkyloxy, such as methoxycarbonyl, ethoxycarbonyl, propyloxycarbonyl, butyloxycarbonyl and the like, with
5 preference given to methoxycarbonyl and ethoxycarbonyl.

The optionally substituted amino group at R^8 and R^{15} is preferably amino optionally substituted by one or two, the same or different C_1 - C_2 alkyl. Examples thereof include amino, methylamino, dimethylamino, ethylamino, diethylamino and the
10 like, with preference given to methylamino and dimethylamino.

The acyl group at R^8 and R^{15} is, for example, acetyl, propionyl, butyryl, isobutyryl and the like, with preference given to acetyl.

The optionally substituted alkyloxy group at R^8 and R^{15}
15 is preferably, C_1 - C_4 linear or branched chain, which may be substituted by one or more, the same or different substituents mentioned below. Examples of these "substituents" include fluorine atom, chlorine atom and the like. Specific examples thereof include methoxy, ethoxy, propoxy, isopropoxy, butyloxy,
20 trifluoromethoxy, 2,2,2-trifluoroethyloxy and the like, with preference given to methoxy and 2,2,2-trifluoroethyloxy.

The alkylthio group at R^8 and R^{15} is preferably C_1 - C_4 linear or branched chain, such as methylthio, ethylthio, propylthio, isopropylthio, butylthio and the like, with
25 preference given to methylthio and ethylthio.

The C_1 - C_4 hydroxyalkyl group at R^7 is linear or branched chain, such as 1-hydroxymethyl, 1-hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 1-hydroxybutyl and the like, with preference given to 1-hydroxyethyl.

30 The C_1 - C_4 alkylsulfonyl group at R^7 is linear or branched chain, such as methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl and the like, with preference given to ethylsulfonyl.

The optionally substituted saturated or unsaturated

heterocyclic group at R⁷ is a 5 or 6-membered heterocyclic group optionally containing 1 - 3 hetero atom(s) selected from the group consisting of nitrogen atom, oxygen atom and sulfur atom, such as a group derived from furan, thiophene, pyrrole, pyrazole, oxazole, isoxazole, thiazole, isothiazole, imidazole, furazan, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, pyridine, pyrimidine, pyrazine, pyridazine, oxazoline, thiazoline, imidazoline, tetrahydrofuran, tetrahydrothiophene, pyran and the like. Preferred are groups derived from thiophene, pyrazole, oxazole, isoxazole, thiazole, imidazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, 1,3,4-oxadiazole and the like and more preferred are groups derived from oxazole, thiazole, 1,3,4-thiadiazole, 1,3,4-oxadiazole and the like. These may be substituted by one or two the same or different substituents mentioned below.

Examples of these substituents include optionally substituted aryl group (e.g., phenyl or naphthyl optionally substituted by halogen atom, amino, nitro, hydroxy, C₁-C₄ alkyl, C₁-C₄ alkoxy and the like), C₁-C₁₈ alkyl group (as defined above, preferably methyl, ethyl, isopropyl, tert-butyl, isobutyl etc.), C₁-C₂ halogenated alkyl group (as defined above), C₁-C₈ alkoxy group (as defined above, preferably methoxy, ethoxy, isopropoxy etc.), halogen atom (e.g., fluorine atom, chlorine atom, bromine atom or iodine atom), C₂-C₄ alkenyl group (e.g., vinyl, 1-propenyl, allyl, 1-butenyl, 2-butenyl, isopropenyl etc.), C₁-C₄ hydroxyalkyl group (e.g., hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl, 3-hydroxypropyl, 4-hydroxybutyl etc.), alkoxyalkyl group (e.g., methoxymethyl, ethoxymethyl, propyloxymethyl, methoxyethyl, ethoxyethyl etc.), alkyloxycarbonyl group (e.g., methoxycarbonyl, ethoxycarbonyl, propyloxycarbonyl, butyloxycarbonyl etc.), optionally substituted amino group (e.g., amino, methylamino, dimethylamino, ethylamino, diethylamino etc.), acyl group (e.g., acetyl, propionyl, butyryl, isobutyryl etc.), acetamido group,

carboxyl group, optionally substituted alkyloxy group (e.g., methoxy, ethoxy, propoxy, isopropoxy, butyloxy, trifluoromethoxy, 2,2,2-trifluoromethoxy etc.), alkylthio group (e.g., methylthio, ethylthio, propylthio, isopropylthio, butylthio etc.), cyano group and the like.

Examples of the optionally substituted fused heterocyclic group at R^7 include groups derived from benzofuran, benzothiophene, indole, benzoxazole, benzothiazole, 1,2-benzoisoxazole, 1,2-benzisothiazole, benzimidazoline and the like, with preference given to benzoxazol-2-yl and benzothiazol-2-yl. These may be substituted by one or more, the same or different substituents mentioned below. Examples of these substituents include halogen atom (e.g., fluorine atom, chlorine atom, bromine atom etc.), haloalkyl group (e.g., fluoromethyl, difluoromethyl, trifluoromethyl etc.), C_1-C_4 alkyl group (linear or branched, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl etc.), C_1-C_8 alkoxy group (linear or branched, such as methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, hexyloxy, heptyloxy, octyloxy etc.), hydroxy group, nitro group, cyano group, amino group and the like.

The alkoxycarbonyl group at R^6 is preferably C_1-C_4 linear or branched chain, such as ethoxycarbonyl, methoxycarbonyl, propoxycarbonyl, butoxycarbonyl and the like, with preference given to ethoxycarbonyl.

The C_1-C_4 alkyl group at R^{14} is linear or branched, such as methyl, ethyl, propyl, isopropyl, butyl, tert-butyl and the like, with preference given to methyl and ethyl.

The alkoxy group at R^{10} and R^{11} is, for example, linear or branched chain alkoxy having 1 to 4, preferably 1 or 2, carbon atoms, such as methoxy, ethoxy and the like, with preference given to methoxy.

The "cycloalkylene group" of the optionally substituted C_3-C_8 cycloalkylene group at Y' is, for example, cyclopropylene,

5 cyclobutylene, cyclopentylene, cyclohexylene, cycloheptylene,
cyclooctylene and the like. Examples of the substituent include
C₁-C₄ alkyl group (e.g., methyl, ethyl, propyl, isopropyl,
butyl, isobutyl, tert-butyl etc.), C₁-C₈ alkoxy group (e.g.,
10 methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tert-
butoxy, pentyloxy, hexyloxy, heptyloxy, octyloxy etc.), hydroxy
group, oxo group and the like. Examples of the optionally
substituted C₃-C₈ cycloalkylene include 2-methoxycyclopentylene,
2-methylcyclohexylene, 2,6-dimethylcyclohexylene, 3-
10 ethylcycloheptylene, 3-hydroxycycloheptylene and the like, with
preference given to 2,6-dimethylcyclohexylene.

The C₁-C₈ alkylene group at Y' is, for example, linear
or branched, such as methylene, ethylene, trimethylene,
tetramethylene, pentamethylene, hexamethylene, heptamethylene,
15 octamethylene, methylenemethylene, dimethylenemethylene, 1-
methylethylene, 2-methylethylene, 1,1-dimethylethylene, 2,2-
dimethylethylene, ethylenemethylene, diethylenemethylene, 1-
ethylethylene, 2-ethylethylene, 1-methyltrimethylene, 1,1-
dimethyltrimethylene, 2-methyltrimethylene, 2,2-
20 dimethyltrimethylene, 3-methyltrimethylene, 3,3-
dimethyltrimethylene, 1-ethyltrimethylene, 2-ethyltrimethylene,
3-ethyltrimethylene and the like, with preference given to
ethylene, trimethylene and tetramethylene.


The C₁-C₄ alkyleneoxy group at Y' is, for example,
25 linear or branched, such as methyleneoxy, ethyleneoxy,
trimethyleneoxy, tetramethyleneoxy, methylenemethyleneoxy,
dimethylenemethyleneoxy, 1-methylethyleneoxy, 2-methylethyleneoxy,
1,1-dimethylethyleneoxy, 2,2-dimethylethyleneoxy,
ethylenemethyleneoxy, 1-ethylethyleneoxy, 2-ethylethyleneoxy, 1-
30 methyltrimethyleneoxy, 2-methyltrimethyleneoxy, 3-
methyltrimethyleneoxy and the like, with preference given to
ethyleneoxy. The alkyleneoxy group here means both -RO- and -
OR-, wherein R is C₁-C₄ alkylene. For example, ethyleneoxy
means both -CH₂CH₂O- and -OCH₂CH₂-.

The optionally substituted 4 to 7-membered heterocycle having 1 or 2 hetero atom(s) selected from the group consisting of nitrogen atom and oxygen atom in the ring at Q' is, for example, a group derived from 3,5-dihydroimidazole,
 5 imidazolidine, pyrrolidine, oxazolidine, oxetane, oxolane, oxane, perhydroazepine, imidazolidine, oxepane, azetidine and the like. Examples of these substituents include C₁-C₁₈ alkyl group (e.g., methyl, ethyl etc.), C₂-C₄ alkoxyalkyl group (e.g., 2-methoxyethyl etc.), optionally substituted aryl group (as
 10 defined above, e.g., phenyl etc.), optionally substituted aralkyl group (as defined above, e.g., benzyl etc.), oxo group, thioxo group and the like. Preferable examples of the heterocycle group include groups derived from 3,5-dihydro-2-methylimidazole, 3,5-dihydro-2,3-dimethylimidazole, 3,5-
 15 dihydro-2-methyl-3-phenylimidazole, 3,5-dihydro-3-ethyl-2-methylimidazole, 3-benzyl-3,5-dihydro-2-methylimidazole, 1,3-dimethylimidazolidine, pyrrolidine, 1-methylpyrrolidine, 1-(2-methoxyethyl)pyrrolidine, oxazolidine and 5,5-dimethyloxane.

The C₁-C₄ alkyl group at R is, for example, methyl,
 20 ethyl, propyl, isopropyl, butyl, isobutyl and the like, with preference given to methyl and ethyl.

Examples of the X include hydrogen atom, hydroxy, methoxy, ethoxy, isopropoxy, acetoxy and the like, with particular preference given to hydroxy.

25 As R¹, a group of the following formula is preferable:



wherein R⁵ is optionally substituted phenyl group or naphthyl group, Z is void and R⁶ is hydrogen atom.

Specific examples of R¹ include
 30 1-benzylpiperidin-4-ylamino,
 4-phenylcyclohexyl-1-ylamino,

4-hydroxy-4-(4-chlorophenyl)piperidino,
 4-hydroxy-4-(2-naphthyl)piperidino,
 4-hydroxy-4-(benzo(b)thiophen-2-yl)piperidino,
 4-benzylpiperidino,
 5 4-(4-fluorobenzyl)piperidino,
 4-(4-chlorobenzyl)piperidino,
 4-(4-bromobenzyl)piperidino,
 4-phenylpiperidino,
 4-(4-fluorophenyl)piperidino,
 10 4-(4-chlorophenyl)piperidino,
 4-(4-bromophenyl)piperidino,
 4-(4-methoxyphenyl)piperidino,
 4-(4-methylphenyl)piperidino,
 4-(4-trifluoromethylphenyl)piperidino,
 15 4-(3-chlorophenyl)piperidino,
 4-(3-fluorophenyl)piperidino,
 4-(3-trifluoromethylphenyl)piperidino,
 4-(3-bromophenyl)piperidino,
 4-(3-methoxyphenyl)piperidino,
 20 4-(3-methylphenyl)piperidino,
 4-(2-fluorophenyl)piperidino,
 4-(2-chlorophenyl)piperidino,
 4-(2-bromophenyl)piperidino,
 4-(2-methylphenyl)piperidino,
 25 4-(2-methoxyphenyl)piperidino,
 4-(3,4-dichlorophenyl)piperidino,
 4-(3,4-dimethylphenyl)piperidino,
 4-(3,4-dimethoxyphenyl)piperidino,
 4-(3,4-methylenedioxyphenyl)piperidino,
 30 4-(2,3-dimethoxyphenyl)piperidino,
 4-(2,3-dimethylphenyl)piperidino,
 4-(2,3-dichlorophenyl)piperidino,
 4-(3,5-dimethoxyphenyl)piperidino,
 4-(3,5-dimethylphenyl)piperidino,

4-(3,5-dichlorophenyl)piperidino,
 4-(2,6-dimethoxyphenyl)piperidino,
 4-(3,4,5-trimethoxyphenyl)piperidino,
 4-(naphthalen-1-yl)piperidino,
 5 4-(naphthalen-2-yl)piperidino,
 4-(6-methoxynaphthalen-2-yl)piperidino,
 4-(benzo(b)thiophen-2-yl)piperidino,
 4-(benzo(b)furan-2-yl)piperidino,
 4-(indol-2-yl)piperidino,
 10 4-(4-fluorobenzyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-chlorobenzyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-bromobenzyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-phenyl-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-fluorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 15 4-(4-chlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-bromophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-methoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-methylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-trifluoromethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 20 4-(3-chlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-fluorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-trifluoromethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-bromophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-methoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 25 4-(3-methylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-fluorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-chlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-bromophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-methylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 30 4-(2-methoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4-dichlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4-dimethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4-methylenedioxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,

- 4-(2,3-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2,3-dimethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2,3-dichlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,5-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 5 4-(3,5-dimethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,5-dichlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2,6-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4,5-trimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(naphthalen-1-yl)-3,6-dihydro-2H-pyridin-1-yl,
 10 4-(naphthalen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(6-methoxynaphthalen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(benzo(b)thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(benzo(b)furan-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(indol-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 15 4-(1,4-benzodioxan-6-yl)piperidino,
 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 4-(benzo(b)furan-5-yl)piperidino,
 4-(chroman-6-yl)piperidino,
 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 20 4-(2,3-dihydrobenzo(b)furan-6-yl)piperidino,
 4-(2,3-dihydrobenzo(b)thiophen-5-yl)piperidino,
 4-(2,3-dihydrobenzo(b)thiophen-6-yl)piperidino,
 4-(benzo(b)furan-5-yl)piperidino,
 4-(benzo(b)furan-6-yl)piperidino,
 25 4-(benzo(b)thiophen-5-yl)piperidino,
 4-(benzo(b)thiophen-6-yl)piperidino,
 4-(4-methoxy-3-methylphenyl)piperidino,
 4-(indan-5-yl)piperidino,
 4-(inden-5-yl)piperidino,
 30 4-(1H-indolin-5-yl)piperidino,
 4-(1-methylindolin-5-yl)piperidino,
 1,3-dihydrobenzo(c)furan-1-spiro-4'-piperidin-1'-yl,
 4-(chroman-7-yl)piperidino,
 4-(2H-chromen-6-yl)piperidino,

- 4-(3-chloro-4-methoxyphenyl)piperidino,
4-(4-chloro-3-methoxyphenyl)piperidino,
4-(3-chloro-4-methylphenyl)piperidino,
4-(4-chloro-3-methylphenyl)piperidino,
5 4-(3-chloro-4-fluorophenyl)piperidino,
4-(4-chloro-3-fluorophenyl)piperidino,
4-(3-chloro-4-trifluoromethylphenyl)piperidino,
4-(4-chloro-3-trifluoromethylphenyl)piperidino,
4-(1H-indol-6-yl)piperidino,
10 4-(1-methylindol-6-yl)piperidino,
4-(1,3-dihydrobenzo(c) furan-5-yl)piperidino,
4-(3,4-dihydro-1H-benzo(c) oxin-6-yl)piperidino,
4-(3,4-dihydro-2H-benzo(b) thiin-6-yl)piperidino,
4-(3,4-dihydro-2H-benzo(b) thiin-7-yl)piperidino,
15 4-(2-methyl-2,3-dihydrobenzo(b) furan-5-yl)piperidino,
4-(2,2-dimethyl-2,3-dihydrobenzo(b) furan-5-yl)piperidino,
4-(1-methyl-2-oxoindolin-5-yl)piperidino,
4-(4-chloro-2,2-dimethyl-2,3-dihydrobenzo(b) furan-5-yl)piperidino,
20 4-(7-chloro-2,2-dimethyl-2,3-dihydrobenzo(b) furan-5-yl)piperidino,
4-(2,2-dimethyl-4-methyl-2,3-dihydrobenzo(b) furan-5-yl)piperidino,
4-(2,2-dimethyl-7-methyl-2,3-dihydrobenzo(b) furan-5-yl)piperidino,
25 4-(2,4,6-trimethylphenyl)piperidino,
4-(2H-1-oxoisoindolin-5-yl)piperidino,
4-(2-methyl-1-oxoisoindolin-5-yl)piperidino,
4-(quinolin-6-yl)piperidino,
30 4-(isoquinolin-6-yl)piperidino,
4-(4,5-dimethylthiophen-2-yl)piperidino,
4-(4,5-dichlorothiophen-2-yl)piperidino,
4-(4,5-dimethylfuran-2-yl)piperidino,
4-(4,5-dichlorofuran-2-yl)piperidino,

4-(2-methylpyridin-4-yl)piperidino
and the like.

As R¹, preferred are

- 1-benzylpiperidin-4-ylamino,
5 4-phenylcyclohexyl-1-ylamino,
4-hydroxy-4-(4-chlorophenyl)piperidino,
4-hydroxy-4-(2-naphthyl)piperidino,
4-hydroxy-4-(benzo (b) thiophen-2-yl)piperidino,
4-benzylpiperidino,
10 4-(4-fluorobenzyl)piperidino,
4-(4-chlorobenzyl)piperidino,
4-(4-bromobenzyl)piperidino,
4-phenylpiperidino,
4-(4-fluorophenyl)piperidino,
15 4-(4-chlorophenyl)piperidino,
4-(4-bromophenyl)piperidino,
4-(4-methoxyphenyl)piperidino,
4-(4-methylphenyl)piperidino,
4-(4-trifluoromethylphenyl)piperidino,
20 4-(3-chlorophenyl)piperidino,
4-(3-fluorophenyl)piperidino,
4-(3-trifluoromethylphenyl)piperidino,
4-(3-bromophenyl)piperidino,
4-(3-methoxyphenyl)piperidino,
25 4-(3-methylphenyl)piperidino,
4-(2-fluorophenyl)piperidino,
4-(2-chlorophenyl)piperidino,
4-(2-bromophenyl)piperidino,
4-(2-methylphenyl)piperidino,
30 4-(2-methoxyphenyl)piperidino,
4-(3,4-dichlorophenyl)piperidino,
4-(3,4-dimethylphenyl)piperidino,
4-(3,4-dimethoxyphenyl)piperidino,
4-(3,4-methylenedioxyphenyl)piperidino,

- 4-(2,3-dimethoxyphenyl)piperidino,
 4-(2,3-dimethylphenyl)piperidino,
 4-(2,3-dichlorophenyl)piperidino,
 4-(3,5-dimethoxyphenyl)piperidino,
 5 4-(3,5-dimethylphenyl)piperidino,
 4-(3,5-dichlorophenyl)piperidino,
 4-(2,6-dimethoxyphenyl)piperidino,
 4-(3,4,5-trimethoxyphenyl)piperidino,
 4-(naphthalen-1-yl)piperidino,
 10 4-(naphthalen-2-yl)piperidino,
 4-(6-methoxynaphthalen-2-yl)piperidino,
 4-(benzo(b)thiophen-2-yl)piperidino,
 4-(benzo(b)furan-2-yl)piperidino,
 4-(indol-2-yl)piperidino,
 15 4-(4-fluorobenzyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-chlorobenzyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-bromobenzyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-phenyl-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-fluorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 20 4-(4-chlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-bromophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-methoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-methylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(4-trifluoromethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 25 4-(3-chlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-fluorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-trifluoromethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-bromophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3-methoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 30 4-(3-methylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-fluorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-chlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-bromophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2-methylphenyl)-3,6-dihydro-2H-pyridin-1-yl,

4-(2-methoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4-dichlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4-dimethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 5 4-(3,4-methylenedioxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2,3-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2,3-dimethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2,3-dichlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,5-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 10 4-(3,5-dimethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,5-dichlorophenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(2,6-dimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(3,4,5-trimethoxyphenyl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(naphthalen-1-yl)-3,6-dihydro-2H-pyridin-1-yl,
 15 4-(naphthalen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(6-methoxynaphthalen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(benzo(b)thiophen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(benzo(b)furan-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 4-(indol-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
 20 4-(1,4-benzodioxan-6-yl)piperidino,
 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 4-(benzo(b)furan-5-yl)piperidino,
 4-(chroman-6-yl)piperidino,
 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 25 4-(benzo(b)furan-5-yl)piperidino,
 4-(2,2-dimethyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 4-(7-chloro-2,2-dimethyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 4-(2,2-dimethyl-4-methyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 30 4-(2,2-dimethyl-7-methyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino,
 4-(2,4,6-trimethylphenyl)piperidino,
 4-(2H-1-oxoisindolin-5-yl)piperidino,

4-(2-methyl-1-oxoisoindolin-5-yl)piperidino,
4-(quinolin-6-yl)piperidino,
4-(isoquinolin-6-yl)piperidino,
4-(4,5-dimethylthiophen-2-yl)piperidino,
5 4-(4,5-dichlorothiophen-2-yl)piperidino,
4-(4,5-dimethylfuran-2-yl)piperidino,
4-(4,5-dichlorofuran-2-yl)piperidino,
4-(2-methylpyridin-4-yl)piperidino
and the like.

10 As R¹, more preferred are

4-(3,4-dimethylphenyl)piperidin-1-yl,
4-(1-naphthyl)piperidin-1-yl,
4-(2-naphthyl)piperidin-1-yl,
4-(6-methoxynaphthalen-2-yl)piperidin-1-yl,
15 4-(3,4-dimethylphenyl)-3,6-dihydro-2H-pyridin-1-yl,
4-(1-naphthyl)-3,6-dihydro-2H-pyridin-1-yl,
4-(2-naphthyl)-3,6-dihydro-2H-pyridin-1-yl,
4-(6-methoxynaphthalen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
4-(benzo(b)furan-5-yl)piperidino,
20 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino,
4-(2,2-dimethyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino,
4-(4-chloro-3-fluorophenyl)piperidino,
4-(4-chloro-3-methylphenyl)piperidino,
4-(3,4-dichlorophenyl)piperidino
25 and the like.


As R¹,

4-(naphthalen-1-yl)piperidino,
4-(naphthalen-2-yl)piperidino,
4-(naphthalen-1-yl)-3,6-dihydro-2H-pyridin-1-yl,
30 4-(naphthalen-2-yl)-3,6-dihydro-2H-pyridin-1-yl,
4-(3,4-dichlorophenyl)piperidino,
4-(4-chloro-3-methylphenyl)piperidino,
4-(2,2-dimethyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino
and the like are particularly preferable.

As R³, hydrogen atom and C₁-C₄ alkyl (e.g., methyl, ethyl, propyl, isopropyl, butyl etc.) are preferable and hydrogen atom is particularly preferable.

W is preferably void.


5 As R⁷, a group of the following formula is preferable:



wherein

R⁸ is hydrogen atom, phenyl group, C₁-C₄ alkyl group, C₁-C₂ halogenated alkyl group, halogen atom, C₂-C₄ alkenyl group, C₁-C₄ hydroxyalkyl group, alkoxyalkyl group, alkyloxycarbonyl group, optionally substituted amino group, acetamido group, carboxyl group, acyl group, optionally substituted alkyloxy group, alkylthio group or cyano group, and

15 R⁹ is a group of the following formula




wherein R¹⁰ and R¹¹ are each independently hydrogen atom, C₁-C₁₈ alkyl group, optionally substituted aryl group, optionally substituted aralkyl group or alkoxy group, and R¹² is hydrogen atom, optionally substituted aryl group, C₁-C₁₈ alkyl group, C₁-C₈ alkoxy group or acyl group.

Ra, Rb and Rc are each specifically hydrogen atom,

fluorine atom, chlorine atom, bromine atom, methyl, ethyl, methoxy, methylenedioxy, hydroxy, acetyl and the like, with preference given to all Ra, Rb, Rc being hydrogen atom.


Examples of the group of the following formula




5

include groups derived from β -propiolactone, γ -butyrolactone, 5,5,-dimethyl- γ -butyrolactone, γ -valerolactone, δ -valerolactone, 6,6-dimethyl- δ -valerolactone, γ -caprolactone, ϵ -caprolactone, 6,6-dimethyl- ϵ -caprolactone, 2-azetidinone, 2-pyrrolidinone, δ -
10 valerolactam, ϵ -caprolactam, hydantoin, 3,5-dihydroimidazol-4-one and the like.

A preferable group of the following formula




includes the groups of the following formulas




15

, which are specifically groups derived from γ -butyrolactone, δ -valerolactone, 2-pyrrolidinone and the like, with particular preference given to a group of the following formula




20 Specific examples are groups derived from γ -butyrolactone and δ -valerolactone.

Preferable embodiment of the formula (I) includes the compounds of the following formulas:



The phenoxypropylamine compound of the present invention is a compound of the following formula (I)




5 wherein each symbol is as defined above.

The preferable compound of the above-mentioned formula (I) is a compound (compound A) wherein each symbol of the formula (I) is as follows:

a bond represented by a solid line and a dotted line shows a
10 double bond;

X is a hydrogen atom, a hydroxy group, a C₁-C₈ alkoxy group, an acyloxy group or an oxo group;
R¹ is a group of the following formula



wherein

Y is O or S,

m and n are each independently 0, 1 or 2,

5 Ar is optionally substituted benzene or naphthalene,

R² is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

R⁵ is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,

10 Z is void or -CH₂-, and

R⁶ is hydrogen atom, hydroxy group, acetamido group, carboxyl group, alkoxy carbonyl group, cyano group or C₁-C₈ alkoxy group;

R³ is a hydrogen atom, a C₁-C₁₈ alkyl group or a halogen atom;

15

V is -CH₂-, -O-, -S- or the formula -N(R⁴)-

wherein R⁴ is hydrogen atom, C₁-C₁₈ alkyl group or optionally substituted aralkyl group;

W is void or -CH₂- or -C(=O)-; or

20 V and W are each a hydrogen atom without bonding;

R⁷ is a C₁-C₄ hydroxyalkyl group, an acyl group, an optionally substituted saturated or unsaturated heterocyclic group, an optionally substituted fused heterocyclic group, a C₁-C₄ alkylsulfonyl group or the


25

formula -Q-R⁹

wherein

Q is -C(=O)-, -C(=S)-, -CH₂- or S(=O)₂-, and

R⁹ is a group of the following formula



or -NH-NH-R^{15}

wherein R^{10} and R^{11} are each independently hydrogen atom, $\text{C}_1\text{-C}_{18}$ alkyl group, optionally substituted aryl group, optionally substituted aralkyl group or alkoxy group, R^{12} is hydrogen atom, optionally substituted aryl group, $\text{C}_1\text{-C}_{18}$ alkyl group, $\text{C}_1\text{-C}_8$ alkoxy group or acyl group, and R^{15} is hydrogen atom, phenyl group, $\text{C}_1\text{-C}_4$ alkyl group, $\text{C}_1\text{-C}_2$ halogenated alkyl group, halogen atom, $\text{C}_2\text{-C}_4$ alkenyl group, $\text{C}_1\text{-C}_4$ hydroxyalkyl group, alkoxyalkyl group, alkyloxycarbonyl group, optionally substituted amino group, acetamido group, carboxyl group, acyl group, optionally substituted alkyloxy group, alkylthio group or cyano group; and

Ra , Rb and Rc are each independently a hydrogen atom, a $\text{C}_1\text{-C}_{18}$ alkyl group, a hydroxy group, a $\text{C}_1\text{-C}_8$ alkoxy group, a halogen atom, an acyl group, a nitro group or an amino group;


provided that when V and W are not directly bonded and V and W are both hydrogen atoms, R^7 should not be a group of the formula -CO-R^9 wherein R^9 is as defined above.

In the above-mentioned compound A, a compound A wherein each symbol of the formula (I) is as follows is more preferable:

a bond represented by a solid line or a dotted line shows a double bond;

X is a hydroxy group;

R¹ is a group of the following formula



wherein

R⁵ is optionally substituted phenyl group or naphthyl group,

Z is void, and


R⁶ is hydrogen atom;

R³ is a hydrogen atom or C₁-C₄ alkyl group;

V is -CH₂-, -O-, -S- or the formula -N(R⁴)- wherein R⁴ is hydrogen atom, C₁-C₆ alkyl group or optionally substituted aralkyl group;

W is void; or

R⁷ is a group of the following formula



or the formula -CO-R⁹


wherein

R⁸ is hydrogen atom, phenyl group, C₁-C₄ alkyl group, C₁-C₂ halogenated alkyl group, halogen atom, C₂-C₄ alkenyl group, C₁-C₄ hydroxyalkyl group,

alkoxyalkyl group, alkyloxycarbonyl group,

optionally substituted amino group, acetamido group,


carboxyl group, acyl group, optionally substituted
alkyloxy group, alkylthio group or cyano group, and
R⁹ is a group of the following formula



5 wherein R¹⁰ and R¹¹ are each independently hydrogen
atom, C₁-C₁₈ alkyl group, optionally substituted aryl
group, optionally substituted aralkyl group or
alkoxy group, and R¹² is hydrogen atom, optionally
substituted aryl group, C₁-C₁₈ alkyl group, C₁-C₈
10 alkoxy group or acyl group; and

Ra, Rb and Rc are each a hydrogen atom.

In the above-mentioned compound A, that having the
following formula (I') is more preferable:




15 wherein each symbol is as defined for compound A.

Specific examples of the above-mentioned compound A are:

- (1) 1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-
propyloxy)benzo(b)furan-2-ylcarbonyl)pyrrolidine,
- (2) 4-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-
20 propyloxy)benzo(b)furan-2-ylcarbonyl)morpholine,
- (4) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-
N,N-dimethylbenzo(b)furan-2-carboxamide,

- (12) 1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)thiophen-2-ylcarbonyl)pyrrolidine,
(13) 4-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)thiophen-2-ylcarbonyl)morpholine,
5 (15) 4-(2-hydroxy-3-(4-(naphthalen-1-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide,
(17) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide,
(20) 4-(7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-ylcarbonyl)morpholine,
10 (21) 7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)furan-2-carboxamide,
(27) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethyl-1H-indole-2-carboxamide,
15 (30) 4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethyl-1-methylindole-2-carboxamide,
(35) 1-(2-(5-methyl-1,2,4-oxadiazol-3-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
(37) 1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
20 (38) 1-(2-(5-trifluoromethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
(39) 1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
25 (42) 1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indol-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
(44) 1-(2-(3-methyl-1,2,4-oxadiazol-5-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
(48) 1-(2-(5-methyloxazol-2-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol,
30 (81) 3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol,
(88) 1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol, and



wherein

E is an oxygen atom or a sulfur atom;

Q' is optionally substituted 4 to 7-membered

5 . heterocycle having 1 or 2 hetero atom(s) selected from the group consisting of nitrogen atom and oxygen atom in the ring,


and V is hydrogen atom; and

Ra, Rb and Rc are each independently a hydrogen atom, a C₁-C₁₈


10 alkyl group, a hydroxy group, a C₁-C₈ alkoxy group, a halogen atom, an acyl group, a nitro group or an amino group.

In the above-mentioned compound B, that wherein each symbol is as follows is more preferable:

15 a group of the following formula



is a group of the following formula



wherein

20 E is an oxygen atom or a sulfur atom,

q is 0, 1, 2 or 3,

R^{4'}, R^{7'} and R^{8'} are each independently a hydrogen atom, a C₁-C₁₈ alkyl group, an optionally substituted aryl group or an optionally substituted aralkyl group, and

25 other symbols are as defined in the aforementioned compound B.


In the above-mentioned compound B, that wherein each

symbol is as follows is more preferable:

a bond represented by a solid line and a dotted line shows a double bond;

X is a hydroxy group;

5 R¹ is a group of the following formula:



wherein


R⁵ is optionally substituted phenyl group or naphthyl group,

10 Z is void, and


R⁶ is hydrogen atom;

R³ is a hydrogen atom or C₁-C₄ alkyl group;

a group of the formula




15 is a group of the following formula



wherein q is 1 and R^{4'} is hydrogen atom or C₁-C₄ alkyl group); and

R_a, R_b and R_c are each a hydrogen atom.

20 In the above-mentioned compound B, that having the following formula (I'') is particularly preferable:




wherein each symbol is as defined for compound B.

Specific examples of the above-mentioned compound B are as follows:

- 5 (306) 5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-1,3-dimethylimidazolidine-2,4-dione,
- (307) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone,
- (308) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-
- 10 propyloxy)benzylidene)- γ -butyrolactone,
- (309) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone,
- (310) α -(2'-(3-(4-(3-fluoro-4-methylphenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,
- 15 (311) α -(2'-(3-(4-(3,4-dimethylphenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,
- (312) α -(2'-(3-(4-(4-chloro-3-fluorophenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,
- (313) α -(2'-(3-(4-(4-chloro-3-trifluoromethylphenyl)-
- 20 piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone,
- (314) α -(2'-(2-hydroxy-3-(4-(naphthalen-1-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone,
- (315) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- δ -valerolactone,
- 25 (316) α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -valerolactone,
- (319) 3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-2-pyrrolidone,
- (322) 3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-


propyloxy)benzylidene)-1-methyl-2-pyrrolidone, and
 (325) α -(2'-(2-hydroxy-3-(4-(6-methoxynaphthalen-2-yl)piperidino)propyloxy)benzylidene)- γ -butyrolactone, wherein
 the number in the parenthesis affixed to each compound is an
 5 Example number.

A synthetic intermediate of compound A may be a compound
 of the following formula (II)



wherein each symbol in the formula is as defined below:

- 10 X is a hydrogen atom, a hydroxy group, a C₁-C₈ alkoxy group or an acyloxy group or an oxo group;
 R¹ is a group of the following formula



- 15 wherein
 Y is O or S,
 m and n are each independently 0, 1 or 2,
 Ar is optionally substituted benzene or naphthalene,
 R² is optionally substituted aryl group or optionally
 20 substituted aromatic heterocyclic group,
 R⁵ is optionally substituted aryl group or optionally substituted aromatic heterocyclic group,
 Z is void or -CH₂-, and
 R⁶ is hydrogen atom, hydroxy group, acetamido group,

carboxyl group, alkoxycarbonyl group, cyano group
or C₁-C₈ alkoxy group;

R³ is a hydrogen atom, a C₁-C₁₈ alkyl group or a halogen
atom;

5 V is -CH₂-, -O-, -S- or the formula -N(R⁴)-
wherein

R⁴ is hydrogen atom, C₁-C₁₈ alkyl group or optionally
substituted aralkyl group;


W is void, -CH₂- or -C(=O)-; or

10 V and W are each a hydrogen atom without bonding;

R¹⁴ is a hydrogen atom or a C₁-C₄ alkyl; and

Ra, Rb and Rc are each independently a hydrogen atom, a C₁-C₁₈
alkyl group, a hydroxy group, a C₁-C₈ alkoxy group,
a halogen atom, an acyl group, a nitro group or an
15 amino group.

A synthetic intermediate of compound B may be a compound
of the following formula (III)



wherein each symbol in the formula is as defined below:

20 R is a hydrogen atom, a C₁-C₄ alkyl group, an allyl group
or a 2,3-epoxypropan-1-yl group;

a bond represented by a solid line and a dotted line shows a
double bond or a single bond;

E is an oxygen atom or a sulfur atom;


25 R³ is a hydrogen atom, a C₁-C₁₈ alkyl group or a halogen
atom;

Q' is an optionally substituted 4 to 7-membered
heterocycle having 1 or 2 hetero atom(s) selected from
the group consisting of nitrogen atom and oxygen atom


in the ring; and

Ra, Rb and Rc are each independently a hydrogen atom, a C₁-C₁₈ alkyl group, a hydroxy group, a C₁-C₈ alkoxy group, a halogen atom, acyl group, a nitro group or an amino group.

As a synthetic intermediate for the above-mentioned compound B, a compound of the formula (III) wherein each symbol is as defined below is preferable:
the group of the following formula



is a group of the following formula



wherein

E is an oxygen atom or a sulfur atom,

q is 0, 1, 2 or 3, and

R₄', R₇' and R₈' are each independently hydrogen atom, C₁-C₁₈ alkyl group, optionally substituted aryl group or optionally substituted aralkyl group, and

other symbols are as defined with regard to the formula

(III).

The pharmaceutically acceptable salts of compounds of the formulas (I), (II) and (III) include acid addition salts with inorganic acids (e.g., hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid etc.) or organic acids (e.g., acetic acid, propionic acid, succinic acid, glycolic acid, lactic acid, malic acid, tartaric acid, citric acid, maleic acid, fumaric acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, camphorsulfonic


acid, ascorbic acid, terephthalic acid, adipic acid etc.).

The compounds of the formulas (I), (II) and (III) and pharmaceutically acceptable salts thereof may be present in the form of a hydrate or a solvate. These hydrates and solvates
5 are also encompassed in the present invention. When the compound of the formula (I) has an asymmetric atom, at least two kinds of optical isomers exist. The optical isomers and racemates thereof are encompassed in the present invention.


The compound of the formula (I) can be synthesized by
10 the following methods. Each symbol in the following reaction formulas is as defined above, unless particularly specified.

The compound of the formula (I) and synthetic intermediates of the formulas (II) and (III) can be produced according to the following reaction formulas A - Z and Q' - T',
15 as well as methods analogous to the following examples and the like. In the formulas, the symbol A refers to a leaving group (or nucleofugal group) well known in the organic synthesis, such as chlorine atom, bromine atom, iodine atom, mesylate, tosylate, nosylate, triflate and the like. Leaving groups (or
20 nucleofugal groups) are well known to those of ordinary skill in the art of organic syntheses.


Reaction Formula A




Reaction Formula B




Reaction Formula C



Reaction Formula D



Reaction Formula E



Reaction formula A: A method comprising reacting a phenol derivative 1 and 2,3-epoxypropane compound 2 having a leaving group (or nucleofugal group) at 1-position, followed by

reaction with $H-R^1$,

Reaction formula B: A method comprising reacting $H-R^1$ and 2,3-epoxypropane compound 2 having a leaving group (or nucleofugal group) at 1-position to give compound 4, which is reacted with
5 a phenol derivative 1,


Reaction formula C: A method comprising reacting a phenol derivative 1 and 2-propanone 5 having leaving groups (or nucleofugal groups) at 1,3-position to give compound 6, which is reacted with $H-R^1$ to give a product 7, followed by reduction
10 thereof,

Reaction formula D: A method comprising reacting $H-R^1$ and 2-propanone compound 5 having leaving groups (or nucleofugal groups) at 1,3-position to give compound 8, which is reacted with phenol derivative 1 to give a product 7, followed by
15 reduction thereof,


Reaction formula E: A method comprising reacting phenol derivative 1 and allyl compound 9 (e.g., 3-allyl bromide etc.) having a leaving group (or nucleofugal group) at 3-position to give a product 10, which is epoxidated and successively reacted
20 with $H-R^1$, and the like. The methods for synthesis of the compound of the formula (I) are not limited to those mentioned above.

Particularly, the optically active compound of the formula (I) ($X=OH$) can be synthesized by the following reaction
25 formulas F - L and the like. In these reaction formulas, the symbol R^* means a part other than carboxyl group of optically active carboxylic acid.


Reaction Formula F




Reaction Formula G




Reaction Formula H




Reaction Formula I




Reaction Formula J



Reaction Formula K



Reaction Formula L



Reaction formula F: A method comprising asymmetric epoxidation of intermediate 10 obtained by the above-mentioned reaction formula E, using optically active base and asymmetric ligand in

1 catalytic or stoichiometric amounts to give optically active
intermediate 3, which is reacted with $H-R^1$,

Reaction formula G: A method comprising reacting phenol
derivative 1 and optically active 2,3-epoxypropane derivative 2
5 having a leaving group (or nucleofugal group) at 1-position to
give compound 3, which is reacted with $H-R^1$,

Reaction formula H: A method comprising reacting $H-R^1$ and
optically active 2,3-epoxypropane derivative 2 having a leaving
group (or nucleofugal group) at 1-position to give compound 4,
10 which is reacted with phenol derivative 1,

Reaction formula I: A method comprising condensing a racemic
mixture of the compound of the formula (I) with optically
active carboxylic acid 11 to convert the compound to optically
active ester 12, which is followed by crystallization, column
15 chromatography and the like to resolve the compound into two
diastereomers,

Reaction formula J: A method comprising asymmetric reduction of
intermediate 7 obtained by the above-mentioned reaction
formulas C and D, using a chiral ligand,


20 Reaction formula K: A method comprising forming a salt in a
racemic mixture of the compound of the formula (I) and
optically active carboxylic acid 11, whereby both isomers are
resolved based on difference in crystallinity,

Reaction formula L: A method comprising condensing a racemic
25 mixture of the compound of the formula (I) with carboxylic acid
13 to once convert the compound to an ester, and hydrolyzing
the ester enantioselectively using an enzyme.


The methods for synthesizing the optically active
compound of the formula (I) ($X=OH$) are not limited to those
30 mentioned above.

A compound of the formula (I) wherein X is hydrogen atom
can be synthesized as in the following reaction formulas M - N
and the like.

Reaction Formula M



Reaction Formula N




Reaction formula M: A method comprising reacting phenol derivative 1 and propane derivative 15 having leaving groups or nucleofugal groups at 1,3-positions to synthesize intermediate 16, and condensing the intermediate 16 and H-R¹ in the presence of deoxidizing agent,

Reaction formula N: A method comprising reacting H-R¹ and propane derivative 15 having leaving groups or nucleofugal groups at 1,3-positions to synthesize intermediate 17 and condensing the intermediate 17 and phenol derivative 1 in the presence of deoxidizing agent.

Of the compounds of the formula (I), a compound wherein X is alkoxy can be derived from the compound of the formula (I) wherein X is OH as in the following reaction formula O wherein R¹³ is alkyl group.


Reaction Formula O



Reaction formula O: A method comprising alkylating hydroxy group of a compound of the formula (I) wherein X is hydroxy group, in the presence of deoxidizing agent.

- 5 Of the compounds of the formula (I), a compound wherein R^7 is the formula: $-Q-R^9$ wherein Q is $-C(=O)-$ or $-CH_2-$ can be derived from carboxylic acid derivative 18, as in the following the reaction formula P.

Reaction Formula P




- 10 Reaction formula P: A method comprising condensing carboxylic acid derivative 18 with $H-R^9$ in the presence of amidating agent to synthesize amide compound ($Q=CO$), and reducing the amide compound to synthesize amino compound ($Q=CH_2$). The amidating agent to be used for this method is exemplified by
- 15 dicyclohexylcarbodiimide (DCC), diethyl cyanophosphate, diphenylphosphoryl azide (DPPA), 1,1'-carbonylbis-1H-imidazole (CDI), 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride (WSC) and the like. The reducing agent to be used for the reduction is exemplified by lithium aluminum


hydride, diisopropyl aluminum hydride, diborane, sodium borohydride and the like.

Of the phenol derivatives 1 used in Reaction formulas A, B, D, E, G, H, M and N, a compound wherein R⁷ is the formula:
 5 -Q-R⁹ can be synthesized according to the following reaction formulas Q - S and the like. In these reaction formulas, the symbol PG means a protecting group (e.g., methyl, ethyl, methoxymethyl, ethoxymethyl, trimethylsilyl, benzyl, acetyl, benzoyl etc.) that can be eliminated easily in the organic
 10 synthesis.


Reaction Formula Q



Reaction Formula R



Reaction Formula S



Reaction formula Q: A production method comprising condensing carboxylic acid derivative 19 with $H-R^9$, using amidating agent, and then eliminating the protecting group to give phenol derivative (1, $Q=CO$). As the amidating agent,


5 dicyclohexylcarbodiimide (DCC), diethyl cyanophosphate, diphenylphosphoryl azide (DPPA), 1,1'-carbonylbis-1H-imidazole (CDI), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC) and the like can be used.

Reaction formula R: A production method comprising reducing
10 amide compound 20 with a reducing agent, and eliminating the protecting group to give phenol derivative (1, $Q=CH_2$). As the reducing agent, lithium aluminum hydride, diisopropyl aluminum hydride, diborane, sodium borohydride and the like can be used.


Reaction formula S: A production method comprising condensing,
15 sulfonic chloride derivative 22 with $H-R^9$ using a deoxidizing agent, and eliminating the protecting group to give phenol derivative (1, $Q=SO_2$).

In the following, the reaction formulas T - Z are shown as typical synthetic methods of representative compounds
20 wherein R^7 is optionally substituted heterocycle in the reaction formulas A - H, M and N. In these reaction formulas, the symbol PG is as defined above.


Reaction Formula T



Reaction Formula U



Reaction Formula V



Reaction formula T: A production method of phenol derivative 1 having a 1,3,4-oxadiazole ring, which method comprises

5 cyclization of diacylhydrazine derivative 24 with a dehydrating agent, and deprotection. The phenol derivative 1 having a 1,3,4-oxadiazole ring can be also synthesized by reacting azo compound and triphenylphosphine, in the presence of a

deoxidizing agent, followed by deprotection. As the

10 dehydrating agent, polyphosphoric acid, oxalyl chloride, phosphorus trichloride, sulfuric acid, phosphorus oxychloride, thionyl chloride, oxalyl chloride and the like can be used. As

the azo compound, diethyl azodicarboxylate (DEAD), diisopropyl azodicarboxylate (DIAD) and the like can be used.


Reaction formula U: A production method of phenol derivative 1 having a 1,2,4-oxadiazole ring, which method comprises

- 5 condensing carboxylic acid derivative 19 and hydroxyimino compound 26 using an amidating agent to give compound 27, which is subjected to cyclization using a dehydrating agent or by heating for dehydration, followed by deprotection. As the dehydrating agent, polyphosphoric acid, phosphorus
- 10 pentachloride, phosphorus trichloride, sulfuric acid, phosphorus oxychloride, thionyl chloride, oxalyl chloride and the like can be used.


Reaction formula V: A production method of phenol derivative 1 having a 1,2,4-oxadiazole ring, which method comprises

- 15 condensing nitrile derivative 29 and hydroxylamine to give compound 30, to which acid anhydride 31 is added to allow cyclization by heating for dehydration, followed by deprotection.

Reaction Formula W



Reaction Formula X




Reaction formula W: A production method of phenol derivative 1 having a 1,3,4-thiadiazole ring, which method comprises

- 5 conversion of hydrazone compound 33 into thione compound with a sulfidation agent to give compound 34, which is cyclized with compound 35 by heating, followed by deprotection. As the sulfidation agent, Lawesson reagent, diphosphorus pentasulfide and the like can be used.


- 10 Reaction formula X: A production method of phenol derivative 1 having a thiazole ring, which method comprises cyclization of

compound 37 and thioamide compound 38 by heating, followed by deprotection.

Reaction Formula Y



Reaction Formula Z




5 Reaction formula Y: A production method of phenol derivative 1 having a isoxazole ring, which method comprises cyclization of

hydroxyimino compound 40, using a dehydrating agent or by heating for dehydration, followed by deprotection. As the dehydrating agent, polyphosphoric acid, phosphorus pentachloride, phosphorus trichloride, sulfuric acid, 5 phosphorus oxychloride, thionyl chloride, oxalyl chloride and the like can be used.

Reaction formula Z: A production method of phenol derivative 1 having an oxazole ring, which method comprises condensing acid halide compound 42 and acetylene compound 43 to give compound 10 44, followed by cyclization using mercury(II) acetate and deprotection.

Of the phenol derivatives 1 to be used for the reaction formulas A, B, D, E, G, H, M and N, a compound wherein R and W are bonded to form a ring can be also synthesized by the 15 methods of the following reaction formulas Q' - T'.


Reaction Formula Q'



Reaction formula Q': The phenol derivative 1, which is a benzylidene compound, can be synthesized by reacting phenol derivative 18 and carbonyl (thiocarbonyl) derivative 19 in the 20 presence of a base such as sodium hydride, sodium alcoholate and the like. Further, the phenol derivative 1, which is a benzyl compound, can be synthesized by reducing the obtained phenol derivative 1 (benzylidene compound) in the presence of a catalyst such as palladium carbon and the like.

25 When R and W are bonded to form a ring, which is a lactam or hydantoin (or their sulfur derivatives), and R' is other than hydrogen atom, phenol derivative 1 can be also synthesized by the method of the reaction formula R'.


Reaction Formula R'



Reaction formula R': Synthesis is available by lithiation of a lactam derivative or hydantoin derivative (or their sulfur derivatives) wherein $R^{7'}$ is other than hydrogen atom, with an organic lithium reagent such as $n\text{-BuLi}$ and the like, reacting the same with a phenol derivative 20, wherein hydroxyl group is protected, to once convert to benzyl alcohol compound 21, followed by treatment with an acid. Moreover, by reduction of phenol derivative 1 (benzylidene compound) in the presence of a catalyst such as palladium carbon and the like, phenol derivative 1, which is a benzyl compound, can be synthesized.

The phenol derivative 1 (or its sulfur derivative) containing hydantoin, wherein $R^{7'}$ and $R^{8'}$ are the same substituents, can be also synthesized by the method of the reaction formula S'.


Reaction Formula S'



Reaction formula S': The phenol derivative 20, wherein hydroxyl group is protected, is reacted with hydantoin (or its sulfur derivative) along with a base such as sodium hydride, sodium alcoholate and the like to give benzylidene compound 22, followed by deprotection to give phenol derivative 1, wherein R^{7'} and R^{8'} are hydrogen atoms. Moreover, by reacting benzylidene compound 22, which is an intermediate, with W-R^{7'} having a nucleofugal group to synthesize intermediate 23, and deprotecting the same, the phenol derivative 1 (or its sulfur derivative) containing hydantoin, wherein R^{7'} and R^{8'} are the same, can be synthesized.

The phenol derivative 1 (or its sulfur derivative) containing 3,5-dihydroimidazol-4-one can be also synthesized by the method of the reaction formula T'.

Reaction Formula T'



Reaction formula T': The phenol derivative 20, wherein hydroxyl group is protected, is reacted with glycine derivative 24 to give benzylidene compound 25, which is then reacted with amine $R^{7'}-NH_2$ to give intermediate 26. This intermediate is deprotected to give phenol derivative 1 (or its sulfur derivative) containing 3,5-dihydroimidazol-4-one.

There are many methods for obtaining phenol derivative 1 other than those mentioned above that are known to synthesis chemists, and therefore, the methods for obtaining the compound are not limited to those shown above.

These reactions and applications ultimately leading to the formula (I) of the present invention are well known to those of ordinary skill in the field of organic chemical synthesis. The improvements to adopt the conditions and reagents for the synthesis of specific compounds of the formula (I) inclusive of the inventive compound, beyond those described, are known to synthesis chemists. For more detailed description, respective synthetic examples are shown under Examples.

The compounds of the formula (I) obtained as mentioned above have selective affinity for as well as simultaneous antagonistic activity against 5-HT_{1A} receptors and have a 5-HT reuptake inhibitory action. Therefore, the compounds can

provide effective pharmaceutical agents for diseases accompanying serotonergic neurotransmission functional disorders. They are also effective as 5HT_{1A} antagonists having a selective serotonin reuptake inhibitory action, or as
5 selective serotonin reuptake inhibitors having a 5HT_{1A} antagonistic action.

That is, the inventive compounds show quick expression of the anti-depressive effect and are useful as a so-called rapid onset antidepressant. They are also useful for the
10 treatment of mammals inclusive of human for central nervous system diseases mediated by 5-HT, such as schizophrenia, anxiety neurosis, obsessive-compulsive disorder (OCD), panic disorder, social anxiety disorder, seasonal emotional disorder, Anorexia Nervosa, Bulimia Nervosa, nocturnal enuresis,
15 children's hyperlocomotion, post-traumatic stress disorder (PTSD), senile dementia, hemicrania, stroke, Alzheimer's disease, recognition disorder, hypertension, gastrointestinal injury, feeding disorders, premenstrual syndrome (PMS), abnormal body temperature regulation and sexual disorder, pain,
20 as well as abnormal cardiovascular system, drug abuse and the like.

When the compound of the present invention is used as a pharmaceutical agent, a systemic administration of a pharmacologically acceptable amount of the compound of the
25 formula (I) or a pharmacologically acceptable acid addition salt thereof to a mammal is included. The dose requires careful control for each case, and in consideration of the age, body weight and condition of the subject, administration route, as well as nature and severity of disease, the general daily
30 dose in the case of parenteral administration is 0.01 - 100 mg/kg, preferably 0.1 - 1 mg/kg, and that in the case of oral administration is 0.5 - 10 mg/kg, preferably 1 - 5 mg/kg. The administration method in the present invention includes oral, rectal and parenteral (e.g., intramuscular, intravenous,

percutaneous and subcutaneous) administrations.

For anti-depression, the compound of the present invention may be administered as a single therapeutic agent or may be administered as a mixture with other therapeutic agents.

5 For therapy, the compound is generally given as a pharmacological composition containing the compound of the formula (I) or a pharmaceutically acceptable salt thereof in an amount sufficient to show an anti-depressive effect, and a pharmaceutically acceptable carrier. A pharmacological
10 composition containing about 1 - 500 mg of the active ingredient per unit dose is desirable.

According to a conventional method, it is prepared into tablets, lozenges, capsules, powders, aqueous or oily suspensions, syrups, elixirs, aqueous solutions and the like.
15 The pharmacological composition to be used naturally shows properties that vary depending on the objective administration route. For example, an oral composition may be tablet or capsule, and may contain a conventional excipient such as binder (starch etc.) and moistening agent (sodium laurylsulfate
20 etc.). A solution or suspension of the present invention containing a conventional pharmacological vehicle may be used for parenteral administration, such as an aqueous solution for intravenous injection and oily suspension for intramuscular injection.

25 **Examples**

The present invention is described in detail in the following by Starting Material Synthesis Examples, Examples, Formulation Examples and Experimental Examples. The present invention is not limited in any way by these examples.

30 **Starting Material Synthesis Example 1**

(S)-1-(4-glycidyoxybenzo(b)furan-2-ylcarbonyl)pyrrolidine

To a solution (30 ml) of (S)-1-(4-hydroxybenzo(b)furan-2-ylcarbonyl)pyrrolidine (1.3 g) in N,N-dimethylformamide (DMF) were added potassium carbonate (2.2 g) and (S)-glycidyl

nosylate (1.7 g), and the mixture was stirred for 10 hr at room temperature, followed by pouring into water. After extraction with ethyl acetate, the organic layer was washed with water, dried over anhydrous magnesium sulfate, and concentrated under
5 reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.2 g) as a yellow oil.

¹H-NMR(CDCl₃)δ:1.93(penth, J=6.4, 2H), 2.00(penth, J=6.4, 2H), 2.79(dd, J=4.9, 2.9, 1H), 2.93(t, J=4.9, 1H), 3.38-3.43(m, 1H),
10 3.69(t, J=6.8, 2H), 3.92(t, J=6.8, 2H), 4.08(dd, J=11.2, 5.8, 1H), 4.36(dd, J=11.2, 3.0, 1H), 6.00(d, J=8.3, 1H), 7.15(d, J=8.3, 1H), 7.28(t, J=8.3, 1H), 7.47(s, 1H)

Starting Material Synthesis Example 2

(S)-4-(4-glycidyoxybenzo(b)furan-2-ylcarbonyl)morpholine

15 To a suspension (30 ml) of sodium hydride (0.52 g) in DMF was dropwise added a solution (30 ml) of 4-(4-hydroxybenzo(b)furan-2-ylcarbonyl)morpholino in DMF at a reaction temperature of 4°C over 10 min, and the mixture was stirred for 30 min. Thereto was added a solution (10 ml) of
20 (S)-glycidyl nosylate (3.4 g) in DMF, and the mixture was stirred for 30 min and poured into water. After extraction with ethyl acetate, the organic layer was washed with water, dried over anhydrous magnesium sulfate and then concentrated under reduced pressure. The obtained residue was purified by
25 silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.3 g) as a yellow oil.

¹H-NMR(CDCl₃)δ:2.81(dd, J=4.9, 2.4, 1H), 2.96(t, J=4.9, 1H), 3.42-3.44(m, 1H), 3.78-4.07(m, 8H), 4.09(dd, J=10.8, 5.9, 1H), 4.40(dd, J=10.8, 3.0, 1H), 6.69(d, J=8.3, 1H), 7.16(d,
30 J=8.3, 1H), 7.32(t, J=8.3, 1H), 7.44(s, 1H)

Starting Material Synthesis Example 3

methyl (S)-4-glycidyoxybenzo(b)furan-2-carboxylate

To a solution (60 ml) of methyl 4-hydroxybenzo(b)furan-2-carboxylate (3.6 g) in DMF were added (S)-glycidyl nosylate

5 (5.1 g) and potassium carbonate (6.5 g) and the mixture was stirred at room temperature for 8 hr. The reaction mixture was concentrated under reduced pressure and ethyl acetate was added to the residue. The mixture was washed with water, dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to give the title compound (4.1 g) as a yellow crystalline compound.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.82(dd, $J=4.9, 3.0$, 1H), 2.96(t, $J=4.9$, 1H), 3.41-3.45(m, 1H), 3.97(s, 3H), 4.09(dd, $J=10.8, 5.9$, 1H), 10 4.40(dd, $J=10.8, 3.0$, 1H), 6.69(d, $J=8.3$, 1H), 7.22(d, $J=8.3$, 1H), 7.36(t, $J=8.3$, 1H), 7.68(s, 1H)

Starting Material Synthesis Example 4

4-(8-methoxy-2H-chromen-3-ylcarbonyl)morpholine

To a solution (200 ml) of 8-methoxy-2H-chromene-3-15 carboxylic acid (10.0 g) in DMF were added triethylamine (8.6 ml) and diethyl cyanophosphate (10.0 ml) and the mixture was stirred at room temperature for 3 hr. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with water and dried over anhydrous20 magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/ethyl acetate) to give the title compound (3.5 g) as a brown oil.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 3.69-3.78(m, 8H), 4.94(s, 2H), 6.60(s, 1H), 25 6.71(d, $J=5.2$, 1H), 6.87-6.90(m, 2H)

Starting Material Synthesis Example 5

4-(8-hydroxy-2H-chromen-3-ylcarbonyl)morpholine

To a solution (70 ml) of 4-(8-methoxy-2H-chromen-3-ylcarbonyl)morpholine (3.5 g) in methylene chloride was added30 dropwise boron tribromide (9.5 g) at -78°C . The reaction temperature was set to room temperature and the mixture was stirred for 2 hr. The reaction mixture was poured into water and stirred for 1 hr. The organic layer was separated and washed with water and dried over anhydrous magnesium sulfate.

The solvent was evaporated under reduced pressure to give the title compound (3.3 g) as brown crystals.

¹H-NMR(CDCl₃)δ:3.69-3.73(brs, 8H), 4.95(s, 2H), 5.83(brs, 1H), 6.61(s, 1H), 6.65(d, J=7.3, 1H), 6.83(t, J=7.3, 1H), 7.89(d, J=7.3, 1H)

Starting Material Synthesis Example 6

(S)-4-(8-glycidyloxy-2H-chromen-3-ylcarbonyl)morpholine

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 4-(8-hydroxy-2H-chromen-3-ylcarbonyl)morpholine (3.3 g), potassium carbonate (3.5 g) and (S)-glycidyl nosylate (3.3 g), the title compound (3.1 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ:2.74(dd, J=4.9, 2.4, 1H), 2.91(t, J=4.9, 1H), 3.37-3.39(m, 1H), 3.69-3.73(brs, 8H), 4.03(dd, J=11.7, 5.8, 1H), 4.11-4.13(m, 1H), 4.28(dd, J=11.7, 3.4, 1H), 4.94(s, 2H), 6.60(s, 1H), 6.75(d, J=7.3, 1H), 6.87(t, J=7.3, 1H), 6.91(d, J=7.3, 1H)

Starting Material Synthesis Example 7

8-methoxy-N,N-dimethyl-2H-chromene-3-carboxamide

By the reactions in the same manner as in Starting Material Synthesis Example 4 using 8-methoxy-2H-chromene-3-carboxylic acid (8.0 g), triethylamine (14.0 ml) and diethyl cyanophosphate (8.2 ml), the title compound (3.2 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ:3.83(s, 6H), 4.84(s, 2H), 6.45(d, J=8.3, 1H), 6.50(d, J=8.3, 1H), 6.99(s, 1H), 7.13(t, J=8.3, 2H)

Starting Material Synthesis Example 8

(S)-8-glycidyloxy-N,N-dimethyl-2H-chromene-3-carboxamide

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 8-methoxy-N,N-dimethyl-2H-chromene-3-carboxamide (3.2 g) and boron tribromide (11.0 g), a brown oil (3.0 g) was obtained. To a solution (50 ml) of this brown oil in DMF were added potassium carbonate (3.8 g) and (S)-glycidyl nosylate (3.8 g), and the mixture was stirred at

room temperature for 10 hr and poured into water. After extraction with ethyl acetate, the organic layer was washed with water, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was
5 purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.2 g) as yellow crystals, melting point 115-117°C.

Starting Material Synthesis Example 9

ethyl 4-benzyloxy-1-methylindole-2-carboxylate

10 To a solution (100 ml) of ethyl 4-benzyloxy-1H-indole-2-carboxylate (12.0 g) in DMF was added sodium hydride (1.6 g) and the mixture was stirred at room temperature for 10 min. To this reaction mixture was added methyl iodide (2.2 g) and the mixture was stirred for one more hour. The reaction mixture
15 was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous solution of ammonium chloride and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give the title compound (13.2 g) as a brown oil.

20 ¹H-NMR(CDCl₃)δ:1.39(t, J=6.9, 3H), 4.06(s, 3H), 4.35(q, J=6.9, 2H), 5.22(s, 2H), 6.66(d, J=7.8, 1H), 6.98(t, J=7.8, 1H), 7.40(t, J=7.4, 1H), 7.45-7.51(m, 6H)

Starting Material Synthesis Example 10

ethyl 4-hydroxy-1-methylindole-2-carboxylate

25 To a solution (200 ml) of ethyl 4-benzyloxy-1-methylindole-2-carboxylate (13.0 g) in ethanol was added 10% palladium-carbon (1.3 g), and the mixture was stirred at room temperature for 8 hr under a hydrogen atmosphere. The palladium-carbon was filtered off with celite and the reaction
30 mixture was concentrated under reduced pressure to give the title compound (8.0 g) as a brown oil.

¹H-NMR(CDCl₃)δ:1.40(t, J=6.9, 3H), 4.05(s, 3H), 4.37(q, J=6.9, 2H), 6.52(d, J=7.8, 1H), 6.95(t, J=7.8, 1H), 7.19(t, J=7.4, 1H), 7.41(s, 1H)

Starting Material Synthesis Example 11

Ethyl 4-benzyloxy-1-(2-methylpropyl)indole-2-carboxylate

By the reactions in the same manner as in Starting Material Synthesis Example 9 using ethyl 4-benzyloxy-1H-indole-2-carboxylate (10.0 g), sodium hydride (1.6 g) and isobutyl iodide (3.3 ml), the title compound (6.0 g) was obtained as a brown oil.

¹H-NMR(CDCl₃) δ: 0.89 (d, J=6.3, 6H), 1.39 (t, J=7.3, 3H), 2.22 (penth, J=6.3, 1H), 4.25-4.42 (m, 2H), 4.35 (q, J=7.3, 1H), 5.21 (s, 2H), 6.54 (d, J=7.8, 1H), 7.00 (d, J=7.8, 1H), 7.20 (t, J=7.8, 1H), 7.33-7.1 (m, 5H)

Starting Material Synthesis Example 12

ethyl 4-hydroxy-1-(2-methylpropyl)indole-2-carboxylate

By the reactions in the same manner as in Starting Material Synthesis Example 10 using ethyl 4-benzyloxy-1-(2-methylpropyl)indole-2-carboxylate (6.0 g) and 10% palladium-carbon (0.6 g), the title compound was obtained as pale-brown crystals.

¹H-NMR(CDCl₃) δ: 0.89 (d, J=6.3, 6H), 1.40 (t, J=7.3, 3H), 2.21 (penth, J=6.3, 1H), 4.25-4.42 (m, 2H), 4.35 (q, J=7.3, 1H), 6.49 (d, J=7.8, 1H), 6.96 (d, J=7.8, 1H), 7.16 (t, J=7.8, 1H), 7.42 (s, 1H)

Starting Material Synthesis Example 13

3-chloro-6-methoxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide

3.0 g of 3-chloro-6-methoxy-benzo(b)thiophene-2-carboxylic acid (7.0 g) synthesized from 4-methoxycinnamic acid (10.0 g) and thionyl chloride (15 ml) according to the method described in J. Med. Chem. 1992, 35, 958-965 was reacted with dimethylamine hydrochloride and triethylamine in THF to give the title compound (1.9 g) as a brown oil.

¹H-NMR(CDCl₃): 3.09 (bs, 3H), 3.12 (bs, 3H), 3.89 (s, 3H), 7.10 (d, 1H, J=8.8), 7.26 (s, 1H), 7.71 (d, 1H, J=8.8)

Starting Material Synthesis Example 14

(S)-3-chloro-6-glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-

carboxamide

3-Chloro-6-methoxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide (1.9 g) was dissolved in methylene chloride (100 ml) and the mixture was cooled to -78°C. Boron tribromide (4 ml) was added dropwise, and after the temperature rose to room temperature, the mixture was poured into water and extracted with chloroform. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure. The obtained residue was dissolved in DMF (40 ml). By the reactions in the same manner as in Starting Material Synthesis Example 1 using potassium carbonate (3.0 g) and (S)-glycidyl nosylate (2.1 g), the title compound (10 g) was obtained as a brown oil.

¹H-NMR(CDCl₃): 2.80 (dd, 1H, J=4.8, 2.9), 2.95 (t, 1H, J=4.8), 3.11 (bs, 3H), 3.17 (bs, 3H), 3.41 (m, 1H), 4.00 (dd, 1H, J=5.9, 10.8), 4.35 (dd, 1H, J=3.0, 11.5), 7.13 (dd, 1H, J=2.5, 8.7), 7.26 (s, 1H), 7.72 (d, 1H, J=8.8)

Starting Material Synthesis Example 15

4-(methoxymethyloxy)benzo(b)thiophene-2-carboxylic acid

4-(Methoxymethyloxy)benzo(b)thiophene (83 g) was dissolved in THF (700 ml) and the mixture was cooled to -78°C. At this temperature, a solution (363 ml) of n-butyllithium in hexane was added dropwise. The temperature was raised to 0°C and then cooled again to -35°C, and carbon dioxide was bubbled. After the completion of the reaction, the reaction mixture was poured into water, and in the presence of ice, hydrochloric acid was added to adjust its pH to 1 and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure to give the title compound (80 g).

¹H-NMR(CDCl₃): 3.55 (s, 3H), 5.37 (s, 2H), 7.04 (d, 1H, J=7.8), 7.41 (t, 1H, J=7.8), 7.50 (d, 1H, J=8.2), 8.36 (s, 1H)

Starting Material Synthesis Example 16

4-(methoxymethyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide

4-(Methoxymethyloxy)benzo(b)thiophene-2-carboxylic acid
5 (9.6 g) obtained in Starting Material Synthesis Example 15 was dissolved in dimethylformamide (75 ml). Triethylamine (17 ml) and dimethylamine hydrochloride (4.9 g) were added and the mixture was stirred. After 15 min, diethyl cyanophosphate (10 ml) was added, and the mixture was stirred at room temperature
10 for 3 hr. Aqueous hydrochloric acid was added under cooling to make the reaction mixture acidic (pH 1), and then the mixture was stirred at 45°C for 5 hr. The reaction mixture was poured into water, extracted three times with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate.
15 After filtration, the solvent was evaporated under reduced pressure. To the obtained residue was added 6N aqueous hydrochloric acid and the mixture was stirred with heating at 50°C for 1 hr. The reaction mixture was extracted with ethyl acetate and the organic layer was dried over anhydrous
20 magnesium sulfate. The solvent was evaporated under reduced pressure to give the title compound (9.0 g).
¹H-NMR(CDCl₃): 3.17(bs, 3H), 3.28(bs, 3H), 6.76(d, 1H, J=7.8), 7.23(t, 1H, J=7.8), 7.36(d, 1H, J=7.8), 7.81(s, 1H).

Starting Material Synthesis Example 17

25 (S)-4-glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide

To a solution of N,N-dimethyl-4-(hydroxymethyloxy)benzo-
(b)thiophene-2-carboxamide (9.0 g) in DMF (100 ml) was added potassium carbonate (8.0 g), and (S)-glycidyl nosylate (8.0 g) was further added. The mixture was stirred at 60°C for 2 hr.
30 The reaction mixture was concentrated under reduced pressure and water was added. The mixture was extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained crystals were recrystallized from ethyl acetate to give the

title compound (7.5 g).

¹H-NMR(CDCl₃): 2.81(dd, 1H, J=2.4, 4.9), 2.96(t, 1H, J=4.4),
3.00-3.21(bs, 6H), 3.44-3.48(m, 1H), 4.08(dd, 1H, J=5.8, 11.2),
4.41(dd, 1H, J=2.4, 11.2), 6.76(d, 1H, J=7.8), 7.32(t, 1H,
5 J=7.8), 7.45(d, 1H, J=8.3), 7.73(s, 1H)


Starting Material Synthesis Example 18

(S)-4-(4-glycidyoxybenzo(b)thiophen-2-ylcarbonyl)morpholine

By the reactions in the same manner as in Starting
Material Synthesis Example 16 using 4-(methoxymethoxy)-
10 benzo(b)thiophene-2-carboxylic acid (3.5 g), morpholine (1.0 g)
and diethyl cyanophosphate (3.1 g), 4-(4-hydroxybenzo(b)-
thiophene-2-carbonyl)morpholine (3.2 g) was obtained as a brown
oil. By the reactions in the same manner as in Starting
Material Synthesis Example 1 using the brown oil (2.0 g) and
15 (S)-glycidyl nosylate (2.1 g), the title compound (2.0 g) was
obtained as brown crystals.

¹H-NMR(CDCl₃): 2.81(dd, 1H, J=1.9, 4.8), 2.97(t, 1H, J=4.8),
3.42-3.48(m, 1H), 3.86-3.95(bs, 8H), 4.05(dd, 1H, J=5.6, 11.2),
4.43(dd, 1H, J=2.9, 11.4), 6.77(d, 1H, J=8.3), 7.33(t, 1H,
20 J=7.8), 7.45(d, 1H, J=7.8), 7.68(s, 1H)

The structural formulas of the compounds obtained from
the starting material synthesis examples 1 to 18 are shown in
the following.



Starting Material Synthesis Example 19

(S)-1-(4-glycidyloxybenzo(b)thiophen-2-ylcarbonyl)pyrrolidine

By the reactions in the same manner as in Starting Material Synthesis Example 16 using 4-(methoxymethyloxy)-benzo(b)thiophene-2-carboxylic acid (3.0 g), pyrrolidine (0.75 g) and diethyl cyanophosphate (2.5 g), 1-(4-hydroxybenzo(b)-thiophene-2-carbonyl)pyrrolidine (2.4 g) was obtained as a brown oil. By the reactions in the same manner as in Starting Material Synthesis Example 1 using the brown oil (2.0 g) and (S)-glycidyl nosylate (2.0 g), the title compound (0.45 g) was obtained as brown crystals.

¹H-NMR(CDCl₃): 1.98-2.10 (bs, 4H), 2.80 (dd, 1H, J=2.9, 4.9), 2.96 (t, 1H, J=4.2), 3.42-3.48 (m, 1H), 3.70 (bs, 2H), 3.87 (bs, 2H), 4.07 (dd, 1H, J=4.8, 11.2), 4.41 (dd, 1H, J=2.9, 11.2), 6.74 (d, 1H, J=7.8), 7.32 (t, 1H, J=7.8), 7.44 (d, 1H, J=8.3), 8.00 (s, 1H)

Starting Material Synthesis Example 20

(S)-4-glycidyloxy-N-methoxy-N-methylbenzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Starting Material Synthesis Example 16 using 4-(methoxymethyloxy)-benzo(b)thiophene-2-carboxylic acid (4.5 g), N,O-dimethylhydroxylamine hydrochloride (2.1 g) and diethyl cyanophosphate (3.2 g), 4-hydroxy-N-methoxy-N-methylbenzo(b)-thiophene-2-carboxamide (4.0 g) was obtained as a brown oil. By the reactions in the same manner as in Starting Material Synthesis Example 1 using the brown oil (2.0 g) and (S)-glycidyl nosylate (2.0 g), the title compound (1.1 g) was obtained as brown crystals.

¹H-NMR(CDCl₃): 2.78 (dd, 1H, J=2.8, 4.8), 2.98 (t, 1H, J=4.2), 3.42 (s, 3H), 3.43-3.48 (m, 1H), 3.83 (s, 3H), 4.10 (dd, 1H, J=4.9, 11.2), 4.36 (dd, 1H, J=3.5, 11.3), 6.74 (d, 1H, J=7.8), 7.33 (t, 1H, J=8.3), 7.44 (d, 1H, J=8.3), 8.40 (s, 1H)

Starting Material Synthesis Example 21

methyl (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylate

To a solution (70 ml) of methyl (S)-4-glycidyoxybenzo(b)furan-2-carboxylate (4.1 g) obtained in Starting Material Synthesis Example 3 in methanol (70 ml) was added 4-(naphthalen-2-yl)piperidine (3.5 g) at room temperature, and the mixture was refluxed under heating for 2 hr. The solvent was evaporated under reduced pressure and the obtained residue was purified by silica gel column chromatography (chloroform:methanol) to give the title compound (5.6 g) as yellow crystals, melting point 118-119°C.

Starting Material Synthesis Example 22

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylic acid

To a solution (140 ml) of methyl (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)benzo(b)furan-2-carboxylate (5.6 g) in methanol was added 2.0 M aqueous potassium hydroxide solution (100 ml) and the mixture was refluxed under heating for 2 hr. The reaction mixture was poured into water and the aqueous solution was made acidic (pH=1) with conc. hydrochloric acid. The solution was extracted with a mixed solvent of chloroform-methanol (2:1) and the organic layer was washed with saturated brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The obtained residue was crystallized from ethyl acetate, and the crystals were collected by filtration and dried to give hydrochloride (4.7 g) of the title compound as pale-yellow crystals, melting point 234-235°C (decomposition).

Starting Material Synthesis Example 23

ethyl (S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)benzo(b)furan-2-carboxylate

By the reactions in the same manner as in Starting Material Synthesis Example 11 using ethyl (S)-7-

(glycidylloxy)benzo(b)furan-2-carboxylate (5.3 g) and 4-(naphthalen-2-yl)piperidine (3.0 g), the title compound (5.2 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ:1.41(t, J=7.3, 3H), 1.87-1.98(m, 4H), 2.23(t, J=7.3, 1H), 2.25-2.63(m, 1H), 2.48-2.79(m, 4H), 3.05(d, J=10.7, 1H), 3.05(d, J=10.7, 1H), 3.23(d, J=10.7, 1H), 4.10-4.28(m, 3H), 4.45(q, J=7.3, 2H), 6.72(d, J=8.3, 1H), 7.21(d, J=8.3, 1H), 7.35-7.49(m, 4H), 7.67-7.70(m, 2H), 7.75-7.82(m, 3H)

Starting Material Synthesis Example 24

10 (S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-benzo(b)furan-2-carboxylic acid

To a solution of ethyl (S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)benzo(b)furan-2-carboxylate (5.2 g) in methanol (50 ml) was added 10% saturated aqueous sodium hydroxide solution (50 ml) and the mixture was refluxed under heating for 1 hr. The reaction mixture was made acidic (pH 1) with conc. hydrochloric acid and extracted with chloroform. The organic layer was washed with water and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give the title compound (4.0 g) as a brown oil.

¹H-NMR(DMSO-d₆)δ:1.81-2.20(m, 4H), 2.80-3.17(m, 2H), 4.01(dd, J=9.3, 3.4, 1H), 4.12(dd, J=9.3, 3.4, 1H), 6.75(d, J=8.3, 1H), 7.19(d, J=8.3, 1H), 7.48(t, J=8.3, 1H), 7.44-7.51(m, 3H), 7.77(s, 1H), 7.87-7.90(m, 3H), 8.04(s, 1H)

Starting Material Synthesis Example 25

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1H-indole-2-carboxylic acid

To a solution of ethyl 4-hydroxy-1H-indole-2-carboxylate (1.3 g) in DMF (50 ml) were added potassium carbonate and (S)-glycidyl nosylate (1.0 g) and the mixture was stirred one day. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with water and dried over anhydrous magnesium sulfate and the solvent was

Starting Material Synthesis Example 27

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1-(2-methylpropyl)indole-2-carboxylic acid

By the reactions in the same manner as in Starting
5 Material Synthesis Example 25 using ethyl 4-hydroxy-1-(2-methylpropyl)indole-2-carboxylate (5.0 g) obtained in Starting Material Synthesis Example 12, (S)-glycidyl nosylate (4.5 g) and 4-(naphthalen-2-yl)piperidine (5.3 g), ethyl (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1-(2-
10 methylpropyl)indole-2-carboxylate (7.5 g) was obtained. This was dissolved in ethanol (40 ml) and water (30 ml) and potassium hydroxide (4.0 g) were added. The mixture was refluxed for 2.5 hr. From the obtained reaction mixture, ethanol was evaporated under reduced pressure and 1N aqueous
15 hydrochloric acid solution (30 ml) was added under ice-cooling. The mixture was extracted with chloroform. The obtained organic layer was washed with saturated brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and isopropyl ether was added to the obtained
20 oil. The obtained crystals were collected by filtration to give the title compound (6.7 g) as pale-yellow crystals.
 $^1\text{H-NMR}(\text{CD}_3\text{OD})\delta$: 0.84-0.86 (m, 7H), 2.15-2.23 (m, 5H), 3.11-3.65 (m, 4H), 3.65 (m, 2H), 4.18-4.25 (m, 2H), 4.40 (d, $J=7.3$, 2H), 4.58 (m, 1H), 6.60 (d, $J=7.8$, 1H), 7.10 (d, $J=8.3$, 1H), 7.24 (dd, $J=7.8$,
25 8.3, 1H), 7.46-7.47 (m, 4H), 7.74-7.86 (m, 4H)

Starting Material Synthesis Example 28

1-(hydroxyimino)-1-(4-methoxybenzo(b)furan-2-yl)methylamine

To a solution (40 ml) of 4-methoxybenzo(b)furan-2-carbonitrile (2.8 g) in ethanol were added hydroxylamine
30 hydrochloride (1.2 g) and sodium hydrogencarbonate (3.0 g). The mixture was refluxed under heating for 1.5 hr. The inorganic material was filtered off and the reaction mixture was concentrated under reduced pressure to give the title compound (3.4 g) as brown crystals.

¹H-NMR(CDCl₃) δ: 3.94(s, 3H), 6.68(d, J=7.8, 1H), 7.13(d, J=7.8, 1H), 7.19(s, 1H), 7.26(t, J=7.8, 1H)

Starting Material Synthesis Example 29

3-(4-methoxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole

5 1-(Hydroxyimino)-1-(4-methoxybenzo(b)furan-2-yl)methylamine (3.4 g) was dissolved in acetic anhydride (40 ml) and the mixture was refluxed under heating for 14 hr. The reaction mixture was concentrated under reduced pressure and the obtained residue was recrystallized from acetonitrile to
10 give the title compound (1.1 g) as pale-brown crystals.

¹H-NMR(CDCl₃) δ: 2.68(s, 3H), 3.97(s, 3H), 6.70(d, J=8.3, 1H), 7.22(d, J=8.3, 1H), 7.33(t, J=8.3, 1H), 7.58(s, 1H)

Starting Material Synthesis Example 30

3-(4-hydroxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole

15 By the reactions in the same manner as in Starting Material Synthesis Example 5 using 3-(4-methoxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole (1.1 g) and boron tribromide (4.2 ml), the title compound (0.75 g) was obtained as yellow crystals.

20 ¹H-NMR(DMSO-d₆) δ: 2.65(s, 3H), 6.68(d, J=7.8, 1H), 7.12(d, J=8.3, 1H), 7.23(dd, J=7.8, 8.3, 1H), 7.60(s, 1H), 10.30(s, 1H)

Starting Material Synthesis Example 31

(S)-3-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole

25 By the reactions in the same manner as in Starting Material Synthesis Example 1 using 3-(4-hydroxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole (0.75 g) and (S)-glycidyl nosylate (0.93 g), the title compound (0.45 g) was obtained as white crystals.

30 ¹H-NMR(CDCl₃) δ: 2.69(s, 3H), 2.83(dd, J=4.9, 2.5, 1H), 2.96(t, J=4.9, 1H), 3.43-3.45(m, 1H), 4.13(dd, J=11.2, 4.4, 1H), 4.40(dd, J=11.2, 3.0, 1H), 6.71(d, J=7.8, 1H), 7.25(d, J=8.3, 1H), 7.32(dd, J=8.3, 7.8, 1H), 7.62(s, 1H)

Starting Material Synthesis Example 32

1-(hydroxyimino)-1-(7-methoxybenzo(b)furan-2-yl)methylamine

By the reactions in the same manner as in Starting Material Synthesis Example 28 using 7-methoxybenzo(b)furan-2-carbonitrile (3.0 g), hydroxylamine hydrochloride (1.4 g) and sodium hydrogencarbonate (2.1 g), the title compound (3.3 g) was obtained as brown crystals.

$^1\text{H-NMR}(\text{CD}_3\text{OD})\delta$: 3.97(s, 3H), 6.89-6.91(m, 1H), 7.11-7.17(m, 3H)
















Starting Material Synthesis Example 33

3-(7-methoxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 29 using 1-(hydroxyimino)-1-(7-methoxybenzo(b)furan-2-yl)methylamine (3.3 g), the title compound (1.7 g) was obtained as white crystals.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.68(s, 3H), 4.03(s, 3H), 6.90(d, J=7.8, 1H), 7.21(d, J=7.8, 1H), 7.25(t, J=7.8, 1H), 7.45(s, 1H)

The structural formulas of the compounds obtained in Starting Material Synthesis Examples 19 to 33 are shown in the following.



Starting Material Synthesis Example 34

3-(7-hydroxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 3-(7-methoxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole (1.7 g) and boron tribromide (6.5 ml), the title compound (1.5 g) was obtained as white crystals. ¹H-NMR(DMSO-d₆)δ:2.65(s, 3H), 6.68(d, J=7.8, 1H), 7.12(d, J=8.3, 1H), 7.23(dd, J=7.8, 8.3, 1H), 7.60(s, 1H), 10.30(s, 1H)

Starting Material Synthesis Example 35

(S)-3-(7-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 3-(7-hydroxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole (1.5 g) and (S)-glycidyl nosylate (1.8 g), the title compound (1.7 g) was obtained as white crystals.

¹H-NMR(CDCl₃)δ:2.69(s, 3H), 2.81(dd, J=4.9, 2.4, 1H), 2.94(t, J=4.9, 1H), 3.46-3.48(m, 1H), 4.26(dd, J=11.2, 5.4, 1H), 4.46(dd, J=11.2, 3.4, 1H), 6.95(d, J=7.8, 1H), 7.21(t, J=7.8, 1H), 7.29(d, J=7.8, 1H), 7.46(s, 1H)

Starting Material Synthesis Example 36

N'-(4-methoxybenzo(b)furan-2-ylcarbonyl)acetohydrazide

To a solution (700 ml) of 4-methoxybenzo(b)furan-2-carboxylic acid (43.4 g) in THF was added 1,1'-carbonylbis-1H-imidazole (CDI) (38.4 g) and the mixture was stirred at room temperature for 1 hr. Acetohydrazine (17.6 g) was added to this reaction mixture, and the mixture was stirred for 1 more hr. The reaction mixture was poured into water, and the precipitated crystals were collected by filtration and dried to give the title compound (38.4 g) as pale-brown crystals.

¹H-NMR(DMSO-d₆)δ:1.91(s, 3H), 3.93(s, 3H), 6.86(d, J=7.8, 1H), 7.25(d, J=7.8, 1H), 7.42(t, J=7.8, 1H), 7.61(s, 1H), 9.92(s, 1H), 10.46(s, 1H)

Starting Material Synthesis Example 37

2-(4-methoxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole

To a solution (400 ml) of N'-(4-methoxybenzo(b)furan-2-ylcarbonyl)acetohydrazide (15.6 g) in 1,2-dichloroethane were
5 added triethylamine (21 ml) and triphenylphosphine (19.8 g) and
the reaction temperature was set to 5°C. To this reaction
mixture was added dropwise diethyl azodicarboxylate (40%
toluene solution) (33 ml) over 15 min. The reaction
temperature was set to room temperature and the mixture was
10 stirred for 1.5 hr and washed with saturated aqueous solution
of ammonium chloride. After partitioning, the obtained organic
layer was dried over anhydrous magnesium sulfate and the
solvent was evaporated under reduced pressure. The obtained
residue was concentrated under reduced pressure and purified by
15 silica gel column chromatography (chloroform/ethyl acetate) to
give the title compound (4.6 g) as pale-yellow crystals.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.65(s, 3H), 3.97(s, 3H), 6.72(d, J=8.3, 1H),
7.22(d, J=8.3, 1H), 7.36(t, J=8.3, 1H), 7.56(s, 1H)

Starting Material Synthesis Example 38

20 2-(4-hydroxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole

By the reactions in the same manner as in Starting
Material Synthesis Example 5 using 2-(4-methoxybenzo(b)furan-2-
yl)-5-methyl-1,3,4-oxadiazole (6.5 g) and boron tribromide (27
ml), the title compound (3.3 g) was obtained as yellow crystals.
25 $^1\text{H-NMR}(\text{DMSO}-d_6)\delta$: 2.60(s, 3H), 6.71(d, J=8.3, 1H), 7.16(d, J=8.3,
1H), 7.29(t, J=8.3, 1H), 7.68(s, 1H)

Starting Material Synthesis Example 39

(S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole

30 By the reactions in the same manner as in Starting
Material Synthesis Example 1 using 2-(4-hydroxybenzo(b)furan-2-
yl)-5-methyl-1,3,4-oxadiazole (3.3 g) and (S)-glycidyl nosylate
(3.7 g), the title compound (1.1 g) was obtained as white
crystals.

¹H-NMR(CDCl₃)δ:2.65(s, 3H), 2.83(dd, J=4.9, 2.4, 1H), 2.96(t, J=4.9, 1H), 3.43-3.46(m, 1H), 4.09(dd, J=11.2, 5.8, 1H), 4.42(dd, J=11.2, 2.9, 1H), 6.72(d, J=8.3, 1H), 7.23(d, J=8.3, 1H), 7.34(t, J=8.3, 1H), 7.59(s, 1H)

5 **Starting Material Synthesis Example 40**

2-(7-methoxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole

7-Methoxybenzo(b)furan-2-carboxylic acid (10 g) was dissolved in tetrahydrofuran (100 ml) and CDI (12.6 g) and acetohydrazine (4.0 g) were added. The mixture was stirred at
10 room temperature for 2 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (19 g). This oily product (19 g) was dissolved in 1,2-dichloroethane (300
15 ml) and triphenylphosphine (39 g) and triethylamine (25 ml) were added. The mixture was stirred under ice-cooling. Diisopropyl azodicarboxylate (40% toluene solution) (75 g) was added and then the mixture was stirred at room temperature for 3 hr. The reaction mixture was poured into ice water and
20 extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (8.0 g) as pale-yellow crystals.

25 ¹H-NMR(CDCl₃)δ:2.65(s, 3H), 4.05(s, 3H), 6.92(d, J=7.8, 1H), 7.23-7.28(m, 2H), 7.51(s, 1H)

Starting Material Synthesis Example 41

N'-(4-(methoxymethyloxy)benzo(b)thiophen-2-ylcarbonyl)-acetohydrazide

30 4-(Methoxymethyloxy)benzothiophene-2-carboxylic acid (7 g) was dissolved in tetrahydrofuran (100 ml) and CDI (7.3 g) and acetohydrazine (2.4 g) was added. The mixture was stirred at room temperature for 3 hr. The precipitated crystals were collected by filtration to give the title compound (3.9 g).

¹H-NMR (DMSO-d₆) δ: 1.99 (s, 3H), 3.32 (bs, 2H), 3.51 (s, 3H), 5.37 (s, 2H), 7.03 (d, J=7.8, 1H), 7.36 (t, J=7.8, 1H), 7.52 (d, J=7.8, 1H), 8.32 (s, 1H)

Starting Material Synthesis Example 42

5 2-(4-(methoxymethyloxy)benzo(b)thiophen-2-yl)-5-methyl-1,3,4-oxadiazole

N'-(4-(Methoxymethyloxy)benzo(b)thiophen-2-ylcarbonyl)acetohydrazide (2.4 g) was dissolved in 1,2-dichloroethane (50 ml) and triphenylphosphine (3.2 g) and
10 triethylamine (2 ml) were added. The mixture was stirred under ice-cooling. Diethyl azodicarboxylate (40% toluene solution) (5.2 g) was added and the mixture was stirred at room temperature for 1 hr. The reaction mixture was poured into ice water and extracted with chloroform. The organic layer was
15 dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title product (1.4 g) as pale-yellow crystals.

¹H-NMR (CDCl₃) δ: 2.61 (s, 3H), 3.54 (s, 3H), 5.38 (s, 2H), 7.05 (d, J=7.8, 1H), 7.38 (t, J=7.8, 1H), 7.52 (d, J=7.8, 1H), 8.12 (s, 1H)
20

Starting Material Synthesis Example 43

2-(4-hydroxybenzo(b)thiophen-2-yl)-5-methyl-1,3,4-oxadiazole

2-(4-(Methoxymethyloxy)benzo(b)thiophen-2-yl)-5-methyl-1,3,4-oxadiazole (1.4 g) was dissolved in a mixed solvent (10
25 ml) of acetic acid-water (1:1) and the mixture was heated at 80°C for 4 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily compound
30 (1.4 g).

¹H-NMR (DMSO-d₆) δ: 2.61 (s, 3H), 6.83 (d, J=7.8, 1H), 7.32 (t, J=7.8, 1H), 7.44 (d, J=7.8, 1H), 8.07 (s, 1H), 10.44 (bs, 1H)

Starting Material Synthesis Example 44

N'-(4-benzyloxy-1H-indol-2-ylcarbonyl)acetohydrazide

Ethyl 4-benzyloxyindol-2-carboxylate (10 g) was dissolved in a mixed solvent (200 ml) of dioxane-water (1:1) and potassium hydroxide (3.8 g) was added. The mixture was refluxed under heating for 2 hr. The reaction mixture was poured into ice water, made acidic with hydrochloric acid and extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 4-benzyloxyindol-2-carboxylic acid as pale-yellow crystals (9.0 g). The crystals were dissolved in dimethylformamide (100 ml) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC, 7.6 g), 1-hydroxybenzotriazole hydrochloride (HOBT, 6.9 g), triethylamine (7.0 ml) and acetohydrazine (2.6 g) were added thereto. The mixture was stirred at room temperature for 6 hr. The reaction mixture was poured into ice water and the precipitated crystals were collected by filtration to give the title compound (10 g).
¹H-NMR(DMSO-d₆)δ:1.93(s, 3H), 5.22(s, 2H), 6.62(d, J=7.8, 1H), 7.04(d, J=7.8, 1H), 7.11(t, J=7.8, 1H), 7.36-7.45(m, 5H), 7.54(s, 1H), 9.85(s, 1H), 10.20(s, 1H), 11.67(s, 1H)

Starting Material Synthesis Example 45

4-benzyloxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indole

N'-(4-Benzyloxy-1H-indol-2-ylcarbonyl)acetohydrazide (7.5 g) was dissolved in tetrahydrofuran (250 ml) and triphenylphosphine (9.0 g) and triethylamine (6 ml) were added. The mixture was stirred under ice-cooling. Diisopropyl azodicarboxylate (40% toluene solution) (17.7 g) was added and the mixture was stirred at 50°C for 2 hr. The reaction mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (6.0 g) as yellow crystals.

¹H-NMR(DMSO-d₆)δ:2.59(s, 3H), 5.25(s, 2H), 6.65(d, J=7.8, 1H), 7.07(d, J=7.8, 1H), 7.15(m, 2H), 7.34(m, 1H), 7.41(m, 2H), 7.53(m, 2H), 12.21(s, 1H)

Starting Material Synthesis Example 46

5 N'-(7-methoxybenzo(b)furan-2-ylcarbonyl)benzohydrazide

7-Methoxybenzo(b)furan-2-ylcarbonylbenzohydrazide (10 g) was dissolved in dichloromethane (100 ml) and triethylamine (9.0 ml) and benzoyl chloride (7.8 g) were added thereto. The mixture was stirred at room temperature for 3 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound 10 (5.0 g) as white crystals.

¹H-NMR(DMSO-d₆)δ:4.00(s, 3H), 7.08(d, J=7.8, 1H), 7.27(t, J=7.8, 1H), 7.35(d, J=7.8, 1H), 7.47-7.60(m, 3H), 7.68(s, 1H), 7.94(m, 2H), 10.57(s, 1H), 10.76(s, 1H)

Starting Material Synthesis Example 47

20 2-(7-methoxybenzo(b)furan-2-yl)-5-phenyl-1,3,4-oxadiazole

N'-(7-Methoxybenzo(b)furan-2-ylcarbonyl)benzohydrazide (5.0 g) was dissolved in thionyl chloride (20 ml) and the mixture was stirred with heating at 80°C for 1 hr. Thionyl chloride was evaporated under reduced pressure and water was added to the residue. The mixture was extracted with ethyl acetate and the organic layer was washed with saturated aqueous solution of sodium hydrogencarbonate, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (3.7 g) as pale-yellow crystals.

30 ¹H-NMR(DMSO-d₆)δ:4.02(s, 3H), 7.12(d, J=7.8, 1H), 7.29(t, J=7.8, 1H), 7.38(d, J=7.8, 1H), 7.63-7.68(m, 3H), 7.88(s, 1H), 8.13(m, 2H)

Starting Material Synthesis Example 48

N'-(4-methoxybenzo(b)furan-2-ylcarbonyl)trifluoroacetohydrazide

To a solution (250 ml) of 4-methoxybenzo(b)furan-2-ylcarbohydrazide (9.5 g) in methylene chloride was added trifluoroacetic anhydride (8.5 ml), and the mixture was stirred at room temperature for 2 hr. The reaction mixture was concentrated under reduced pressure and the residue was crystallized from hexane. The crystals were collected by filtration and dried to give the title compound (10.5 g) as yellow crystals.

¹H-NMR(DMSO-d₆)δ:3.94(s, 3H), 6.89(d, J=8.3, 1H), 7.28(d, J=8.3, 1H), 7.45(t, J=8.3, 1H), 7.66(s, 1H), 11.04(s, 1H), 11.70(s, 1H)

Starting Material Synthesis Example 49

2-(4-methoxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 37 using N'-(4-methoxybenzo(b)furan-2-ylcarbonyl)trifluoroacetohydrazide (5.2 g), triethylamine (7.2 ml), triphenylphosphine (9.0 g) and diethyl azodicarboxylate (40% toluene solution, 6.2 ml), the title compound (4.0 g) was obtained as pale-yellow crystals.

¹H-NMR(CDCl₃)δ:3.98(s, 3H), 6.71(d, J=8.3, 1H), 7.18(d, J=8.3, 1H), 7.48(t, J=8.3, 1H), 7.95(s, 1H)

Starting Material Synthesis Example 50

2-(4-hydroxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 2-(4-methoxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole (4.0 g) and boron tribromide (15 ml), the title compound (3.6 g) was obtained as yellow crystals.

¹H-NMR(DMSO-d₆)δ:6.73(d, J=8.3, 1H), 7.22(d, J=8.3, 1H), 7.36(t, J=8.3, 1H), 10.52(s, 1H)

Starting Material Synthesis Example 51

(S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-trifluoromethyl-

1,3,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 2-(4-hydroxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole (3.3 g) and (S)-glycidyl nosylate (3.7 g), the title compound (1.1 g) was obtained as white crystals.

¹H-NMR(CDCl₃)δ: 2.83(dd, J=4.9, 2.4, 1H), 2.99(t, J=4.9, 1H), 3.44-3.46(m, 1H), 4.12(dd, J=11.2, 5.9, 1H), 4.44(dd, J=11.2, 2.9, 1H), 6.76(d, J=8.3, 1H), 7.27(d, J=8.3, 1H), 7.42(t, J=8.3, 1H), 7.83(s, 1H)

Starting Material Synthesis Example 52

N'-(7-methoxybenzo(b)furan-2-ylcarbonyl)trifluoroacetohydrazide

To a solution (300 ml) of 7-methoxybenzo(b)furan-2-ylcarbonylhydrazide (14.0 g) in methylene chloride was added trifluoroacetic anhydride (11.5 ml) and the mixture was stirred at room temperature for 1 hr. The reaction mixture was concentrated under reduced pressure and the residue was crystallized from hexane, collected by filtration and dried to give the title compound (16.1 g) as white crystals.

¹H-NMR(DMSO-d₆)δ: 7.11(d, J=7.8, 1H), 7.28(t, J=7.8, 1H), 7.35(d, J=7.8, 1H), 7.69(s, 1H), 11.10(s, 1H)

Starting Material Synthesis Example 53





















2-(7-methoxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole

To a solution (280 ml) of N'-(7-methoxybenzo(b)furan-2-ylcarbonyl)trifluoroacetohydrazide (14.6 g) in 1,2-dichloroethane were added thionyl chloride (4.2 ml) and DMF (0.1 ml) and the mixture was refluxed under heating for 4.5 hr.

The solvent was evaporated under reduced pressure and the obtained residue was purified by silica gel column chromatography (chloroform/ethyl acetate) to give the title compound (2.4 g) as pale-yellow crystals.

¹H-NMR(CDCl₃)δ: 4.06(s, 3H), 6.99(d, J=6.9, 1H), 7.22(d, J=6.9, 1H), 7.26-7.31(m, 2H), 7.72(s, 1H)

The structural formulas of the compounds obtained in Starting Material Synthesis Examples 34 to 53 are shown in the following.



Starting Material Synthesis Example 54

2-(7-hydroxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole

5 By the reactions in the same manner as in Starting Material Synthesis Example 5 using 2-(7-methoxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole (2.4 g) and boron tribromide (5.0 ml), the title compound (2.2 g) was obtained as yellow crystals.

10 ¹H-NMR(DMSO-d₆)δ:6.96(d, J=7.3, 1H), 7.19(t, J=7.3, 1H), 7.29(t, d=7.3, 1H), 8.00(s, 1H), 10.50(s, 1H)

Starting Material Synthesis Example 55

(S)-2-(7-glycidyloxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole

15 By the reactions in the same manner as in Starting Material Synthesis Example 1 using 2-(7-hydroxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole (2.4 g) and (S)-glycidyl nosylate (2.2 g), the title compound (1.0 g) was obtained as white crystals.

20 ¹H-NMR(CDCl₃)δ:2.81-2.85(m, 1H), 2.96-2.98(m, 1H), 3.42-3.50(m, 1H), 4.23(dd, J=11.2, 5.8, 1H), 4.52(dd, J=11.2, 3.4, 1H), 7.04(d, J=7.8, 1H), 7.30(t, J=7.8, 1H), 7.33(d, J=7.8, 1H), 7.71(s, 1H)

Starting Material Synthesis Example 56

25 5-(4-methoxybenzo(b)furan-2-yl)-3-methyl-1,2,4-oxadiazole

To a solution (50 ml) of 4-methoxybenzo(b)furan-2-carboxylic acid (1.9 g) in THF were added thionyl chloride (0.9 ml) and DMF (0.1 ml), and the mixture was refluxed under heating for 20 min. The solvent was evaporated under reduced pressure and the obtained residue was dissolved in pyridine (50 ml) and acetamide oxime hydrochloride (1.3 g) was added. The mixture was refluxed under heating for 1 hr and the solvent was evaporated under reduced pressure. The obtained residue was purified by silica gel column chromatography (chloroform:ethyl

acetate=6:1) to give the title compound (1.0 g) as pale-yellow crystals.

¹H-NMR(CDCl₃)δ:2.51(s, 3H), 3.98(s, 3H), 6.73(d, J=7.8, 1H), 7.24(d, J=8.3, 1H), 7.38(dd, J=7.8,8.3,1H), 7.73(s, 1H)

5 **Starting Material Synthesis Example 57**

5-(4-hydroxybenzo(b)furan-2-yl)-3-methyl-1,2,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 5-(4-methoxybenzo(b)furan-2-yl)-3-methyl-1,2,4-oxadiazole (0.98 g) and boron tribromide (3.1 ml), the title compound (0.72 g) was obtained as yellow crystals.

¹H-NMR(CD₃OD)δ:2.44(s, 3H), 6.69(d, J=8.3, 1H), 7.10(d, J=8.3, 1H), 7.31(t, J=8.3, 1H), 7.79(s, 1H)

Starting Material Synthesis Example 58

15 (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-3-methyl-1,2,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 5-(4-hydroxybenzo(b)furan-2-yl)-3-methyl-1,2,4-oxadiazole (3.3 g) and (S)-glycidyl nosylate (3.7 g), the title compound (1.1 g) was obtained as white crystals.

¹H-NMR(CDCl₃)δ:2.51(s, 3H), 2.83(dd, J=4.8, 2.4, 1H), 2.96(t, J=4.8, 1H), 3.42-3.46(m, 1H), 4.11(dd, J=11.2, 5.8, 1H), 4.42(dd, J=11.2, 2.9, 1H), 6.73(d, J=8.3, 1H), 7.26(d, J=8.3, 1H), 7.39(t, J=8.3, 1H), 7.78(s, 1H)

Starting Material Synthesis Example 59

methyl 4-hydroxybenzo(b)thiophene-2-carboxylate

4-(Methoxymethyloxy)benzo(b)thiophene-2-carboxylic acid (7 g) was dissolved in methanol (140 ml) and thionyl chloride (2.0 ml) was added under ice-cooling. The mixture was refluxed under heating for 2 hr and the reaction mixture was concentrated under reduced pressure. Water was added and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and after

filtration, the solvent was evaporated under reduced pressure to give the title compound (6.0 g).

$^1\text{H-NMR}(\text{CDCl}_3)$: 3.95 (s, 3H), 6.82 (d, 1H, $J=4.8$), 7.23-7.38 (m, 2H), 8.30 (s, 1H)

5 Starting Material Synthesis Example 60

5-(4-hydroxybenzo(b)thiophen-2-yl)-3-methyl-1,2,4-oxadiazole

Methyl 4-hydroxybenzo(b)thiophene-2-carboxylate (6.0 g) was dissolved in dimethylformamide (80 ml) and sodium hydride (1.7 g) was added under ice-cooling. The mixture was stirred for 30 min and chloromethyl methyl ether (3 g) was added. The mixture was stirred at room temperature for 3 hr. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure. Tetrahydrofuran (100 ml) was added and the reaction mixture was ice-cooled, and sodium hydride (1.6 g) and acetamide oxime (3.0 g) were added in the presence of molecular sieves (4A). The mixture was refluxed under heating for 30 min and the tetrahydrofuran solution obtained earlier was added to the solution. The mixture was refluxed under heating for 1 hr, and after cooling, poured into water and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure. Thereto were added tetrahydrofuran (35 ml) and 6N hydrochloric acid (20 ml), and the mixture was stirred at 50°C for 30 min. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure to give the title compound (2.4 g).

$^1\text{H-NMR}(\text{CDCl}_3)$: 2.50 (s, 3H), 5.70 (bs, 1H), 6.78 (d, 1H, $J=7.6$), 7.34 (t, 1H, $J=7.8$), 7.47 (d, 1H, $J=8.3$), 8.33 (s, 1H)

Starting Material Synthesis Example 61

(S)-5-(4-glycidyloxybenzo(b)thiophen-2-yl)-3-methyl-1,2,4-

oxadiazole

Synthesized according to a method similar to the method of Starting Material Synthesis Example 1.

¹H-NMR(CDCl₃): 2.48(s, 3H), 2.83(dd, 1H, J=2.4,4.9), 2.98(t, 1H, J=4.4), 3.42-3.48(m, 1H), 4.14(dd, 1H, J=5.9,11.3), 4.41(dd, 1H, J=3.0,10.8), 6.80(d, 1H, J=7.8), 7.40(t, 1H, J=7.8), 7.48(d, 1H, J=8.3), 8.35(s, 1H)

Starting Material Synthesis Example 62

1-(4-methoxybenzo(b)furan-2-yl)butan-1,3-dione

2-Acetyl-4-methoxybenzo(b)furan (2.4 g) was dissolved in ethyl acetate (50 ml), and sodium hydride (1.5 g) was added under ice-cooling. The mixture was stirred at room temperature for 10 min, and the mixture was refluxed under heating for 1 hr.

After cooling, the mixture was poured into water and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (0.7 g).

¹H-NMR(CDCl₃): 2.21(s, 3H), 3.96(s, 3H), 6.25(s, 1H), 6.68(d, 1H, J=7.8), 6.68(d, 1H, J=7.6), 7.15(d, 1H, J=7.8), 7.33(t, 1H, J=7.8), 7.56(s, 1H)

Starting Material Synthesis Example 63

(S)-3-(4-glycidyloxybenzo(b)furan-2-yl)-1,5-dimethylpyrazole

1-(4-Methoxybenzo(b)furan-2-yl)butan-1,3-dione (1.0 g) was dissolved in methanol (30 ml) and methylhydrazine (0.3 g) was added thereto. The mixture was refluxed under heating for 20 min. The reaction solvent was evaporated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane/acetone). To the obtained oil was added methylene chloride (30 ml), and the mixture was cooled to -40°C, and boron tribromide (1 ml) was added dropwise. After the completion of the reaction, the mixture was poured into water and extracted with ethyl acetate. The organic layer was dried

over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure to give 3-(4-hydroxybenzo(b)furan-2-yl)-1,5-dimethylpyrazole (0.85 g) as a brown oil. Using this and (S)-glycidyl nosylate (0.75 g) and in the same manner as in Starting Material Synthesis Example 1, the title compound (0.53 g) was obtained as a brown oil.

¹H-NMR(CDCl₃): 2.33(s, 3H), 2.82(dd, 1H, J=2.8, 4.8), 2.94(t, 1H, J=4.4), 3.86(s, 3H), 4.13(dd, 1H, J=5.4, 11.2), 4.36(dd, 1H, J=3.4, 11.2), 6.40(s, 1H), 6.65(d, 1H, J=6.3), 7.06(s, 1H), 7.08-7.12(m, 2H)

Starting Material Synthesis Example 64

4-methoxymethylbenzo(b)thiophene-2-carboxylic acid

To a solution (400 ml) of 4,5,6,7-tetrahydrobenzo(b)thiophen-4-one (70.0 g) in methanol was added dropwise a solution (200 ml) of bromine (75.0 g) in methanol at room temperature. After the completion of the reaction, the reaction mixture was poured into water and extracted with chloroform, and the solvent was evaporated under reduced pressure. The obtained residue was dissolved in DMF (500 ml) and lithium bromide (30.0 g) was added. The reaction mixture was stirred with heating at a reaction temperature of 110°C for 1 hr. The reaction mixture was poured into ice water, extracted with chloroform, the extract was dried over magnesium sulfate and concentrated under reduced pressure to give 4-hydroxymethylbenzo(b)thiophene (50.5 g). This was dissolved in DMF (300 ml) and boron hydride (15.0 g) was added. The mixture was stirred at room temperature for 1 hr. To this reaction mixture was added dropwise methoxymethyl chloride (16.5 g). After the completion of the reaction, the reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was dried and concentrated under reduced pressure. The residue was dissolved in THF (400 ml) and n-BuLi (1.6M hexane solution) (250 ml) was added dropwise at -78°C. After the mixture was stirred for 30 min, and carbonic acid gas was

blown in until the reaction ended. The reaction mixture was poured into water, and the aqueous layer was made acidic with hydrochloric acid, and after extraction with ethyl acetate, the solvent was evaporated under reduced pressure to give the title
5 compound as white crystals, melting point 212-214°C.

Starting Material Synthesis Example 65

2-(7-hydroxybenzo(b)furan-2-yl)-5-methyloxazole

7-Methoxybenzo(b)furan-2-carboxylic acid (6.0 g) was dissolved in chloroform (30 ml), and dimethylformamide (1 ml)
10 was added. Thionyl chloride (4.0 ml) was added, and the mixture was stirred with heating at 50°C for 2 hr. The reaction solvent was evaporated under reduced pressure, and tetrahydrofuran (100 ml) was added. The mixture was cooled and a solution of propargylamine (1.65 g) and triethylamine (12 ml)
15 in tetrahydrofuran was added dropwise with stirring. The mixture was stirred at room temperature for 2 hr, and poured into water and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure.
20 This product (4 g) was dissolved in acetic acid (40 ml) and mercury(II) acetate (0.5 g) was added. The mixture was refluxed for 2 hr. After cooling, acetic acid was evaporated under reduced pressure and aqueous potassium carbonate solution was added, and the mixture was extracted with ethyl acetate.
25 The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure to give pale-yellow crystals (1.5 g). The crystals were dissolved in methylene chloride (30 ml), and the mixture was cooled to -20°C. Boron tribromide (0.8 ml) was added
30 dropwise and the mixture was stirred at 0°C for 1 hr. The reaction mixture was poured into water, and extracted with tetrahydrofuran. The organic layer was dried over anhydrous magnesium sulfate, and after filtration, the solvent was evaporated under reduced pressure to give the title compound

(1.0).

$^1\text{H-NMR}$ (DMSO-d_6): 2.42 (s, 3H), 6.92-6.95 (m, 2H), 7.01-7.13 (m, 1H), 7.18-7.35 (m, 1H), 7.63 (d, 1H, $J=2.8$)

Starting Material Synthesis Example 66

5 5-(7-methoxybenzo(b)furan-2-yl)-3-methylisoxazole

Thionyl chloride (10 ml) was added dropwise to methanol (100 ml) with stirring under ice-cooling. 7-Methoxybenzo(b)furan-2-carboxylic acid (10 g) was successively added, and the mixture was refluxed under heating for 1 hr.

10 After cooling, the solvent was evaporated under reduced pressure and the precipitated yellow crystals were collected by filtration to give methyl 7-methoxybenzo(b)furan-2-carboxylate (11.2 g). This was used in the next reaction without purification. Acetone oxime (4.8 g) was dissolved in

15 tetrahydrofuran (100 ml), and butyllithium (1.6M hexane solution) (80 ml) was added dropwise to this solution at -5°C with stirring. Thereafter, the mixture was stirred under ice-cooling for 1 hr, and a solution (50 ml) of methyl 7-methoxybenzo(b)furan-2-carboxylate (11.2 g) in tetrahydrofuran

20 was added. The mixture was stirred at room temperature for 20 hr. A solution of sulfuric acid (28 g) dissolved in tetrahydrofuran (120 ml) - water (30 ml) was prepared, into which the reaction mixture was poured. The mixture was refluxed under heating for 2 hr. After cooling, the reaction

25 mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.1 g).

30 $^1\text{H-NMR}$ (CDCl_3) δ : 2.38 (s, 3H), 4.04 (s, 3H), 6.57 (s, 1H), 6.88 (d, $J=7.8$, 1H), 7.22 (m, 3H)

Starting Material Synthesis Example 67

4-(4-methoxybenzo(b)furan-2-yl)-2-methylthiazole

To a solution (30 ml) of 4-methoxybenzo(b)furan-2-yl α -bromomethyl ketone (2.7 g) in ethanol was added thioacetamide (0.75 g), and the mixture was refluxed under heating for 6 hr. The precipitated crystals were collected by filtration and
5 dried to give the title compound (2.7 g) as pale-brown crystals.
 $^1\text{H-NMR}$ (DMSO- d_6) δ : 2.72(s, 3H), 3.91(s, 3H), 6.81(d, J=7.3, 1H), 7.13(s, 1H), 7.21(d, J=7.3, 1H), 7.27(t, J=7.3, 1H), 7.90(s, 1H)

Starting Material Synthesis Example 68

10 2-(2'-hydroxystyryl)-5-methyl-1,3,4-oxadiazole

2-(Methoxymethyloxy)cinnamic acid (4.0 g) and CDI (3.1 g) were successively added to tetrahydrofuran (40 ml) and the mixture was stirred. One hour later, acetylhydrazide (1.4 g) was added, and the mixture was stirred for 3 more hr. The
15 reaction mixture was poured into water and extracted with ethyl acetate to give an oil (3.5 g). This oil was dissolved in dichloroethane (300 ml) and triphenylphosphine (5 g) and triethylamine (3.3 ml) were added to this solution. Then DEAD (8.3 g) was added under ice-cooling. The mixture was stirred
20 at room temperature for 2 hr, and aqueous potassium carbonate solution was added, and reaction mixture was extracted with chloroform. The organic solvent was dried and concentrated, and the residue was purified by silica gel column chromatography (hexane/acetone) to give an oil (2.2 g). This
25 oil was stirred with heating in a mixed solvent of water (20 ml) and hydrochloric acid (20 ml) for 2 hr, and after cooling, poured into water. The mixture was extracted with ethyl acetate to give the title compound (1.5 g) as a brown oil.
 $^1\text{H-NMR}$ (CDCl_3): 2.58(s, 3H), 6.45(bs, 1H), 6.90(t, J=7.8, 1H),
30 6.98(d, J=7.5, 1H), 7.19(d, J=7.5, 1H), 7.40(t, J=8.0, 1H), 7.42(d, J=15.8, 1H), 7.68(d, J=15.8, 1H)

Starting Material Synthesis Example 69

2-(2'-hydroxystyryl)benzothiazole

Salicylaldehyde (6.1 g) and 2-methylthiazole (7.5 g)

were mixed and conc. hydrochloric acid (1.5 ml) was added thereto. The mixture was stirred with heating at 100°C for 9 hr. The reaction mixture was cooled, and aqueous potassium hydroxide solution was added. The aqueous layer was washed
5 with ether and made acidic with hydrochloric acid and extracted again with ethyl acetate. The organic solvent was dried and concentrated to give the title compound (2.5 g) as pale-yellow crystals, melting point 235-236°C.

Starting Material Synthesis Example 70

10 5-(2'-hydroxystyryl)-3-methyl-1,2,4-oxadiazole

Acetamide oxime (7.5 g), molecular sieves (4A) (10 g) and sodium hydride (5 g) were added to tetrahydrofuran (200 ml) and the mixture was refluxed under heating. To this reaction mixture was added dropwise ethyl 2-(methoxymethoxy)cinnamate
15 (12 g) and the mixture was continuously heated for 2 hr. After cooling, the mixture was poured on ice and extracted with ethyl acetate. The organic layer was concentrated under reduced pressure. Thereto were added tetrahydrofuran (10 ml) and 6N hydrochloric acid (20 ml) and the mixture was stirred with
20 heating at 50°C for 30 min to allow precipitation of crystals. The crystals were collected by filtration and dried to give the title compound (6.0 g) as white crystals, melting point 184-186°C.

Starting Material Synthesis Example 71

25 (S)-(4-glycidyloxy)benzo(b)furan-2-yl methyl ketone

To a suspension (40 ml) of sodium hydride (0.22 g) in DMF was added dropwise a solution (10 ml) of 4-hydroxybenzo(b)furan-2-yl methyl ketone (0.80 g) in DMF under ice-cooling and the mixture was stirred at room temperature for
30 30 min. To this reaction mixture was added dropwise a solution (10 ml) of (S)-glycidyl nosylate (1.4 g) in DMF under ice-cooling, and the mixture was stirred for 2 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with water and dried

over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (0.61 g) as yellow crystals.

5 ¹H-NMR(CDCl₃): 2.60 (s, 3H), 2.82 (dd, J=4.4, 5.9, 1H), 2.97 (t, J=4.4, 1H), 3.43-3.46 (m, 1H), 4.09 (dd, J=10.8, 5.9, 1H), 4.42 (dd, J=10.8, 3.0, 1H), 6.69 (d, J=7.8, 1H), 7.20 (d, J=8.3, 1H), 7.39 (t, J=8.3, 1H), 7.65 (s, 1H)

Starting Material Synthesis Example 72


10 (S)-4-glycidyloxy-3-methylbenzo(b)furan-2-yl methyl ketone

To a suspension (60 ml) of sodium hydride (1.4 g) in DMF was added dropwise a solution (30 ml) of 4-hydroxy-3-methylbenzo(b)furan-2-yl methyl ketone (6.1 g) in DMF under ice-cooling and the mixture was stirred at room temperature for
15 30 min. To this reaction mixture was added dropwise under ice-cooling a solution (30 ml) of (S)-glycidyl nosylate (9.1 g) in DMF, and the mixture was stirred for 2 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with water and dried
20 over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.1 g) as pale-yellow crystals.


¹H-NMR(CDCl₃): 2.59 (s, 3H), 2.79 (s, 3H), 2.83 (dd, J=4.9, 2.3, 1H), 2.96 (t, J=4.3, 1H), 3.43-3.45 (m, 1H), 4.08 (dd, J=11.2, 5.4, 1H), 4.37 (dd, J=11.2, 3.0, 1H), 6.62 (d, J=7.8, 1H), 7.11 (d, J=8.3, 1H), 7.34 (t, J=8.3, 1H)

The structural formulas of the compounds obtained in Starting Material Synthesis Examples 54 to 72 are shown in the
30 following.


54




55




56




57




58




59




60




61




62




63




64




65




66




67




68




69




70



71



72



Starting Material Synthesis Example 73

N'-(4-methoxybenzo(b)furan-2-ylcarbonyl)propionohydrazide

To a solution (200 ml) of (4-methoxybenzo(b)furan-2-ylcarbonyl)hydrazide (8.5 g) in THF was added propionic anhydride (8.1 g) and the mixture was stirred at room temperature for 1 hr. The reaction mixture was concentrated under reduced pressure and the residue was crystallized from diisopropyl ether, collected by filtration and dried to give the title compound (8.3 g) as brown crystals.

¹H-NMR(DMSO-d₆)δ:1.05(t, J=7.8,3H), 2.19(q, J=7.8,2H), 3.93(s, 3H), 6.86(d, J=7.8, 1H), 7.25(d, J=8.3, 1H), 7.41(t, J=8.3, 1H), 7.62(s, 1H), 9.89(s, 1H), 10.46(s, 1H)

Starting Material Synthesis Example 74

2-(4-methoxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole

N'-(4-Methoxybenzo(b)furan-2-ylcarbonyl)-propionohydrazide (8.3 g) obtained in Starting Material Synthesis Example 73 was added to phosphorus oxychloride (60 ml) and the mixture was stirred at 90°C for 1 hr. After cooling, the reaction mixture was poured into ice water and extracted with ethyl acetate. After washing with water, the organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to give the title compound (4.5 g) as yellow crystals.

¹H-NMR(CDCl₃)δ:1.46(t, J=7.8,3H), 2.99(q, J=7.8,2H), 3.97(s, 3H), 6.72(d, J=7.8, 1H), 7.22(d, J=8.3, 1H), 7.36(t, J=8.3, 1H), 7.57(s, 1H)

Starting Material Synthesis Example 75

2-(4-hydroxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole

To a solution (60 ml) of 2-(4-methoxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole (4.5 g) obtained in Starting Material Synthesis Example 74 in methylene chloride was added boron tribromide (11.8 ml), and the mixture was stirred at room temperature for 2 hr. The reaction mixture was poured into ice water and stirred for 1 hr and extracted with a mixed solvent

of chloroform - methanol (2:1). After washing with water, the organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to give the title compound (3.1 g) as pale-yellow crystals.

- 5 $^1\text{H-NMR}$ (DMSO- d_6) δ : 1.33 (t, $J=7.8$, 3H), 2.96 (q, $J=7.8$, 2H), 6.71 (d, $J=8.3$, 1H), 7.16 (d, $J=8.8$, 1H), 7.29 (t, $J=8.3$, 1H), 7.69 (s, 1H), 10.37 (s, 1H)

Starting Material Synthesis Example 76

(S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-

10 oxadiazole

- By the reactions in the same manner as in Starting Material Synthesis Example 1 using 2-(4-hydroxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole (3.1 g) obtained in Starting Material Synthesis Example 75, (S)-glycidyl nosylate (3.5 g) and potassium carbonate (5.6 g), the title compound (3.8 g) was obtained as pale-yellow crystals.

- $^1\text{H-NMR}$ (CDCl₃) δ : 1.47 (t, $J=7.8$, 3H), 2.83 (dd, $J=3.9$, 2.4, 1H), 2.96 (t, $J=3.9$, 1H), 2.99 (q, $J=7.8$, 2H), 3.42-3.48 (m, 1H), 4.11 (dd, $J=11.3$, 5.9, 1H), 4.42 (dd, $J=11.3$, 3.0, 1H), 6.72 (d, $J=8.3$, 1H), 7.25 (d, $J=8.3$, 1H), 7.32 (t, $J=8.3$, 1H), 7.61 (s, 1H)

Starting Material Synthesis Example 77

5-(4-methoxybenzo(b)furan-2-yl)-3-methylisoxazole

- To a solution (160 ml) of acetone oxime (5.0 g) in THF was added dropwise n-butyllithium (1.6 M hexane solution) over 15 min under ice-cooling and the mixture was stirred for 1 hr. Thereto was added dropwise a solution (60 ml) of methyl 4-methoxybenzo(b)furan-2-carboxylate (6.7 g) in THF and the mixture was stirred at room temperature for 2 hr. The reaction mixture was poured into ice water, and conc. sulfuric acid (4 ml) was added carefully. The mixture was stirred for 20 min more. The aqueous layer was neutralized with sodium hydrogencarbonate and extracted with ethyl acetate. After washing with water, the organic layer was dried over anhydrous magnesium sulfate and the solvent was evaporated under reduced

pressure to give the title compound (3.0 g) as yellow crystals.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.38(s, 3H), 3.96(s, 3H), 6.46(s, 1H), 6.69(d, J=7.8, 1H), 7.15(d, J=8.3, 1H), 7.29(t, J=8.3, 1H), 7.32(s, 1H)

Starting Material Synthesis Example 78

5 5-(4-hydroxybenzo(b)furan-2-yl)-3-methylisoxazole

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 5-(4-methoxybenzo(b)furan-2-yl)-3-methylisoxazole (3.0 g) and boron tribromide (7.6 ml), the title compound (2.6 g) was obtained as pale-yellow crystals.

10 $^1\text{H-NMR}(\text{DMSO}-d_6)\delta$: 2.30(s, 3H), 6.68(d, J=7.8, 1H), 6.85(s, 1H), 7.10(d, J=8.3, 1H), 7.22(t, J=8.3, 1H), 7.49(s, 1H)

Starting Material Synthesis Example 79

(S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-3-methylisoxazole

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 5-(4-hydroxybenzo(b)furan-2-yl)-3-methylisoxazole (2.6 g), (S)-glycidyl nosylate (3.1 g) and potassium carbonate (5.0 g), the title compound (2.8 g) was obtained as brown crystals.

20 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.82(dd, J=4.9, 2.4, 1H), 2.96(t, J=4.9, 1H), 3.43-3.46(m, 1H), 4.11(dd, J=11.2, 5.4, 1H), 4.39(dd, J=11.2, 3.0, 1H), 6.49(s, 1H), 6.70(d, J=8.3, 1H), 7.17(d, J=8.3, 1H), 7.28(t, J=8.3, 1H), 7.36(s, 1H)

Starting Material Synthesis Example 80

2-(4-methoxybenzo(b)furan-2-yl)-5-methyl-1,3,4-thiadiazole

25 To a solution (50 ml) of N'-(4-methoxybenzo(b)furan-2-ylthiocarbonyl)acetohydrazide (1.1 g) in toluene was added methanesulfonic acid (1.0 ml) and the mixture was stirred at 80°C for 30 min. The reaction mixture was concentrated under reduced pressure and the residue was dissolved in ethyl acetate and aqueous potassium carbonate solution, followed by partitioning. After washing with water, the organic layer was dried over anhydrous magnesium sulfate and the solvent was evaporated under reduced pressure to give the title compound (0.82 g) as yellow crystals.

¹H-NMR(CDCl₃)δ:2.85(s, 3H), 3.97(s, 3H), 6.70(d, J=7.8, 1H), 7.18(d, J=8.3, 1H), 7.32(t, J=8.3, 1H), 7.57(s, 1H)

Starting Material Synthesis Example 81

2-(4-hydroxybenzo(b)furan-2-yl)-5-methyl-1,3,4-thiadiazole

5 By the reactions in the same manner as in Starting Material Synthesis Example 5 using 2-(4-methoxybenzo(b)furan-2-yl)-5-methyl-1,3,4-thiadiazole (0.98 g) and boron tribromide (2.3 ml), the title compound (0.89 g) was obtained as pale-yellow crystals.

10 ¹H-NMR(DMSO-d₆)δ:2.80(s, 3H), 6.70(d, J=7.3, 1H), 7.13(d, J=8.3, 1H), 7.25(t, J=8.3, 1H), 7.67(s, 1H)

Starting Material Synthesis Example 82

(S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-thiadiazole

15 By the reactions in the same manner as in Starting Material Synthesis Example 1 using 2-(4-hydroxybenzo(b)furan-2-yl)-5-methyl-1,3,4-thiadiazole (1.1 g), (S)-glycidyl nosylate (1.2 g) and potassium carbonate (3.0 g), the title compound (1.0 g) was obtained as yellow crystals.

20 ¹H-NMR(CDCl₃)δ:2.82(dd, J=4.9, 3.0, 1H), 2.96(t, J=4.9, 1H), 3.42-3.46(m, 1H), 4.13(dd, J=10.8, 5.9, 1H), 4.40(dd, J=10.8, 3.0, 1H), 6.71(d, J=7.8, 1H), 7.20(d, J=8.3, 1H), 7.31(t, J=8.3, 1H), 7.61(s, 1H)

Starting Material Synthesis Example 83

25 N-propargyl-4-methoxybenzo(b)furan-2-carboxamide

4-Methoxybenzo(b)furan-2-carboxylic acid (44.0 g) and propargylamine (12 g) were dissolved in dimethylformamide (200 ml), and WSC (48.0 g), HOBt (43.0 g) and triethylamine (50 ml) were added thereto at room temperature. The mixture was
30 stirred for 4 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous solution of ammonium chloride, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound as yellow crystals

(45.0 g).

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 3.32(s, 1H), 3.92(s, 3H), 4.06(m, 2H), 6.65(d, $J=7.8$, 1H), 7.18(d, $J=7.8$, 1H), 7.26(t, $J=7.8$, 1H), 7.36(s, 1H), 8.86(m, 1H)

5 **Starting Material Synthesis Example 84**

4-methoxy-2-(5-methyl-1,3-oxazol-2-yl)benzo(b)furan

To a solution (200 ml) of N-propargyl-4-methoxybenzo(b)furan-2-carboxamide (45.0 g) obtained in Starting Material Synthesis Example 83 in acetic acid was added
10 mercury acetate (7.0 g), and the mixture was refluxed under heating for 3 hr. After cooling, the solvent was evaporated under reduced pressure and water was added. The mixture was neutralized with potassium carbonate and extracted with ethyl acetate. The solvent was evaporated under reduced pressure and
15 the residue was purified by silica gel column chromatography (chloroform) to give the title compound (15.0 g) as yellow crystals.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.42(s, 3H), 3.96(s, 3H), 6.70(d, $J=7.8$, 1H), 6.90(s, 1H), 7.20(d, $J=7.8$, 1H), 7.29(t, $J=7.8$, 1H), 7.38(s,
20 1H)


Starting Material Synthesis Example 85

2-(4-hydroxybenzo(b)furan-2-yl)-5-methyloxazole

To a solution (100 ml) of 4-methoxy-2-(5-methyl-1,3-oxazol-2-yl)benzo(b)furan (15.0 g) obtained in Starting
25 Material Synthesis Example 84 in dichloromethane was added dropwise boron tribromide (14 ml) under ice-cooling. The mixture was stirred at room temperature for 3 hr and poured into ice water. The mixture was stirred at room temperature for 3 more hr. The crystals were collected by filtration and
30 dissolved in ethyl acetate. 1N HCl was added and the mixture was stirred one day. The organic layer was washed with saturated aqueous solution of sodium hydrogencarbonate, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (11.0 g) as yellow crystals.

$^1\text{H-NMR}$ (DMSO-d_6) δ : 2.41 (s, 3H), 6.68 (d, $J=7.8$, 1H), 7.04 (s, 1H), 7.10 (d, $J=7.8$, 1H), 7.21 (t, $J=7.8$, 1H), 7.45 (s, 1H), 10.17 (bs, 1H)

The structural formulas of the compounds obtained in
 5 Starting Material Synthesis Examples 73 to 85 are shown in the following.



Starting Material Synthesis Example 86

4-methoxy-2-(trimethylstannyl)benzo(b)furan

To a solution of 4-methoxybenzo(b)furan (2.50 g) in THF (50 ml) was added n-butyllithium (1.54 M hexane solution) (16.5 ml) at -78°C and the mixture was stirred at the same temperature for 20 min. To this solution was added trimethyltin chloride (5.00 g) and the mixture was further stirred at the same temperature for 1 hr. The reaction mixture was warmed to room temperature and water (200 ml) was added. The mixture was extracted with ethyl acetate, and the obtained organic layer was washed with water and saturated brine and dried over magnesium sulfate. This solution was concentrated under reduced pressure to give the title compound (5.86 g) as pale-yellow crystals.

¹H-NMR(CDCl₃) δ: 0.35(s, 3H), 3.90(s, 3H), 6.58(d, J=5.0, 1H), 6.98(s, 1H), 7.10-7.40(m, 2H)

Starting Material Synthesis Example 87

2-(5-ethylthiophen-2-yl)-4-methoxybenzo(b)furan

To a solution of 4-methoxy-2-(trimethylstannyl)-benzo(b)furan (3.00 g) and 2-bromo-5-ethylthiophene (1.84 g) in THF (25 ml) was added bistriphenylphosphinepalladium dichloride (224 mg), and the mixture was stirred with refluxing overnight. After cooling, ethyl acetate was added to the reaction mixture and the mixture was filtered through celite. The filtrate was washed with water and saturated brine, and the organic layer was dried over magnesium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography to give the title compound (1.04 g) as a yellow oil.

¹H-NMR(CDCl₃) δ: 1.32(t, J=8.0, 3H), 2.84(q, J=8, 2H), 3.93(s, 3H), 6.60-6.65(m, 2H), 6.75(d, J=2.0, 1H), 6.85(s, 1H), 7.15(d, J=7.0, 1H), 7.18-7.23(m, 2H)

Starting Material Synthesis Example 88

2-(5-ethylthiophen-2-yl)-4-hydroxybenzo(b)furan

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 2-(5-ethylthiophen-2-yl)-4-methoxybenzo(b)furan (1.00 g) and boron tribromide (1.0 ml), the title compound (774 mg) was obtained as colorless crystals, melting point 87-90°C.

Starting Material Synthesis Example 89

(S)-2-(5-ethylthiophen-2-yl)-4-glycidyloxybenzo(b)furan

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 2-(5-ethylthiophen-2-yl)-4-hydroxybenzo(b)furan (750 mg), (S)-glycidyl nosylate (875 mg) and potassium carbonate (1.27 g), a crude product of the title compound was quantitatively obtained as a yellow oil.

¹H-NMR(CDCl₃) δ: 1.32 (t, J=8.0, 3H), 2.83 (dd, J=3.9, 2.4, 1H), 2.96 (t, J=3.9, 1H), 2.75-2.95 (m, 2H), 3.35-3.45 (m, 1H), 4.05 (dd, J=11.0, 6.0, 1H), 4.34 (dd, J=11.0, 3.0, 1H), 6.62 (dd, J=8.0, 1.0, 1H), 6.75 (d, J=1.0, 1H), 6.88 (s, 1H), 7.10-7.20 (m, 2H)

Starting Material Synthesis Example 90

4-methoxy-2-(1-methylimidazol-2-yl)benzo(b)furan

By the reactions in the same manner as in Starting Material Synthesis Example 87 using 4-methoxy-2-(trimethylstannyl)benzo(b)furan (5.26 g), 2-bromo-1-methylimidazole (2.72 g) and bistriphenylphosphinepalladium dichloride (593 mg), the title compound (2.06 g) was obtained as a yellow oil.

¹H-NMR(CDCl₃) δ: 3.97 (s, 6H), 6.69 (d, J=8.0, 1H), 7.22 (d, J=8.0, 1H), 6.96 (s, 1H), 7.24 (d, J=8.0, 1H), 7.20-7.35 (m, 4H)

Starting Material Synthesis Example 91

4-hydroxy-2-(1-methylimidazol-2-yl)benzo(b)furan

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 4-methoxy-2-(1-methylimidazol-2-yl)benzo(b)furan (2.00 g) and boron tribromide (2.0 ml), the title compound (1.21 g) was obtained as pale-yellow crystals, melting point >265°C (decomposition).

Starting Material Synthesis Example 92

(S)-4-glycidyloxy-2-(1-methylimidazol-2-yl)benzo(b)furan

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 4-hydroxy-2-(1-methylimidazol-2-yl)benzo(b)furan (1.10 g), (S)-glycidyl nosylate (1.33 g) and potassium carbonate (2.13 g), a crude product of the title compound was quantitatively obtained as a pale-yellow oil.

¹H-NMR(CDCl₃)δ: 2.75-2.80 (m, 1H), 2.92 (d, J=4.0, 2H), 3.42-3.45 (m, 1H), 4.11 (dd, J=11.0, 6.0, 1H), 4.37 (dd, J=11.0, 2.0, 1H), 6.67 (d, J=8.0, 1H), 6.95 (s, 1H), 7.10-7.30 (m, 4H)

Starting Material Synthesis Example 93

N-propargyl-4-(methoxymethyloxy)benzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Starting Material Synthesis Example 83 using 4-(methoxymethyloxy)-benzo(b)furan-2-carboxylic acid (10.0 g), propargylamine (2.31 g), WSC (8.87 g), HOBt (7.71 g) and triethylamine (8.76 ml), the title compound (7.15 g) was obtained as pale-brown crystals.

¹H-NMR(CDCl₃)δ: 3.31 (s, 1H), 3.45 (s, 3H), 4.06 (br.s, 2H), 5.38 (s, 2H), 7.04 (d, J=8.0, 1H), 7.38 (t, J=8.0, 1H), 7.60 (d, J=8.0, 1H), 8.25 (s, 1H), 9.26 (m, 1H)

Starting Material Synthesis Example 94

4-hydroxy-2-(5-methyloxazol-2-yl)benzo(b)thiophene

By the reactions in the same manner as in Starting Material Synthesis Example 5 using N-propargyl-4-(methoxymethyloxy)benzo(b)thiophene-2-carboxamide (6.00 g) and mercury acetate (765 mg), the title compound (2.41 g) was obtained as yellow crystals, melting point 188-189°C.

Starting Material Synthesis Example 95

(S)-4-glycidyloxy-2-(5-methyloxazol-2-yl)benzo(b)thiophene

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 4-hydroxy-2-(5-methyloxazol-2-yl)benzo(b)thiophene (2.20 g), (S)-glycidyl nosylate (2.38 g) and potassium carbonate (3.18 g), a crude product of the title

compound was quantitatively obtained as pale-yellow crystals.
 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.84(dd, $J=5.0, 3.0$, 1H), 2.96(t, $J=5.0$, 1H),
3.40-3.48(m, 1H), 4.12(dd, $J=11.0, 6.0$, 1H), 4.39(dd, $J=11.0$,
2.0, 1H), 6.76(d, $J=8.0$, 1H), 6.85(s, 1H), 7.33(d, $J=8.0$, 1H),
5 7.45(d, $J=8.0$, 1H), 8.03(s, 1H)

Starting Material Synthesis Example 96

2-(4,4-dimethyloxazolin-2-yl)-4-methoxybenzo(b) furan

To a solution of 4-methoxybenzo(b)furan-2-carboxylic
acid (15.0 g) in dichloromethane (300 ml) were added DMF (6 ml)
10 and thionyl chloride (17.1 ml), and the mixture was stirred
with refluxing for 2 hr. The solvent was evaporated under
reduced pressure to give acid chloride.

This acid chloride was dissolved in dichloromethane (150
ml) and added dropwise to a solution of 2-amino-2-methyl-1-
15 propanol (10.4 g) in dichloromethane (150 ml) at 0°C . The
mixture was stirred at room temperature for 2 hr and saturated
sodium hydrogencarbonate (500 ml) was added. The mixture was
extracted with chloroform and the organic layer was washed with
water and saturated brine and dried over magnesium sulfate.
20 The solvent was evaporated under reduced pressure and the
residue was purified by silica gel column chromatography to
give an amide compound (14.7 g).













To the obtained amide compound was added thionyl
chloride (75 ml) and the mixture was stirred at room
25 temperature for 5 hr and poured into saturated aqueous solution
of sodium hydrogencarbonate (500 ml). 10% Sodium hydroxide was
added to this mixed solution until it reached pH 12 and the
mixture was extracted with ethyl acetate. The organic layer
was washed with water and saturated brine and dried over
30 magnesium sulfate. The solvent was evaporated under reduced
pressure to give the title compound (7.81 g) as a yellow oil.
 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.40(s, 6H), 3.92(s, 3H), 4.12(s, 2H), 6.65(d,
 $J=8.0$, 1H), 7.15(d, $J=8.0$, 1H), 7.28(t, $J=8.0$, 1H), 7.33(s, 1H)

Starting Material Synthesis Example 97

2-(4,4-dimethyloxazolin-2-yl)-4-hydroxybenzo(b) furan

By the reactions in the same manner as in Starting Material Synthesis Example 5 using 2-(4,4-dimethyloxazolin-2-yl)-4-methoxybenzo(b) furan (6.11 g) and boron tribromide (6.11 ml), the title compound (5.03 g) was obtained as pale-yellow crystals, melting point 187-188°C.

The structural formulas of the compounds obtained in Starting Material Synthesis Examples 86 to 97 are shown in the following.



Starting Material Synthesis Example 98

(S)-4-glycidyoxy-2-(4,4-dimethyloxazolin-2-yl)benzo(b)furan

By the reactions in the same manner as in Starting Material Synthesis Example 1 using 2-(4,4-dimethyloxazolin-2-yl)-4-hydroxybenzo(b)furan (1.50 g), (S)-glycidyl nosylate (1.68 g) and potassium carbonate (2.69 g), a crude product of the title compound was quantitatively obtained as a pale-yellow oil.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.40 (s, 6H), 2.75-2.80 (m, 1H), 2.92 (t, J=4.0, 1H), 3.35-3.45 (m, 1H), 4.00-4.20 (m, 1H), 4.13 (s, 2H), 4.36 (dd, J=11.0, 2.0, 1H), 6.65 (d, J=8.0, 1H), 7.18 (d, J=8.0, 1H), 7.28 (t, J=8.0, 1H), 7.37 (s, 1H)

Starting Material Synthesis Example 99

2-(ethylsulfonyl)-4-methoxybenzo(b)furan

To a solution of 4-methoxybenzo(b)furan (5.00 g) in THF (40 ml) was added n-butyllithium (1.54 M hexane solution) (24.1 ml) at -78°C , and the mixture was stirred at the same temperature for 30 min. To this solution was added sulfur (1.19 g) and the mixture was stirred further at the same temperature for 30 min. Then, bromoethane (4.16 ml) was added and this reaction mixture was stirred at room temperature for 1 hr. A saturated aqueous solution of ammonium chloride (100 ml) was added, and the mixture was extracted with ethyl acetate. The obtained organic layer was washed with water and saturated brine and dried over magnesium sulfate. This solution was concentrated under reduced pressure to give a sulfide compound (3.50 g).

To a solution of this sulfide compound (3.50 g) in dichloromethane (50 ml) was added m-chloroperoxybenzoic acid (70%, 9.13 g) at 0°C , and the mixture was stirred at room temperature for 1 hr. Saturated sodium hydrogencarbonate (50 ml) and saturated sodium thiosulfate (50 ml) were added to the reaction mixture and the mixture was extracted with ethyl acetate. The obtained organic layer was washed with water and

saturated brine and dried over magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography to give the title compound (3.53 g) as pale-brown crystals.

- 5 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.34 (t, $J=8.0$, 1H), 3.29 (q, $J=8.0$, 1H), 3.94 (s, 3H), 6.72 (d, $J=8.0$, 1H), 7.16 (d, $J=8$, 1H), 7.40 (t, $J=8.0$, 1H), 7.61 (s, 1H)

Starting Material Synthesis Example 100

2-(ethylsulfonyl)-4-hydroxybenzo(b) furan

- 10 By the reactions in the same manner as in Starting Material Synthesis Example 5 using 2-(ethylsulfonyl)-4-methoxybenzo(b) furan (3.50 g) and boron tribromide (7.0 ml), the title compound (2.85 g) was obtained as colorless crystals, melting point 145-147°C.

Starting Material Synthesis Example 101

(S)-2-(ethylsulfonyl)-4-glycidyloxybenzo(b) furan

- By the reactions in the same manner as in Starting Material Synthesis Example 1 using 2-(ethylsulfonyl)-4-hydroxybenzo(b) furan (2.75 g), (S)-glycidyl nosylate (3.48 g) and potassium carbonate (5.05 g), a crude product of the title compound (3.62 g) was obtained as a pale-yellow oil.

- 20 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.34 (t, $J=8.0$, 1H), 2.79 (dd, $J=4.0$, 2.0, 2H), 2.94 (t, $J=4.0$, 1H), 3.29 (q, $J=8.0$, 1H), 3.35-3.45 (m, 1H), 4.08 (dd, $J=10.0$, 4.0, 2H), 4.39 (dd, $J=10.0$, 2.0, 1H), 6.72 (d, $J=8.0$, 1H), 7.18 (d, $J=8.0$, 1H), 7.39 (t, $J=8.0$, 1H), 7.65 (s, 1H)

Starting Material Synthesis Example 102

2-(N,N-dimethylsulfamoyl)-4-methoxybenzo(b) furan

- To a solution of 4-methoxybenzo(b) furan (5.00 g) in THF (40 ml) was added n-butyllithium (1.54 M hexane solution, 24.1 ml) at -78°C and the mixture was stirred at the same temperature for 30 min. To this solution was added sulfonyl chloride (9.13 g) and the mixture was stirred further at 0°C for 1 hr. The reaction mixture was concentrated under reduced pressure and the condensate was dissolved in acetone (30 ml).

This was added dropwise at room temperature to a mixed solution of aqueous dimethylamine solution (50%, 20 g) and acetone (50 ml), and the mixture was stirred at room temperature for 1 hr and extracted with ethyl acetate. The obtained organic layer
5 was washed with water and saturated brine and dried over magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography to give the title compound (1.32 g) as a yellow oil.

10 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.88(s, 6H), 3.94(s, 3H), 6.71(d, J=8.0, 1H), 7.14(d, J=8.0, 1H), 7.35(t, J=8.0, 1H), 7.45(s, 1H)

Starting Material Synthesis Example 103

2-(N,N-dimethylsulfamoyl)-4-hydroxybenzo(b)furan

By the reactions in the same manner as in Starting
15 Material Synthesis Example 5 using 2-(N,N-dimethylsulfamoyl)-4-methoxybenzo(b)furan (1.30 g) and boron tribromide (2.6 ml), the title compound (1.20 g) was obtained as colorless crystals, melting point 150-153°C.

Starting Material Synthesis Example 104

20 (S)-2-(N,N-dimethylsulfamoyl)-4-glycidyloxybenzo(b)furan

By the reactions in the same manner as in Starting
Material Synthesis Example 1 using 2-(N,N-dimethylsulfamoyl)-4-methoxybenzo(b)furan (1.10 g), (S)-glycidyl nosylate (1.30 g) and potassium carbonate (1.89 g), a crude product of the title
25 compound was quantitatively obtained as a pale-yellow oil.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.80(dd, J=4.0, 1.0, 2H), 2.88(s, 6H), 2.94(dd, J=4.0, 1.0, 1H), 3.35-3.45(m, 1H), 4.07(dd, J=11.0, 4.0, 2H), 4.39(dd, J=11.0, 1.0, 1H), 6.71(d, J=8.0, 1H), 7.16(d, J=8.0, 1H), 7.35(t, J=8.0, 1H), 7.47(s, 1H)

30 **Starting Material Synthesis Example 105**

N-(2-oxobutyl)-4-methoxybenzo(b)furan-2-carboxamide

To a solution of 4-methoxybenzo(b)furan-2-carboxylic acid (10.0 g) in dichloromethane (100 ml) were added DMF (4 ml) and thionyl chloride (11.4 ml), and the mixture was stirred

with refluxing for 2 hr. The solvent was evaporated under reduced pressure to give acid chloride.

This acid chloride was dissolved in THF (50 ml) and added dropwise to a solution of 1-amino-2-butanol (10.0 g) in THF (130 ml) at 0°C. The mixture was stirred at room temperature for 3 hr and water (200 ml) was added. The mixture was extracted with ethyl acetate and the organic layer was washed with water and saturated brine and dried over magnesium sulfate. The solvent was evaporated under reduced pressure and purified by silica gel column chromatography to give an amide compound (7.48 g) as a brown oil.

A solution of the obtained amide compound (3.00 g) in dichloromethane (20 ml) was added dropwise to a suspension of pyridinium chlorochromate (7.39 g) and molecular sieve (4A) (7.50 g) in dichloromethane (120 ml) at room temperature, and the mixture was stirred at room temperature for 2 hr. Ether (300 ml) was added to the reaction mixture and the mixture was dried over magnesium sulfate and filtered through celite. The filtrate was concentrated under reduced pressure and purified by silica gel column chromatography (chloroform-methanol) to give the title compound (2.14 g) as a yellow oil.

¹H-NMR(CDCl₃) δ: 1.15(t, J=8.0, 3H), 2.54(q, J=8.0, 2H), 3.47(d, J=2.0, 1H), 3.94(s, 3H), 4.35(d, J=2.0, 2H), 6.66(d, J=8.0, 1H), 7.12(d, J=8.0, 1H), 7.32(t, J=8.0, 1H), 7.55(s, 1H)

Starting Material Synthesis Example 106

2-(5-ethyloxazol-2-yl)-4-methoxybenzo(b)furan

To a solution of N-(2-oxobutyl)-4-methoxybenzo(b)furan-2-carboxamide (2.00 g) in THF (60 ml) was added Burgess reagent (7.30 g), and the mixture was stirred with refluxing. Water (200 ml) was added to the reaction mixture and the mixture was extracted with ether. The organic layer was washed with water and saturated brine, and dried over magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography to give the

title compound (1.56 g) as colorless crystals, melting point 88-87°C.

Starting Material Synthesis Example 107

2-(5-ethyloxazol-2-yl)-4-hydroxybenzo(b)furan

5 By the reactions in the same manner as in Starting Material Synthesis Example 5 using 2-(5-ethyloxazol-2-yl)-4-methoxybenzo(b)furan (2.00 g) and boron tribromide (2.0 ml), the title compound (1.24 g) was obtained as colorless crystals. ¹H-NMR(CDCl₃)δ:1.33(t, J=8.0, 3H), 2.79(q, J=8.0, 2H),
10 6.04(br.s, 1H), 6.68(d, J=8.0, 1H), 6.92(s, 1H), 7.15(d, J=8.0, 1H), 7.21(t, J=8.0, 1H), 7.42(s, 1H)

Starting Material Synthesis Example 108

(S)-2-(5-ethyloxazol-2-yl)-4-glycidyloxybenzo(b)furan

By the reactions in the same manner as in Starting
15 Material Synthesis Example 1 using 2-(5-ethyloxazol-2-yl)-4-hydroxybenzo(b)furan (1.10 g), (S)-glycidyl nosylate (1.24 g) and potassium carbonate (1.99 g), a crude product of the title compound (1.36 g) was obtained as pale-yellow crystals. ¹H-NMR(CDCl₃)δ:1.34(t, J=8.0, 3H), 2.70-2.2.85(m, 3H), 2.96(t, J=3.0, 1H), 3.40-3.45(m, 1H), 4.14(dd, J=11.0, 4.0, 2H),
20 4.40(dd, J=11.0, 1.0, 1H), 6.70(d, J=8.0, 1H), 6.91(s, 1H), 7.20-7.30(m, 2H), 7.44(s, 1H)

Starting Material Synthesis Example 109

N'-(4-hydroxybenzo(b)furan-2-ylcarbonyl)acetohydrazide

25 N'-(4-Methoxybenzo(b)furan-2-ylcarbonyl)acetohydrazide (4.0 g) obtained in Starting Material Synthesis Example 36 was dissolved in dichloromethane (40 ml) and boron tribromide (4.0 ml) was added dropwise while stirring under ice-cooling. Then the mixture was stirred at room temperature for 5 hr, and the
30 reaction mixture was poured into ice water. The mixture was further stirred as it was at room temperature for 1 hr. The precipitated crystals were collected by filtration, dissolved in chloroform, washed with saturated aqueous solution of sodium hydrogencarbonate, dried over anhydrous sodium sulfate and

concentrated under reduced pressure to give the title compound (2.5 g) as yellow crystals.

¹H-NMR(DMSO-d₆)δ:1.93(s, 3H), 6.69(d, J=7.8, 1H), 7.07(d, J=7.8, 1H), 7.27(t, J=7.8, 1H), 7.65(s, 1H), 9.92(s, 1H), 10.29(s, 1H),
5 10.43(s, 1H)

Starting Material Synthesis Example 110

N-ethoxalyl-N'-(4-methoxybenzofuran-2-ylcarbonyl)hydrazide

4-Methoxybenzo(b)furan-2-ylcarbonylhydrazide (9.0 g) was dissolved in dichloromethane (100 ml) and triethylamine (7.0
10 ml) and ethyl chloroglyoxylate (6.0 g) were added. The mixture was stirred at room temperature for 4 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The
15 residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (7.0 g) as pale-yellow crystals.

¹H-NMR(DMSO-d₆)δ:1.33(t, J=8.0, 3H), 3.95(s, 3H), 4.32(q, J=8.0, 2H), 6.87(d, J=7.8, 1H), 7.26(d, J=7.8, 1H), 7.44(t, J=7.8, 1H),
20 7.65(s, 1H), 10.82(s, 1H), 10.96(s, 1H)

Starting Material Synthesis Example 111

5-ethoxycarbonyl-2-(4-methoxybenzo(b)furan-2-yl)-1,3,4-oxadiazole

N-Ethoxalyl-N'-(4-methoxybenzofuran-2-ylcarbonyl)-
25 hydrazide (7.0 g) was dissolved in phosphorus oxychloride (20 ml) and the mixture was heated at 80°C for 7 hr. Phosphorus oxychloride was evaporated under reduced pressure and water was added to the residue. The mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous
30 solution of sodium hydrogencarbonate, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (chloroform) to give the title compound (5.0 g) as yellow crystals.

¹H-NMR(CDCl₃)δ:1.49(t, J=8.0, 3H), 3.98(s, 3H), 4.57(d, J=8.0, 2H), 6.73(d, J=7.8, 1H), 7.24(d, J=7.8, 1H), 7.41(t, J=7.8, 1H), 7.79(s, 1H)

Starting Material Synthesis Example 112

5 5-ethoxycarbonyl-2-(4-hydroxybenzo(b)furan-2-yl)-1,3,4-oxadiazole

5-Ethoxycarbonyl-2-(4-methoxybenzo(b)furan-2-yl)-1,3,4-oxadiazole (5.0 g) was dissolved in dichloromethane (40 ml) and boron tribromide (4.0 ml) was added dropwise with stirring
10 under ice-cooling. Then, the mixture was stirred at room temperature for 5 hr, and the reaction mixture was poured into ice water. The mixture was stirred at it was at room temperature for 1 hr. The crystals were collected by filtration and dissolved in chloroform. The solution was
15 washed with saturated aqueous solution of sodium hydrogencarbonate, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (2.5 g) as yellow crystals.

¹H-NMR(DMSO-d₆)δ:1.40(t, J=8.0, 3H), 4.48(d, J=8.0, 2H), 6.74(d, J=7.8, 1H), 7.19(d, J=7.8, 1H), 7.35(t, J=7.8, 1H), 7.90(s, 1H),
20 10.46(bs, 1H)

The structural formulas of the compounds obtained in Starting Material Synthesis Examples 98 to 112 are shown in the following.

5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-2,3-dihydro-1,3,4-
oxadiazole-2-thione

131

the mixture was refluxed under heating for 7 hr. The reaction mixture was concentrated under reduced pressure and water was added to the residue. The pH was adjusted to 4 with ammonium chloride. After standing, the precipitated crystals were
5 collected by filtration and dried to give the title compound (4.6 g) as yellow crystals.

$^1\text{H-NMR}(\text{CD}_3\text{OD})\delta$: 3.44 (s, 3H), 5.27 (s, 2H), 6.90 (d, $J=7.8$, 1H), 7.15 (d, $J=8.3$, 1H), 7.25 (t, $J=7.8$, 1H), 7.35 (s, 1H)

Starting Material Synthesis Example 114

10 5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-2-methylthio-1,3,4-oxadiazole

To a suspension (40 ml) of sodium hydride (0.8 g) in THF was added dropwise a solution (30 ml) of 5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-2,3-dihydro-1,3,4-
15 oxadiazole-2-thione (4.6 g) in DMF at room temperature, and the mixture was stirred for 40 min. To this reaction mixture was added dropwise methyl iodide at room temperature, and the mixture was further stirred for 1 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The
20 organic layer was washed with water and dried. The solvent was evaporated under reduced pressure to give the title compound (2.5 g) as brown crystals.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 2.81 (s, 3H), 3.54 (s, 3H), 5.34 (s, 2H), 6.97 (d, $J=7.8$, 1H), 7.27 (d, $J=8.3$, 1H), 7.35 (t, $J=7.8$, 1H), 7.58 (s, 1H)

25 Starting Material Synthesis Example 115

5-(4-hydroxybenzo(b)furan-2-yl)-2-methylthio-1,3,4-oxadiazole

To a solution (20 ml) of 5-(4-(methoxymethyloxy)-benzo(b)furan-2-yl)-2-methylthio-1,3,4-oxadiazole (1.0 g) in THF was added 2N aqueous hydrochloric acid (3.0 ml), and the
30 mixture was refluxed under heating for 7 hr. The solvent was evaporated under reduced pressure to give the title compound (0.62 g) as yellow crystals.

$^1\text{H-NMR}(\text{DMSO}-d_6)\delta$: 2.78 (s, 3H), 6.71 (d, $J=7.9$, 1H), 7.16 (d, $J=8.3$, 1H), 7.29 (t, $J=8.3$, 1H), 7.71 (s, 1H), 10.42 (brs, 1H)

Starting Material Synthesis Example 116

5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-2,3-dihydro-1,3,4-oxadiazol-2-one

To a solution (20 ml) of 4-(methoxymethyloxy)-
5 benzo(b)furan-2-ylcarbohydrazide (1.0 g) in 1,2-dimethoxyethane
were added triphosgene (1.0 g) and triethylamine (1.8 ml), and
the mixture was stirred at room temperature for 20 min. The
reaction mixture was poured into 2N saturated aqueous sodium
hydroxide solution and soluble part of the organic layer was
10 removed with ethyl acetate. The aqueous layer was made acidic
with hydrochloric acid and extracted with ethyl acetate. This
organic layer was washed with water and dried. The solvent was
evaporated under reduced pressure to give the title compound
(1.0 g) as white crystals.

15 $^1\text{H-NMR}(\text{DMSO}-d_6)\delta$: 3.43 (s, 3H), 5.36 (s, 2H), 7.00 (d, $J=7.8$, 1H),
7.37 (d, $J=8.3$, 1H), 7.40 (t, $J=7.8$, 1H), 7.53 (s, 1H)

Starting Material Synthesis Example 117

5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-2-methoxy-1,3,4-oxadiazole

20 To a suspension of sodium hydride (0.17 g) in DMF was
added dropwise a solution (20 ml) of 5-(4-(methoxymethyloxy)-
benzo(b)furan-2-yl)-2,3-dihydro-1,3,4-oxadiazol-2-one (1.0 g)
obtained in Starting Material Synthesis Example 116 in DMF at
room temperature, and the mixture was stirred for 30 min.
25 Thereto was added methyl iodide (0.26 ml), and the mixture was
further stirred for 30 min. The reaction mixture was poured
into ice water and extracted with ethyl acetate. The extract
was washed with water and dried. The solvent was evaporated
under reduced pressure to give the title compound (0.80 g) as
30 pale-yellow crystals.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 3.54 (s, 3H), 3.55 (s, 3H), 5.33 (s, 2H), 6.97 (d,
 $J=7.8$, 1H), 7.23 (d, $J=8.3$, 1H), 7.34 (t, $J=7.8$, 1H), 7.42 (s, 1H)

Starting Material Synthesis Example 118

(S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methoxy-1,3,4-

oxadiazole

To a solution (20 ml) of 5-(4-(methoxymethyloxy)-benzo(b)furan-2-yl)-2-methoxy-1,3,4-oxadiazole (0.80 g) obtained in Starting Material Synthesis Example 117 in THF was
5 added 2N hydrochloric acid (15 ml), and the mixture was refluxed under heating for 2 hr. The reaction mixture was concentrated under reduced pressure to give crude crystals of 5-(4-hydroxybenzo(b)furan-2-yl)-2-methoxy-1,3,4-oxadiazole. This was dissolved in DMF (30 ml) and (S)-glycidyl nosylate
10 (0.83 g) and potassium carbonate (0.89 g) were added. The mixture was stirred at room temperature for 14 hr. This reaction mixture was poured into ice water, and the precipitated crystals were collected by filtration, washed with water and dried to give the title compound (0.50 g) as brown
15 crystals.

¹H-NMR(DMSO-d₆)δ:2.81(dd, J=4.9, 2.5, 1H), 2.87(t, J=4.9, 1H), 3.35-3.46(m, 1H), 3.42(s, 3H), 4.04(dd, J=11.3, 5.9, 1H), 4.53(dd, J=11.3, 2.0, 1H), 6.92(d, J=8.3, 1H), 7.31(d, J=8.3, 1H), 7.41(t, J=8.3, 1H), 7.57(s, 1H)

20 **Starting Material Synthesis Example 119**

2-ethoxy-5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 117 using sodium hydride (0.37 g),
25 5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-2,3-dihydro-1,3,4-oxadiazol-2-one (2.0 g) and ethyl iodide (0.73 ml), the title compound (1.9 g) was obtained as pale-yellow crystals.

¹H-NMR(CDCl₃)δ:1.44(t, J=7.3, 3H), 3.54(s, 3H), 3.90(q, J=7.3, 2H), 5.33(s, 2H), 6.97(d, J=7.8, 1H), 7.23(d, J=8.3, 1H),
30 7.32(t, J=7.8, 1H), 7.42(s, 1H)

Starting Material Synthesis Example 120

(S)-2-ethoxy-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole

To a solution (40 ml) of 2-ethoxy-5-(4-

(methoxymethyloxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole (2.0 g) obtained in the same manner as in Starting Material Synthesis Example 119 in THF was added 2N hydrochloric acid (40 ml), and the mixture was refluxed under heating for 2 hr. The reaction mixture was concentrated under reduced pressure to give crude crystals of 2-ethoxy-5-(4-hydroxybenzo(b)furan-2-yl)-1,3,4-oxadiazole. This was dissolved in DMF (40 ml) and (S)-glycidyl nosylate (1.7 g) and potassium carbonate (3.5 g) were added. The mixture was stirred at room temperature for 14 hr. This reaction mixture was poured into ice water, and after extraction with ethyl acetate, the extract was washed with water and dried. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/ethyl acetate) to give the title compound (1.0 g) as white crystals.

¹H-NMR(CDCl₃)δ:1.44(t, J=7.3, 3H), 2.81(dd, J=4.9, 2.9, 1H), 2.97(t, J=4.3, 1H), 3.42-3.46(m, 1H), 3.90(q, J=7.3, 2H), 4.10(dd, J=11.2, 5.9, 1H), 4.40(dd, J=11.2, 2.9, 1H), 6.72(d, J=8.3, 1H), 7.21(d, J=8.3, 1H), 7.34(t, J=8.3, 1H), 7.44(s, 1H)

Starting Material Synthesis Example 121

2-(1-methylethyloxy)-5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole

By the reactions in the same manner as in Starting Material Synthesis Example 117 using sodium hydride (0.28 g), 5-(4-(methoxymethyloxy)benzo(b)furan-2-yl)-2,3-dihydro-1,3,4-oxadiazol-2-one (1.5 g) and 2-iodopropane (1.1 g), the title compound (1.5 g) was obtained as brown crystals.

¹H-NMR(CDCl₃)δ:1.47(d, J=6.3, 6H), 3.54(s, 3H), 4.44(penth, J=6.3, 1H), 5.33(s, 2H), 6.96(d, J=7.8, 1H), 7.25(d, J=8.3, 1H), 7.33(t, J=7.8, 1H), 7.42(s, 1H)

Starting Material Synthesis Example 122

(S)-2-(1-methylethyloxy)-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole

To a solution (30 ml) of 2-(1-methylethyloxy)-5-(4-

(methoxymethyloxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole (1.5 g) obtained in Starting Material Synthesis Example 121 in THF was added 4N hydrochloric acid (15 ml), and the mixture was refluxed under heating for 3 hr. The reaction mixture was concentrated under reduced pressure to give crude crystals of 2-(1-methylethyloxy)-5-(4-hydroxybenzo(b)furan-2-yl)-1,3,4-oxadiazole. This was dissolved in DMF (30 ml) and (S)-glycidyl nosylate (1.2 g) and potassium carbonate (3.2 g) were added. The mixture was stirred at room temperature for 5 hr. This reaction mixture was poured into ice water, and after extraction with ethyl acetate, the extract was washed with water and dried. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/ethyl acetate) to give the title compound (1.3 g) as white crystals.

¹H-NMR(CDCl₃)δ:1.35(d, J=6.3, 6H), 2.81(dd, J=4.9, 2.9, 1H), 2.88(t, J=4.3, 1H), 3.40-3.42(m, 1H), 4.04(dd, J=11.7, 6.4, 1H), 4.30(penth, J=6.3, 1H), 4.53(dd, J=11.7, 1.9, 1H), 6.91(d, J=8.3, 1H), 7.33(d, J=7.8, 1H), 7.41(t, J=8.3, 1H), 7.55(s, 1H)


Starting Material Synthesis Example 123

4-(2'-methoxybenzylidene)-2-methyl-4H-oxazol-5-one


Anisaldehyde (11.3 g), N-acetylglycine (9.8 g) and sodium acetate (8.2 g) were dissolved in acetic anhydride (200 ml) and the mixture was heated at 100°C for 10 hr. After cooling, the precipitated yellow crystals were collected by filtration to give the title compound (9.52 g), melting point 151-153°C.

The structural formulas of the compounds obtained in Starting Material Synthesis Examples 113 to 123 are shown in the following.


113




114




115




116




117




118




119




120




121



122



123

**Example 1**

(S)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-

propyloxy)benzo(b)furan-2-ylcarbonyl)pyrrolidine

(S)-1-(4-Glycidyloxybenzo(b)furan-2-ylcarbonyl)-
pyrrolidine (1.2 g) obtained in Starting Material Synthesis
Example 1 was dissolved in methanol (40 ml) and 4-(naphthalen-
5 2-yl)piperidine (0.85 g) was added. The mixture was refluxed
under heating for 8 hr. The reaction mixture was evaporated
under reduced pressure and the obtained residue was purified by
silica gel column chromatography (chloroform/methanol) to give
the title compound (1.6 g) as a brown oil.

10 $^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.81-2.20 (m, 8H), 2.22 (t, $J=11.7$, 1H), 2.56-
2.96 (m, 1H), 2.62-2.79 (m, 3H), 3.03 (d, $J=10.8$, 1H), 3.22 (d,
 $J=10.8$, 1H), 4.10-4.28 (m, 3H), 6.73 (d, $J=8.3$, 1H), 7.16 (d,
 $J=8.3$, 1H), 7.33 (t, $J=8.3$, 1H), 7.35-7.50 (m, 3H), 7.51-7.55 (m,
1H), 7.67 (s, 1H), 7.81 (d, $J=8.8$, 3H)

15 **Example 2**

(S)-4-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-
propyloxy)benzo(b)furan-2-ylcarbonyl)morpholine

(S)-4-(4-Glycidyloxybenzo(b)furan-2-ylcarbonyl)-
morpholine (1.3 g) obtained in Starting Material Synthesis
20 Example 2 was dissolved in methanol (40 ml) and 4-(naphthalen-
2-yl)piperidine (0.91 g) was added. The mixture was refluxed
under heating for 8 hr and the reaction solvent was evaporated
under reduced pressure. The obtained residue was purified by
silica gel column chromatography (chloroform/methanol) to give
25 the title compound (1.8 g) as a brown oil.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.86-1.99 (m, 4H), 2.21 (t, $J=11.7$, 1H), 2.53 (t,
 $J=11.2$, 1H), 2.59-2.74 (m, 3H), 3.03 (d, $J=10.8$, 1H), 3.22 (d,
 $J=10.8$, 1H), 3.70-4.03 (m, 8H), 4.10-4.27 (m, 3H), 6.73 (d, $J=8.3$,
1H), 7.15 (d, $J=8.3$, 1H), 7.33 (t, $J=8.3$, 1H), 7.37-7.41 (m, 3H),
30 7.49 (s, 1H), 7.67 (s, 1H), 7.81 (d, $J=8.8$, 3H)

Example 3

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-
methylbenzo(b)furan-2-carboxamide

To a solution (13 ml) of (S)-4-(2-hydroxy-3-(4-

(naphthalen-2-yl)piperidino)propyloxy)benzo(b)furan-2-carboxylic acid (0.12 g) obtained in Starting Material Synthesis Example 22 in DMF were added methylamine hydrochloride (0.18 g), triethylamine (0.1 ml) and diethyl
5 cyanophosphate (0.1 ml), and the mixture was stirred for 1 hr at room temperature. The reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution and water and dried over anhydrous magnesium sulfate. The solvent
10 was evaporated under reduced pressure and the obtained residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.05 g) as a brown oil.

¹H-NMR(CDCl₃)δ: 1.84-1.97(m, 4H), 2.20(t, J=11.7, 1H), 2.45-
15 2.55(m, 1H), 2.59-2.79(m, 3H), 2.99-3.06(m, 1H), 3.03(d, J=5.3, 3H), 3.20(d, J=9.7, 1H), 4.11-4.20(m, 3H), 6.60(br, 1H), 6.70(d, J=8.3, 1H), 7.08(d, J=8.3, 1H), 7.31(t, J=8.3, 1H), 7.35-7.41(m, 3H), 7.59(s, 1H), 7.65(s, 1H), 7.78(d, J=8.8, 3H)

Example 4

20 (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)furan-2-carboxamide

By the reactions as in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylic acid (0.8 g) obtained in
25 Starting Material Synthesis Example 22, dimethylamine hydrochloride (0.15 g), triethylamine (0.49 ml) and diethyl cyanophosphate (0.33 ml), the title compound (0.61 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.84-2.00(m, 4H), 2.22(t, J=11.0, 1H), 2.49-
30 2.55(m, 1H), 2.65-2.77(m, 3H), 3.03(brd, J=10.7, 1H), 3.16(brs, 3H), 3.22(brd, J=10.7, 1H), 3.36(brs, 3H), 4.14-4.24(m, 3H), 6.72(d, J=8.3, 1H), 7.16(d, J=8.3, 1H), 7.31(t, J=8.3, 1H), 7.39-7.48(m, 3H), 7.67(s, 1H), 7.80(d, J=8.8, 3H)

Example 5

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-diethylbenzo(b)furan-2-carboxamide

By the reactions in the same manner as in as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylic acid (0.8 g) obtained in Starting Material Synthesis Example 22, diethylamine (0.24 ml) and diethyl cyanophosphate (0.5 ml), the title compound (0.61 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.19-1.40 (m, 6H), 1.82-2.00 (m, 4H), 2.22 (t, J=12.2, 1H), 2.49-2.55 (m, 1H), 2.64-2.76 (m, 3H), 3.04 (brd, J=11.3, 1H), 3.21 (brd, J=11.3, 1H), 3.43-3.70 (m, 4H), 4.12-4.24 (m, 3H), 6.72 (d, J=8.3, 1H), 7.14 (d, J=8.3, 1H), 7.30 (t, J=8.3, 1H), 7.38-7.48 (m, 3H), 7.67 (s, 1H), 7.80 (d, J=8.3, 3H)

Example 6

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-methoxy-N-methylbenzo(b)furan-2-carboxamide

By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylic acid (0.8 g) obtained in Starting Material Synthesis Example 22, N,O-dimethylhydroxylamine hydrochloride (0.24 g), triethylamine (1.0 ml) and diethyl cyanophosphate (0.27 ml), the title compound (0.64 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.86-1.99 (m, 4H), 2.22 (t, J=10.2, 1H), 2.49-2.53 (m, 1H), 2.63-2.74 (m, 3H), 3.04 (brd, 11.7, 1H), 3.22 (brd, 11.7, 1H), 3.42 (s, 3H), 3.92 (s, 3H), 4.14-4.27 (m, 3H), 6.72 (d, J=7.8, 1H), 7.23 (d, J=7.8, 1H), 7.34 (t, J=7.8, 1H), 7.38-7.48 (m, 3H), 7.63 (s, 1H), 7.67 (s, 1H), 7.79-7.82 (m, 3H)

Example 7

(S)-4-(8-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-2H-chromen-3-ylcarbonyl)morpholine

By the reactions in the same manner as in Example 1 using (S)-4-(8-glycidyloxy-2H-chromen-3-ylcarbonyl)morpholine

(3.1 g) obtained in Starting Material Synthesis Example 6 and 4-(naphthalen-2-yl)piperidine (2.5 g), the title compound (3.5 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.86-1.99(m, 4H), 2.21(t, J=11.7, 1H), 2.49-
5 2.56(m, 1H), 2.63-2.74(m, 3H), 3.03(d, J=11.7, 1H), 3.22(d, J=11.7, 1H), 3.42(s, 3H), 3.84(s, 3H), 4.14-4.27(m, 3H), 6.72(d, J=8.3, 1H), 7.23(d, J=8.3, 1H), 7.43(d, J=8.3, 1H), 7.44-7.48(m, 2H), 7.63(s, 1H), 7.68(s, 1H), 7.78-7.82(m, 3H)

Example 8

10 (S)-4-(8-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-2H-chromen-3-ylmethyl)morpholine maleate

To a suspension of lithium aluminum hydride (0.55 g) in THF was added aluminum chloride (0.63 g) and the mixture was stirred at room temperature for 1 hr. The reaction mixture was
15 made to become 4°C and a solution of (S)-4-(8-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-2H-chromen-3-ylcarbonyl)morpholine (2.5 g) in THF (50 ml) was added dropwise. The mixture was stirred for 30 min and hydrous THF was added. The mixture was further stirred for 30 min at room temperature
20 and the precipitated insoluble matter was filtered off through celite. The solvent was evaporated under reduced pressure to give a brown oil. This was dissolved in ethanol and maleic acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.3 g) as
25 pale-yellow crystals, melting point 164-166°C.

Example 9

(S)-8-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethyl-2H-chromene-3-carboxamide

By the reactions in the same manner as in Example 1
30 using (S)-8-glycidyloxy-N,N-dimethyl-2H-chromene-3-carboxamide (3.2 g) obtained in Starting Material Synthesis Example 8 and 4-(naphthalen-2-yl)piperidine (1.5 g), the title compound (3.7 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.86-1.96 (m, 4H), 2.19 (t, J=11.7, 1H), 2.43-
2.55 (m, 1H), 2.59-2.89 (m, 3H), 2.96 (s, 3H), 2.97 (s, 3H), 2.90-
3.32 (m, 2H), 4.07-4.32 (m, 3H), 6.61 (s, 1H), 6.73 (d, J=8.3, 1H),
6.86 (t, J=8.3, 1H), 6.93 (d, J=8.3, 1H), 7.35-7.47 (m, 3H),
5 7.66 (s, 1H), 7.78-7.80 (m, 3H)

Example 10











(S)-3-chloro-6-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-
propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Example 1, the
10 title compound (0.4 g) was obtained from (S)-3-chloro-6-
glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide (0.6 g)
obtained in Starting Material Synthesis Example 14 and 4-
(naphthalen-2-yl)piperidine (0.45 g).

¹H-NMR(CDCl₃)δ: 1.87-1.96 (m, 3H), 2.05-2.22 (m, 1H), 2.52-2.70 (m,
15 4H), 3.03-3.22 (m, 10H), 4.08-4.20 (m, 3H), 7.13-7.16 (m, 1H),
7.30 (d, 1H, J=1.9), 7.39 (d, 1H, J=8.8), 7.43-7.48 (m, 2H),
7.60 (s, 1H), 7.72 (d, 1H, J=8.3), 7.77-7.82 (m, 3H, J=8.3)

The structural formulas of the compounds obtained in
Examples 1 to 10 are shown in the following.

20



Example 11

(S)-3-chloro-6-(2-hydroxy-3-(4-(naphthalen-1-yl)piperidino)-propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Example 1
5 using (S)-3-chloro-6-glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide (0.6 g) obtained in Starting Material Synthesis Example 14 and 4-(naphthalene-1-yl)piperidine (0.45 g), the title compound (0.5 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.81-2.33(m, 3H), 2.30-2.37(m, 1H), 2.62-2.70(m,
10 4H), 3.11-3.17(m, 8H), 3.21-3.25(m, 1H), 3.35-3.44(m, 1H), 4.02-4.15(m, 2H), 4.18-4.22(m, 1H), 7.15(d, 1H, J=6.8), 7.30(s, 1H), 7.40-7.49(m, 4H), 7.75-7.79(m, 2H), 7.88(d, 1H, J=7.8), 8.10(d, 1H, J=8.3)

Example 12

15 (S)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)thiophen-2-ylcarbonyl)pyrrolidine 2 methanesulfonate monohydrate

By the reactions in the same manner as in Example 1
using (S)-1-(4-glycidyloxybenzo(b)thiophen-2-ylcarbonyl)-
20 pyrrolidine (4.0 g) obtained in similar manner as in Starting Material Synthesis Example 19 and 4-(naphthalen-2-yl)piperidine (2.2 g), a brown oil (4.2 g) was obtained. This was dissolved in ethyl acetate and methanesulfonic acid was added. The precipitated crystals were collected by filtration and dried to
25 give the title compound (3.3 g) as pale-yellow crystals, melting point 88-90°C.

Example 13

(S)-4-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)thiophen-2-ylcarbonyl)morpholine

30 By the reactions in the same manner as in Example 1, the title compound (0.7 g) was obtained from (S)-4-(4-glycidyloxybenzo(b)thiophen-2-ylcarbonyl)morpholine (1.2 g) obtained in Starting Material Synthesis Example 18 and 4-

(naphthalen-2-yl)piperidine (1.0 g), melting point 82-86°C.

Example 14

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-methoxy-N-methylbenzo(b)thiophene-2-carboxamide

5 By the reactions in the same manner as in Example 1, the title compound (0.8 g) was obtained as a brown oil from (S)-4-glycidyloxy-N-methoxy-N-methylbenzo(b)thiophene-2-carboxamide (1.1 g) obtained in Starting Material Synthesis Example 20 and 4-(naphthalen-2-yl)piperidine (0.8 g).

10 ¹H-NMR(CDCl₃)δ: 1.86-1.99(m, 4H), 2.23(t, 1H, J=9.8), 2.47-2.55(m, 1H), 2.63-2.74(m, 3H), 3.05(d, 1H, J=11.2), 3.23(d, 1H, J=11.2), 3.43(s, 3H), 3.83(s, 3H), 4.11-4.15(m, 1H), 4.20-4.27(m, 2H), 6.79(d, 1H, J=7.8), 7.35-7.48(m, 5H), 7.68(s, 1H), 7.81(d, 3H, J=8.3), 8.42(s, 1H)

15 **Example 15**

(S)-4-(2-hydroxy-3-(4-(naphthalen-1-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Example 1, the title compound (0.4 g) was obtained from (S)-4-glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide (0.5 g) obtained in
20 Starting Material Synthesis Example 17 and 4-(naphthalen-1-yl)piperidine (0.4 g), as pale-yellow crystals, melting point 97-100°C.

Example 16

25 (S)-4-(2-hydroxy-3-(4-(6-methoxynaphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Example 1, the title compound (1.2 g) was obtained from (S)-4-glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide (1.7 g) obtained in
30 Starting Material Synthesis Example 17 and 4-(6-methoxynaphthalen-2-yl)piperidine (1.5 g).

¹H-NMR(CDCl₃)δ: 1.82-2.00(m, 3H), 2.07-2.23(m, 1H), 2.52-2.57(m, 1H), 2.63-2.75(m, 3H), 3.02-3.05(m, 1H), 1.10-3.20(bs, 6H), 3.91(s, 3H), 4.09-4.23(m, 3H), 6.79(d, 1H, J=7.9), 7.11(s, 1H),

7.14(d, 1H, J=2.5), 7.30-7.37(m, 2H), 7.44(d, 1H, J=7.8),
7.59(s, 1H), 7.69(s, 1H), 7.70(s, 1H), 7.74(s, 1H)

Example 17

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-
5 N,N-dimethylbenzo(b)thiophene-2-carboxamide (L)-tartrate

By the reactions in the same manner as in Example 1, a brown oil (1.9 g) was obtained from (S)-4-glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide (1.2 g) obtained in Starting Material Synthesis Example 17 and 4-(naphthalen-2-
10 yl)piperidine (0.9 g). This was dissolved in ethanol and (L)-tartaric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.2 g) as white crystals, melting point 173-175°C.

Example 18

15 (S)-4-(2-hydroxy-3-(4-(naphthalen-1-yl)-3,6-dihydro-2H-pyridin-1-yl)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Example 1 using (S)-4-glycidyloxy-N,N-dimethylbenzo(b)thiophene-2-carboxamide (2.0 g) obtained in Starting Material Synthesis
20 Example 17 and 4-(naphthalen-1-yl)-3,6-dihydro-2H-pyridine (2.0 g), the title compound (0.8 g) was obtained.

¹H-NMR(CDCl₃)δ: 2.63-2.81(m, 6H), 3.05-3.40(m, 6H), 2.98-3.02(m, 1H), 3.41-3.44(m, 1H), 4.17-4.23(m, 2H), 4.25-4.33(m, 1H),
6.25(s, 1H), 6.79(d, 1H, J=7.9), 7.32(t, 1H, J=7.9), 7.40-
25 7.58(m, 2H), 7.60(d, 1H, J=10.2), 7.74-7.83(m, 6H)

Example 19

(S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-
methylbenzo(b)furan-2-carboxamide

By the reactions in the same manner as in Example 3
30 using (S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)benzo(b)furan-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 24, methylamine (0.15 g), triethylamine (0.63 ml) and diethyl cyanophosphate (0.37 ml), the title compound (0.75 g) was obtained as a brown

oil.

¹H-NMR(CDCl₃)δ: 1.85-1.97(m, 4H), 2.20(t, J=11.7, 1H), 2.45-2.55(m, 1H), 2.59-2.79(m, 3H), 2.99-3.06(m, 1H), 3.04(d, J=5.3, 3H), 3.20(d, J=9.7, 1H), 4.07-4.27(m, 3H), 4.18-4.38(s, m),
5 6.82(br, 1H), 6.94(d, J=8.3, 1H), 7.18(t, J=8.3, 1H), 7.31(t, J=8.3, 1H), 7.37-7.46(m, 3H), 7.66(s, 1H), 7.79(d, J=8.8, 3H)

Example 20


(S)-4-(7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-ylcarbonyl)morpholine

10 By the reactions in the same manner as in Example 3 using (S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 24, morpholine (0.19 g), triethylamine (0.63 ml) and diethyl cyanophosphate (0.37 ml),
15 the title compound (0.60 g) was obtained as a brown oil.


¹H-NMR(CDCl₃)δ: 1.86-1.99(m, 4H), 2.22(t, J=11.7, 1H), 2.53-2.58(m, 1H), 2.59-2.80(m, 3H), 3.03(d, J=10.8, 1H), 3.23(d, J=10.8, 1H), 3.72-4.03(m, 8H), 4.20-4.36(m, 3H), 6.96(d, J=8.3, 1H), 7.22(t, J=8.3, 1H), 7.25(d, J=8.3, 1H), 7.37-7.41(m, 3H),
20 7.49(s, 1H), 7.66(s, 1H), 7.81(d, J=8.8, 3H)

The structural formulas of the compounds obtained in Examples 11 to 20 are shown in the following.


11




12




13




14




15




16




17




18



19



20



Example 21

(S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)furan-2-carboxamide

By the reactions in the same manner as in Example 3
5 using (S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 24, dimethylamine hydrochloride (0.18 g), triethylamine (0.63 ml) and diethyl cyanophosphate (0.37 ml), the title compound (0.60 g) was
10 obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.81-2.01 (m, 4H), 2.18-2.29 (m, 1H), 2.44-2.58 (m, 1H), 2.61-2.78 (m, 3H), 2.88 (s, 3H), 2.95 (s, 3H), 3.03 (d, J=10.8, 1H), 3.24 (d, J=10.8, 1H), 4.20-4.37 (m, 3H), 6.95 (d, J=7.8, 1H), 7.19 (t, J=7.8, 1H), 7.25 (d, J=7.8, 1H), 7.31 (s, 1H), 7.38-
15 7.48 (m, 3H), 7.66 (s, 1H), 7.80 (d, J=8.8, 3H)

Example 22

(S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-methoxy-N-methylbenzo(b)furan-2-carboxamide

By the reactions in the same manner as in Example 3
20 using (S)-7-(2-hydroxy-3-(4-(naphthalen-2-yl)-piperidino)propyloxy)benzo(b)furan-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 24, N,O-dimethylhydroxylamine (0.21 g), triethylamine (0.63 ml) and diethyl cyanophosphate (0.37 ml), the title compound (0.62 g)
25 was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.83-2.01 (m, 4H), 2.21-2.29 (m, 1H), 2.43-2.58 (m, 1H), 2.63-2.78 (m, 3H), 3.03 (brd, J=10.8, 1H), 3.23 (d, J=10.8, 1H), 3.42 (s, 3H), 3.86 (s, 3H), 4.21-4.38 (m, 3H), 6.98 (d, J=7.8, 1H), 7.20 (t, J=7.8, 1H), 7.38-7.48 (m, 3H), 7.66 (s, 1H), 7.80 (d,
30 J=8.8, 3H)

Example 23

(S)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indol-2-ylcarbonyl)-4-methylpiperazine

By the reactions in the same manner as in Example 3

using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indole-2-carboxylic acid (0.70 g) obtained in Starting Material Synthesis Example 25, N-methylpiperazine (0.16 g), triethylamine (0.44 ml) and diethyl cyanophosphate (0.27 ml), the title compound (0.65 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.73-2.04(m, 4H), 2.16-2.20(m, 1H), 2.34(s, 3H), 2.49-2.79(m, 7H), 3.03(d, J=10.7, 1H), 3.15-3.36(m, 5H), 4.10-4.37(m, 3H), 6.54(d, J=8.3, 1H), 6.93(s, 1H), 7.00(d, J=8.3, 1H), 7.18(t, J=8.3, 1H), 7.38-7.46(m, 3H), 7.67(s, 1H), 7.78(m, 3H), 9.29(s, 1H)

Example 24

(S)-4-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indol-2-ylcarbonyl)morpholine hydrochloride

By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indole-2-carboxylic acid (0.70 g) obtained in Starting Material Synthesis Example 25, morpholine (0.14 g), triethylamine (0.44 ml) and diethyl cyanophosphate (0.27 ml), a brown oil (0.66 g) was obtained. This was dissolved in acetone and 1N solution of hydrochloric acid in methanol was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.65 g) as white crystals, melting point 169-171°C.

Example 25

(S)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indol-2-ylcarbonyl)pyrrolidine 3/2 hydrochloride

By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indole-2-carboxylic acid (0.70 g) obtained in Starting Material Synthesis Example 25, pyrrolidine (0.11 g), triethylamine (0.44 ml) and diethyl cyanophosphate (0.27 ml), the title compound (0.24 g) was obtained as white crystals,

melting point 158-161°C.

Example 26

(R)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indol-2-ylcarbonyl)pyrrolidine

5 By the reactions in the same manner as in Example 3 using (R)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1H-indole-2-carboxylic acid (1.0 g) obtained by the reactions in the same manner as in Starting Material Synthesis Example 25 from (R)-glycidyl nosylate, pyrrolidine (0.30 g),
10 triethylamine (3.0 ml) and diethyl cyanophosphate (0.30 ml), the title compound (0.54 g) was obtained as white crystals, melting point 211-212°C.

Example 27

(R)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-

15 N,N-dimethyl-1H-indole-2-carboxamide

By the reactions in the same manner as in Example 3 using (R)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1H-indole-2-carboxylic acid (1.0 g) obtained by the reactions in the same manner as in Starting
20 Material Synthesis Example 25 from (R)-glycidyl nosylate, dimethylamine hydrochloride (0.3 g), triethylamine (3.0 ml) and diethyl cyanophosphate (0.3 ml), the title compound (0.24 g) was obtained as white crystals, melting point 158-160°C.

Example 28

25 4-(2-hydroxy-3-(2-(2-naphthoxy)ethylamino)propyloxy)-1H-indole-2-carboxamide

By the reactions in the same manner as in Example 1 using 4-glycidyloxy-1H-indole-2-carboxamide (0.70 g) and 2-(2-naphthoxy)ethylamine (0.70 g), the title compound (0.57 g) was
30 obtained as white crystals, melting point 125-126°C.

Example 29

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1-methyl-N-methylindole-2-carboxamide

By the reactions in the same manner as in Example 3

using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-methylindole-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 26, methylamine hydrochloride (0.2 g), triethylamine (1.0 ml) and diethyl
5 cyanophosphate (0.5 ml), a yellow oil (0.8 g) was obtained. To this oil was added isopropyl ether and the precipitated crystals were collected by filtration to give the title compound (0.5 g) as pale-yellow crystals, melting point 180-183°C.

10 ¹H-NMR(CDCl₃)δ: 1.87-1.96 (m, 4H), 2.19-2.50 (m, 1H), 2.50-2.80 (m, 4H), 2.90-3.20 (m, 4H), 3.21 (m, 1H), 4.04 (s, 3H), 4.14-4.18 (m, 3H), 6.19 (brs, 1H), 6.55 (d, J=7.8, 1H), 6.98-7.08 (m, 2H), 7.20-7.22 (m, 1H), 7.38-7.46 (m, 3H), 7.66 (m, 1H), 7.79-7.81 (m, 3H)











Example 30

15 (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1-methyl-N,N-dimethylindole-2-carboxamide

By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-methylindole-2-carboxylic acid (0.6 g) obtained in
20 Starting Material Synthesis Example 26, dimethylamine hydrochloride (0.3 g), triethylamine (1.0 ml) and diethyl cyanophosphate (0.5 ml), the title compound (0.6 g) was obtained as pale-yellow crystals, melting point 146-148°C.

25 ¹H-NMR(CDCl₃)δ: 1.84-1.93 (m, 4H), 2.16-2.20 (m, 1H), 2.50-2.80 (m, 4H), 3.00-3.40 (m, 8H), 3.81 (s, 3H), 4.10-4.30 (m, 3H), 6.54 (d, J=8.4, 1H), 6.77 (s, 1H), 6.96 (d, J=8.3, 1H), 7.18 (dd, J=7.8, 7.8, 1H), 7.24 (s, 1H), 7.36-7.45 (m, 3H), 7.64 (s, 1H), 7.78 (d, J=7.8, 2H)

The structural formulas of the compounds obtained in
30 Examples 21 to 30 are shown in the following.



Example 31

(S)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-methylindole-2-carbonyl)pyrrolidine

By the reactions in the same manner as in Example 3
5 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-methylindole-2-carboxylic acid (1.8 g) obtained in Starting Material Synthesis Example 26, pyrrolidine (0.5 ml), triethylamine (0.5 ml) and diethyl cyanophosphate (1.5 ml), the title compound (0.2 g) was obtained as a yellow oil.

10 ¹H-NMR(CDCl₃)δ: 1.84-2.04(m, 9H), 2.23(m, 1H), 2.50(m, 1H), 2.66-2.80(m, 2H), 3.00-3.30(m, 2H), 3.60-3.80(m, 4H), 3.92(s, 3H), 4.00-4.30(m, 3H), 6.55(d, J=7.8, 1H), 6.89(s, 1H), 6.99(d, J=8.3, 1H), 7.22(dd, J=7.8, 8.3, 1H), 7.38(s, 1H), 7.40-7.47(m, 3H), 7.66(s, 1H), 7.80(d, J=7.3, 2H)

15 **Example 32**

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-methyl-1-(2-methylpropyl)indole-2-carboxamide hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 3
20 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-(2-methylpropyl)indole-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 27, methylamine hydrochloride (0.2 g), triethylamine (0.7 ml) and diethyl cyanophosphate (0.5 ml), a yellow oil (0.8 g) was obtained. A
25 1N solution of hydrochloric acid in isopropyl was added to this oil in isopropyl ether. The precipitated crystals were collected by filtration and dried to give the title compound (0.7 g) as pale-yellow crystals, melting point 108-110°C.

¹H-NMR(CD₃OD)δ: 1.10-1.12(m, 7H), 2.09-2.24(m, 5H), 2.91(s, 3H),
30 3.11-3.60(m, 4H), 3.84-3.92(m, 2H), 4.15-4.25(m, 2H), 4.37(d, J=7.4, 2H), 4.57(m, 1H), 6.60(d, J=7.8, 1H), 7.09(d, J=8.3, 1H), 7.16-7.22(m, 2H), 7.43-7.46(m, 3H), 7.74-7.867(m, 4H)

Example 33

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-

N,N-dimethyl-1-(2-methylpropyl)indole-2-carboxamide
hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-(2-methylpropyl)indole-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 27, dimethylamine hydrochloride (0.2 g), triethylamine (0.7 ml) and diethyl cyanophosphate (0.5 ml), the title compound (0.6 g) was obtained as pale-yellow crystals, melting point 108-110°C.

¹H-NMR(CD₃OD)δ: 1.10-1.12 (m, 7H), 2.03 (m, 1H), 2.10-2.30 (m, 4H), 3.00-3.40 (m, 8H), 3.40-3.60 (m, 2H), 3.80-3.95 (m, 2H), 4.12 (d, J=7.8, 2H), 4.20-4.25 (m, 2H), 4.57 (m, 1H), 6.62 (d, J=7.8, 1H), 6.87 (s, 1H), 7.10 (d, J=8.3, 1H), 7.17 (dd, J=7.8, 8.3m, 1H), 7.43-7.49 (m, 3H), 7.74-7.86 (m, 4H)

Example 34

(S)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-(2-methylpropyl)indole-2-carbonyl)pyrrolidine
hydrochloride

By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-(2-methylpropyl)indole-2-carboxylic acid (1.0 g) obtained in Starting Material Synthesis Example 27, pyrrolidine (0.2 ml), triethylamine (0.7 ml) and diethyl cyanophosphate (0.5 ml), the title compound (0.4 g) was obtained as pale-yellow crystals, melting point 104-106°C.

¹H-NMR(CD₃OD)δ: 0.79-0.81 (m, 7H), 1.91-2.14 (m, 9H), 3.00-3.40 (m, 4H), 3.60-3.80 (m, 6H), 4.15-4.25 (m, 4H), 4.57 (m, 1H), 6.61 (d, J=7.8, 1H), 6.98 (s, 1H), 7.08 (d, J=8.3, 1H), 7.20 (dd, J=7.8, 8.3m, 1H), 7.42-7.69 (m, 3H), 7.72-7.84 (m, 4H)

Example 35

(S)-1-(2-(5-methyl-1,2,4-oxadiazol-3-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Example 3 using (S)-3-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,2,4-

oxadiazole (0.45 g) obtained in Starting Material Synthesis Example 31 and 4-(naphthalen-2-yl)piperidine (0.35 g), the title compound (0.65 g) was obtained as white crystals, melting point 146-148°C.

5 **Example 36**

(S)-1-(2-(5-methyl-1,2,4-oxadiazol-3-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-3-(7-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole (1.7 g) obtained in Starting Material Synthesis Example 35 and 4-(naphthalen-2-yl)piperidine (1.3 g), the title compound (2.0 g) was obtained as white crystals, melting point 169-170°C.

Example 37

15 (S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.33 g) obtained in Starting Material Synthesis Example 39 and 4-(naphthalen-2-yl)piperidine (0.26 g), a brown oil (0.5 g) was obtained. This was dissolved in ethyl acetate and 1N solution of hydrochloric acid in ether was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.30 g) as pale-yellow crystals, 25 melting point 158-160°C.

Example 38

(S)-1-(2-(5-trifluoromethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-trifluoromethyl-1,3,4-oxadiazole (1.0 g) obtained in Starting Material Synthesis Example 51 and 4-(naphthalen-2-yl)piperidine (0.75 g), the title compound (0.5 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.87-2.00 (m, 4H), 2.23 (t, J=11.7, 1H), 2.51-2.58 (m, 1H), 2.63-2.76 (m, 3H), 3.05 (brd, J=10.3, 1H), 3.23 (brd, J=10.3, 1H), 4.15-4.26 (m, 3H), 6.79 (d, J=8.3, 1H), 7.26 (d, J=8.3, 1H), 7.39-7.48 (m, 4H), 7.64 (s, 1H), 7.80 (d, J=8.3, 4H)

5 **Example 39**

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

To a solution of 2-(7-methoxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (8.0 g) obtained in Starting Material
10 Synthesis Example 40 in methylene chloride (100 ml) was added dropwise boron tribromide (10 ml) at -8°C. The mixture was stirred under ice-cooling for 1 hr. The reaction mixture was poured into ice water and the mixture was extracted with chloroform. The organic layer was dried over anhydrous sodium
15 sulfate and concentrated under reduced pressure to give red crystals (6.0 g) of 7-hydroxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan. This compound and (S)-glycidyl nosylate (7.25 g) were dissolved in dimethylformamide (100 ml) and potassium carbonate (11 g) was added. The mixture was heated
20 at 50°C for 2 hr. The reaction mixture was poured into ice water and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product
25 (6.0 g). The oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (50 ml) and the mixture was refluxed under heating for 1 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to
30 give the title compound (3.0 g) as pale-yellow crystals, melting point 140-142°C.

¹H-NMR(DMSO-d₆)δ: 1.77-1.83 (m, 4H), 2.20-2.25 (m, 2H), 2.47-2.66 (m, 3H), 2.62 (s, 3H), 3.04-3.13 (m, 2H), 4.17 (m, 2H), 4.30 (m, 1H), 5.02 (bs, 1H), 7.14 (d, J=7.8, 1H), 7.29 (t, J=7.8, 1H),

7.34(d, J=7.8, 1H), 7.41-7.48(m, 3H), 7.70(s, 1H), 7.72(s, 1H),
7.81-7.84(m, 3H)











Example 40

(S)-1-(2-(5-trifluoromethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-
5 7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1
using (S)-2-(7-glycidyloxybenzo(b)furan-2-yl)-5-
trifluoromethyl-1,3,4-oxadiazole (1.0 g) obtained in Starting
Material Synthesis Example 55 and 4-(naphthalen-2-yl)piperidine
10 (0.80 g), the title compound (0.35 g) was obtained as a brown
oil.

¹H-NMR(CDCl₃)δ: 1.81-2.00(m, 4H), 2.21-2.25(m, 1H), 2.47-2.60(m,
1H), 2.60-2.79(m, 3H), 3.07(d, J=9.8, 1H), 3.21-3.30(m, 1H),
4.23-4.31(m, 3H), 7.02-7.09(m, 1H), 7.21-7.36(m, 3H), 7.40-
15 7.54(m, 3H), 7.68(s, 1H), 7.81(d, J=7.8, 1H)

The structural formulas of the compounds obtained in
Examples 31 to 40 are shown in the following.



Example 41

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)thiophen-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

2-(4-Hydroxybenzo(b)thiophen-2-yl)-5-methyl-1,3,4-oxadiazole (1.4 g) obtained in Starting Material Synthesis Example 43 and (S)-glycidyl nosylate (1.3 g) were dissolved in dimethylformamide (15 ml) and potassium carbonate (1.5 g) was added. The mixture was heated at 50°C for 2 hr. The reaction mixture was poured into ice water and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oil (1.7 g). The oil and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (20 ml) and the mixture was refluxed under heating for 1 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.36 g) as a brown oil.

¹H-NMR(DMSO-d₆)δ: 1.77-1.85(m, 4H), 2.18-2.25(m, 2H), 2.49-2.68(m, 3H), 2.61(s, 3H), 3.05-3.15(m, 2H), 4.18(m, 2H), 4.36(m, 1H), 5.02(bs, 1H), 7.01(d, J=7.8, 1H), 7.32(t, J=7.8, 1H), 7.34(d, J=7.8, 1H), 7.41-7.48(m, 3H), 7.74(s, 1H), 7.81-7.84(m, 3H), 8.07(s, 1H)

Example 42

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indol-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

4-Benzyloxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indole (5.0 g) obtained in Starting Material Synthesis Example 45 was dissolved in a mixed solvent (500 ml) of methanol - dimethylformamide (3:2) and 5% palladium-carbon (0.5 g) was added. The mixture was stirred for 5 hr under a hydrogen flow. The catalyst was removed by filtration through celite and the filtrate was concentrated under reduced pressure. To a

solution of the obtained 4-hydroxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)indole in dimethylformamide were added (S)-glycidyl nosylate (4 g) and potassium carbonate (4.2 g), and the mixture was heated at 50°C for 5 hr. The reaction mixture was poured into ice water and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (chloroform/methanol) to give yellow crystals (1 g). The yellow crystals and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (10 ml) and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.54 g) as yellow crystals, melting point 215-217°C.

¹H-NMR (DMSO-d₆) δ: 1.78-1.83 (m, 4H), 2.22-2.25 (m, 2H), 2.51-2.63 (m, 3H), 2.58 (s, 3H), 3.05-3.13 (m, 2H), 4.05 (m, 1H), 4.16 (m, 2H), 4.89 (bs, 1H), 6.58 (d, J=7.8, 1H), 7.04 (d, J=7.8, 1H), 7.13-7.19 (m, 2H), 7.42 (m, 3H), 7.70 (s, 1H), 7.82 (m, 3H), 12.16 (s, 1H)

Example 43

(S)-3-(4-(naphthalen-2-yl)piperidino)-1-(2-(5-phenyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-7-yloxy)-2-propanol

25 To a solution of 2-(7-methoxybenzo(b)furan-2-yl)-5-phenyl-1,3,4-oxadiazole (3.7 g), obtained in Starting Material Synthesis Example 47, in methylene chloride (100 ml) was added dropwise boron tribromide (4 ml) with stirring at -8°C. The mixture was then stirred for 1 hr under ice-cooling, and the
30 reaction mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give yellow crystals (2.7 g) of 7-hydroxy-2-(5-phenyl-1,3,4-oxadiazol-2-yl)benzo(b)furan. This compound and (S)-glycidyl nosylate (2.6

g) were dissolved in dimethylformamide (50 ml) and potassium carbonate (2.8 g) was added. The mixture was heated at 50°C for 2 hr. The reaction mixture was poured into ice water and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (1.7 g). The oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (20 ml) and the mixture was refluxed under heating for 1 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (2.3 g) as pale-yellow crystals, melting point 78-80°C. ¹H-NMR (DMSO-d₆) δ: 1.77-1.83 (m, 4H), 2.21-2.23 (m, 2H), 2.51-2.66 (m, 3H), 3.05-3.14 (m, 2H), 4.18 (m, 2H), 4.33 (m, 1H), 5.05 (bs, 1H), 7.18 (d, J=7.8, 1H), 7.32 (t, J=7.8, 1H), 7.38-7.44 (m, 4H), 7.65-7.70 (m, 4H), 7.80-7.84 (m, 3H), 7.90 (s, 1H), 8.18 (m, 2H)

Example 44

(S)-1-(2-(3-methyl-1,2,4-oxadiazol-5-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-3-methyl-1,2,4-oxadiazole (0.46 g) obtained in Starting Material Synthesis Example 58 and 4-(naphthalen-2-yl)piperidine (0.43 g), a brown oil (1.0 g) was obtained. This compound was dissolved in ethyl acetate and 1N solution of hydrochloric acid in ether was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.33 g) as brown crystals, melting point 216-218°C (decomposition).

Example 45

(S)-1-(2-(3-methyl-1,2,4-oxadiazol-5-yl)benzo(b)thiophen-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1

using (S)-5-(4-glycidyloxybenzo(b)thiophen-2-yl)-3-methyl-
1,2,4-oxadiazole (1.5 g) obtained in Starting Material
Synthesis Example 61 and 4-(naphthalen-2-yl)piperidine (1.0 g),
the title compound (1.5 g) was obtained as brown crystals,
5 melting point 180-182°C.

Example 46

(S)-1-(2-(3-methyl-1,2,4-oxadiazol-5-yl)benzo(b)thiophen-4-
yloxy)-3-(4-(naphthalen-1-yl)piperidino)-2-propanol
hydrochloride

10 By the reactions in the same manner as in Example 1
using (S)-5-(4-glycidyloxybenzo(b)thiophen-2-yl)-3-methyl-
1,2,4-oxadiazole (0.73 g) obtained in Starting Material
Synthesis Example 61 and 4-(naphthalen-1-yl)piperidine (1.0 g),
a brown oil (1.5 g) was obtained as brown crystals. This
15 compound was dissolved in ethyl acetate and 1N solution of
hydrochloric acid in ether was added. The precipitated
crystals were collected by filtration and dried to give the
title compound (0.5 g) as pale-yellow crystals, melting point
235°C or higher (decomposition).

20 **Example 47**

(S)-1-(2-(1,5-dimethylpyrazol-3-yl)benzo(b)furan-4-yloxy)-3-(4-
(naphthalen-2-yl)piperidino)-2-propanol 1/4 hydrate

By the reactions in the same manner as in Example 1
using (S)-3-(4-glycidyloxybenzo(b)furan-2-yl)-1,5-
25 dimethylpyrazole (0.2 g) obtained in Starting Material
Synthesis Example 63 and 4-(naphthalen-2-yl)piperidine (0.15 g),
the title compound (0.16 g) was obtained, melting point 155-
157°C.









Example 48

30 (S)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-7-yloxy)-3-(4-
(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Starting
Material Synthesis Example 1 using 2-(7-hydroxybenzo(b)furan-2-
yl)-5-methyloxazole (2.0 g) obtained in Starting Material

Synthesis Example 65 and (S)-glycidyl nosylate (1.8 g), (S)-7-glycidyoxy-2-(5-methyloxazol-2-yl)benzo(b)furan (1.5 g) was obtained. Then, by the reactions in the same manner as in Example 1 using 4-(naphthalen-2-yl)piperidine (0.7 g), the
5 title compound (0.26 g) was obtained, melting point 147-149°C.

The structural formulas of the compounds obtained in Examples 41 to 48 are shown in the following.



Example 49

(S)-1-(2-(3-methylisoxazol-5-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

5-(7-Methoxybenzo(b)furan-2-yl)-3-methylisoxazole (2.04 g) obtained in Starting Material Synthesis Example 66 was dissolved in dichloromethane (30 ml) and boron tribromide (3 ml) was added dropwise with stirring at -40°C. The mixture was then stirred for 4 hr under ice-cooling and the reaction mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give red crystals (1.96 g) of 5-(7-hydroxybenzo(b)furan-2-yl)-3-methylisoxazole. This compound and (S)-glycidyl nosylate (2.5 g) were dissolved in dimethylformamide (20 ml) and potassium carbonate (2.48 g) was added. The mixture was heated at 50°C for 3 hr. The reaction mixture was poured into ice water and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (2.38 g). The oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (20 ml) and the solution was refluxed under heating for 1 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (2.93 g) as an oil.

¹H-NMR (DMSO-d₆) δ: 1.93-2.25 (m, 4H), 2.33 (s, 3H), 2.75-3.35 (m, 5H), 3.65 (m, 2H), 4.27 (m, 2H), 4.48 (m, 1H), 5.00 (bs, 1H), 6.91 (s, 1H), 7.11 (d, J=7.8, 1H), 7.27 (t, J=7.8, 1H), 7.34 (d, J=7.8, 1H), 7.45-7.54 (m, 4H), 7.74 (s, 1H), 7.88 (m, 3H)

Example 50

(S)-1-(2-(2-methylthiazol-4-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Starting

Material Synthesis Example 5 using 4-(4-methoxybenzo(b)furan-2-yl)-2-methylthiazole (2.7 g) obtained in Starting Material Synthesis Example 67 and boron tribromide (7.5 g), 4-(4-hydroxybenzo(b)furan-2-yl)-2-methylthiazole (2.0 g) was
5 obtained as yellow crystals. By the reactions in the same manner as in Starting Material Synthesis Example 2 using this compound, (S)-glycidyl nosylate (2.9 g) and potassium carbonate (3.1 g), (S)-4-(4-glycidyoxybenzo(b)furan-2-yl)-2-methylthiazole (2.1 g) was obtained as a brown oil. By the
10 reactions in the same manner as in Example 1 using the brown oil and 4-(naphthalen-2-yl)piperidine (1.5 g), the title compound (0.3 g) was obtained as white crystals, melting point 148-150°C

Example 51

15 (S)-1-(2-(2-(5-methyl-1,3,4-oxadiazol-2-yl)vinyl)phenoxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

To a solution (20 ml) of 2-(2'-hydroxystyryl)-5-methyl-1,3,4-oxadiazole (1.5 g) obtained in Starting Material Synthesis Example 68 in DMF was added potassium carbonate (2.0
20 g), and then (S)-glycidyl nosylate (1.9 g) was added. The mixture was stirred at 40°C for 3 hr. The reaction mixture was concentrated under reduced pressure and water was added. The mixture was extracted with ethyl acetate, and the organic layer was dried over anhydrous sodium sulfate and concentrated under
25 reduced pressure to give an oil (1.3 g). To the oil (1.3 g) was added methanol (50 ml), and then 4-(naphthalen-2-yl)piperidine (1.0 g) was added. The mixture was refluxed under heating for 3 hr. After concentration, the residue was purified by silica gel column chromatography
30 (chloroform/methanol) to give the title compound (1.0 g) as white crystals, melting point 105-106°C.

Example 52

(S)-1-(2-(2-(benzothiazol-2-yl)vinyl)phenoxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

5 To a solution (50 ml) of 2-(2'-hydroxystyryl)-
benzothiazole (2.5 g) obtained in Starting Material Synthesis
Example 69 in DMF was added potassium carbonate (5.0 g), and
then (S)-glycidyl nosylate (2.4 g) was added. The mixture was
10 stirred at 50°C for 2 hr. The reaction mixture was
concentrated under reduced pressure and water was added. The
mixture was extracted with ethyl acetate, and the organic layer
was dried over anhydrous sodium sulfate and concentrated under
reduced pressure to give yellow crystals of (S)-2-(2'-
15 glycidyloxy)styrylbenzothiazole (2.7 g). To the yellow
crystals (1.5 g) was added methanol (50 ml), and then 4-
(naphthalen-2-yl)piperidine (1.0 g) was added. The mixture was
refluxed under heating for 3 hr. After concentration, the
residue was purified by silica gel column chromatography
20 (chloroform/methanol) to give white crystals (1.3 g), melting
point 125-127°C.

Example 53

(S)-1-(2-(2-(benzothiazol-2-yl)vinyl)phenyloxy)-3-(4-
(naphthalen-1-yl)piperidino)-2-propanol

20 By the reactions in the same manner as in Example 53
using (S)-2-(2'-glycidyloxystyryl)benzothiazole (0.9 g) and 4-
(naphthalen-1-yl)piperidine (0.6 g), the title compound (0.98
g) was obtained as white crystals, melting point 146-148°C.

Example 54

25 (S)-1-(2-(2-(3-methyl-1,2,4-oxadiazol-5-yl)vinyl)phenyloxy)-3-
(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride

To a solution (50 ml) of 5-(2'-hydroxystyryl)-3-methyl-
1,2,4-oxadiazole (2.0 g) obtained in Starting Material
Synthesis Example 70 in DMF was added potassium carbonate (3.0
30 g), and then (S)-glycidyl nosylate (2.6 g) was added. The
mixture was stirred at 50°C for 2 hr. The reaction mixture was
concentrated under reduced pressure and water was added. The
mixture was extracted with ethyl acetate and the organic layer
was dried over anhydrous sodium sulfate. The solvent was

evaporated under reduced pressure to give oily (S)-5-(2'-glycidyloxystyryl)-3-methyl-1,2,4-oxadiazole (2.2 g). This compound (1.2 g) was dissolved in methanol (50 ml), and 4-(naphthalen-2-yl)piperidine (1.0 g) was added. The mixture was
5 refluxed under heating for 3 hr. After concentration, the concentrate was purified by silica gel column chromatography (chloroform/methanol), and 1 M solution of hydrochloric acid in methanol was added to the residue obtained. The precipitated crystals were collected by filtration and dried to give the
10 title compound (1.2 g) as white crystals, melting point 184-186°C.

Example 55

(S)-1-(2-(2-(3-methyl-1,2,4-oxadiazol-5-yl)vinyl)phenoxy)-3-(4-(naphthalen-1-yl)piperidino)-2-propanol hydrochloride

15 By the reactions in the same manner as in Example 3 using 5-(2'-hydroxystyryl)-3-methyl-1,2,4-oxadiazole (1.0 g) obtained in Starting Material Synthesis Example 70 and 4-(naphthalen-1-yl)piperidine (1.0 g), the title compound (0.62 g) was obtained as white crystals, melting point 227-229°C
20 (decomposition).

Example 56

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-yl methyl ketone maleate

By the reactions in the same manner as in Example 3
25 using (S)-4-glycidyloxybenzo(b)furan-2-yl methyl ketone (0.52 g) obtained in Starting Material Synthesis Example 71 and 4-(naphthalen-2-yl)piperidine (0.47 g), (S)-4-(2-hydroxy-3-(4-naphthalen-2-yl)piperidino)propyloxy)benzo(b)furan-2-yl methyl ketone (0.87 g) was obtained as a brown oil. This was
30 dissolved in ethyl acetate and maleic acid (0.22 g) was added. The precipitated crystals were recrystallized from a mixed solvent of isopropanol - ethyl acetate to give the title compound (0.76 g) as pale-yellow crystals, melting point 153-

155°C.

Example 57

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-3-methylbenzo(b)furan-2-yl methyl ketone maleate

5 By the reactions in the same manner as in Example 3 using (S)-4-glycidyloxy-3-methylbenzo(b)furan-2-yl methyl ketone (0.60 g) obtained in Starting Material Synthesis Example 72 and 4-(naphthalen-2-yl)piperidine (0.51 g), (S)-4-(2-hydroxy-3-(4-naphthalen-2-yl)piperidino)propyloxy)-3-
10 methylbenzo(b)furan-2-yl methyl ketone (1.1 g) was obtained as a brown oil. This was dissolved in ethyl acetate and maleic acid (0.25 g) was added. The precipitated crystals were recrystallized from a mixed solvent of isopropanol - ethyl acetate to give the title compound (0.82 g) as pale-yellow
15 crystals, melting point 163-164°C.


Example 58

1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-yl)ethanol


(S)-4-(2-Hydroxy-3-(4-(naphthalen-2-yl)piperidino)-
20 propyloxy)benzo(b)furan-2-yl methyl ketone (0.30 g) obtained in Example 56 was dissolved in methanol and sodium borohydride (30 mg) was added at room temperature. The mixture was stirred for 20 min. To the reaction mixture was added saturated aqueous ammonium chloride solution and the solvent was evaporated under
25 reduced pressure. The obtained residue was dissolved in ethyl acetate, and the mixture was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give the title compound (0.24 g) as brown crystals, melting point 143-144°C.

30 The structural formulas of the compounds obtained in Examples 49 to 58 are shown in the following.


49




50




51




52




53




54




55




56



57



58



Example 59

(S)-5-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-3-morpholinomethyl-2-chromenone

Red crystals (2 g) of 5-hydroxy-3-morpholinomethyl-2-chromenone and (S)-glycidyl nosylate (2 g) were dissolved in dimethylformamide (20 ml) and potassium carbonate (3 g) was added. The mixture was heated at 50°C for 5 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (1.10 g). The oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (20 ml) and the mixture was refluxed under heating for 3 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.63 g) as an oil.

¹H-NMR(DMSO-d₆)δ: 2.09-2.22(m, 4H), 2.58(m, 2H), 2.75-3.35(m, 5H), 3.64(m, 8H), 4.01(s, 2H), 4.15(m, 2H), 4.46(m, 1H), 5.00(bs, 1H), 6.99(m, 2H), 7.46-7.57(m, 4H), 7.75(s, 1H), 7.88(m, 3H), 8.31(s, 1H)

Example 60

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-7-yloxy)-3-(4-(naphthalen-1-yl)piperidino)-2-propanol

7-Methoxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan (2 g) was dissolved in dichloromethane (50 ml) and boron tribromide (2 ml) was added dropwise with stirring at -8°C. The mixture was then stirred for 1 hr under ice-cooling and the reaction mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give red crystals (1.5 g) of 7-hydroxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan. This compound and (S)-glycidyl nosylate (2 g) were dissolved in DMF (100 ml) and potassium carbonate (11

g) was added. The mixture was stirred at 50°C for 2 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (2 g). The oily product and 4-(naphthalen-1-yl)piperidine were dissolved in methanol (20 ml) and the mixture was refluxed under heating for 1 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (1.0 g) as a pale-yellow oil.

¹H-NMR(DMSO-d₆)δ: 1.77-1.83(m, 4H), 2.20-2.25(m, 2H), 2.47-2.66(m, 3H), 2.62(s, 3H), 3.04-3.13(m, 2H), 4.17(m, 2H), 4.30(m, 1H), 5.02(bs, 1H), 7.17(d, J=7.8, 1H), 7.32(t, J=7.8, 1H), 7.40(d, J=7.8, 1H), 7.50-7.58(m, 4H), 7.74(s, 1H), 7.81(d, J=7.8, 1H), 7.93(d, J=7.8, 1H), 8.23(d, J=7.8, 1H)

Example 61

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)1H-indol-4-yloxy)-3-(4-(naphthalen-1-yl)piperidino)-2-propanol

To a solution of 4-hydroxy-2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indole in dimethylformamide were added (S)-glycidyl nosylate (2 g) and potassium carbonate (2 g), and the mixture was heated at 50°C for 5 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (chloroform/methanol) to give yellow crystals (0.5 g). The yellow crystals and 4-(naphthalen-1-yl)piperidine were dissolved in methanol (10 ml) and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography

(chloroform/methanol) to give the title compound (0.36 g) as yellow crystals, melting point 203-205°C.

¹H-NMR(DMSO-d₆)δ: 1.81-1.86(m, 4H), 2.33-2.39(m, 2H), 2.51-2.66(m, 3H), 2.58(s, 3H), 3.08-3.16(m, 2H), 4.05(m, 1H), 4.16(m, 2H), 4.92(bs, 1H), 6.58(d, J=7.8, 1H), 7.05(d, J=7.8, 1H), 7.13-7.19(m, 2H), 7.41-7.56(m, 4H), 7.75(d, J=7.8, 1H), 7.90(d, J=7.8, 1H), 8.14(d, J=7.8, 1H), 12.16(s, 1H)

Example 62

(S)-4-(2-acetoxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-methoxy-N-methylbenzo(b)furan-2-carboxamide maleate

(S)-4-(2-Hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-N-methoxy-N-methylbenzo(b)furan-2-carboxamide (0.40 g) obtained in Example 6 was dissolved in pyridine (20 ml) and acetic anhydride (10 ml) was added at room temperature. The mixture was stood for one day. The solvent was evaporated under reduced pressure and the obtained residue was purified by silica gel column chromatography (chloroform/methanol) to give (S)-4-(2-acetoxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N-methoxy-N-methylbenzo(b)furan-2-carboxamide (0.34 g) as a brown oil. This was dissolved in ethanol and maleic acid (0.10 g) was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.25 g) as pale-yellow crystals, melting point 125-127°C.

Example 63

ethyl (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzo(b)furan-2-carboxylate

By the reactions in the same manner as in Example 1 using ethyl (S)-4-glycidyloxybenzo(b)furan-2-carboxylate (3.3 g) and 4-(naphthalen-2-yl)piperidine (2.7 g), the title compound (5.1 g) was obtained as a brown oil.

¹H-NMR(CDCl₃)δ: 1.42(t, J=7.3, 3H), 1.87-1.99(m, 4H), 2.20(t, J=3.1, 1H), 2.50-2.54(m, 1H), 2.63-2.74(m, 3H), 3.05(brd, J=10.7, 1H), 3.23(brd, J=11.2, 1H), 4.13-4.25(m, 3H), 4.45(q, J=7.3, 2H), 6.72(d, J=8.3, 1H), 7.21(d, J=8.3, 1H), 7.35-7.49(m,

4H), 7.67(s, 1H), 7.68(d, J=6.3, 1H), 7.81(d, J=8.3, 3H)

Example 64

4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1H-indole-2-carboxamide

5 By the reactions in the same manner as in Example 1 using 4-glycidyloxy-1H-indole-2-carboxamide (1.8 g) and 4-(naphthalen-2-yl)piperidine (1.4 g), the title compound (1.8 g) was obtained as white crystals, melting point 200-202°C.

Example 65

10 (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide L-tartaric acid

By the reactions in the same manner as in Example 1 using 4-(glycidyloxy)benzo(b)thiophene-N,N-dimethylcarboxamide (3.5 g) and 4-(naphthalen-2-yl)piperidine (2.0 g), an oil (2.5
15 g) was obtained. This was dissolved in a solution of L-tartaric acid (2.0 g) in ethanol. The precipitated crystals were collected by filtration and dried to give the title compound (1.4 g) as white crystals, melting point 173-175°C.

Example 66

20 (S)-1-(7-(2-hydroxy-3-(5,6-dihydro-4-(naphthalen-2-yl)-2H-pyridin-1-yl)propyloxy)benzo(b)furan-2-ylcarbonyl)pyrrolidine

By the reactions in the same manner as in Example 1 using (S)-1-(7-glycidyloxybenzo(b)furan-2-ylcarbonyl)-pyrrolidine (2.1 g) and 5,6-dihydro-4-(naphthalen-2-yl)-2H-
25 pyridine (1.8 g), the title compound (2.8 g) was obtained as white crystals, melting point 114-116°C.

Example 67

(S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-1-isopropyl-N,N-dimethylindole-2-carboxamide

30 By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-isopropylindole-2-carboxylic acid (2.5 g), dimethylamine hydrochloride (0.63 g), triethylamine (2.1 ml) and diethyl cyanophosphate (0.93 ml), the title compound (2.0

g) was obtained as a yellow oil.

¹H-NMR(CDCl₃)δ: 1.62(d, J=6.8, 6H), 1.94-1.97(m, 4H), 2.24(t, J=3.1, 1H), 2.44-2.54(m, 1H), 2.61-2.76(m, 3H), 3.05(brd, J=10.7, 1H), 3.15(s, 6H), 3.23(brd, J=11.2, 1H), 4.13-4.29(m, 3H), 4.79(q, J=6.8, 1H), 6.54(d, J=6.8, 1H), 6.67(s, 1H), 7.13-7.15(m, 2H), 7.38-7.46(m, 3H), 7.66(s, 1H), 7.79(d, J=8.3, 3H)











Example 68

(S)-1-(4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-isopropylindole-2-carbonyl)pyrrolidine maleate

- 10 By the reactions in the same manner as in Example 3 using (S)-4-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)-1-isopropylindole-2-carboxylic acid (2.5 g), pyrrolidine (0.44 g), triethylamine (2.1 ml) and diethyl cyanophosphate (0.93 ml), a brown oil (2.1 g) was obtained.
- 15 This was dissolved in ethanol and maleic acid (0.4 g) was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.2 g) as pale-yellow crystals, melting point 154-155°C.

The structural formulas of the compounds obtained in

20 Examples 59 to 68 are shown in the following.



Example 69

(S)-1-(5-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidin-1-yl)propyloxy)-2H-chromen-3-ylcarbonyl)pyrrolidine

Red crystals (2.0 g) of 1-(5-hydroxy-2H-chromen-3-ylcarbonyl)pyrrolidine and (S)-glycidyl nosylate (2.0 g) were dissolved in dimethylformamide (20 ml), and potassium carbonate (3 g) was added. The mixture was heated at 50°C for 3 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (3.27 g). The oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (20 ml) and the mixture was refluxed under heating for 3 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.12 g) as a brown oil.

¹H-NMR(CDCl₃)δ: 1.91-2.02(m, 8H), 2.17(m, 2H), 2.48-2.70(m, 3H), 2.96(m, 1H), 3.15(m, 1H), 3.54(m, 4H), 3.73(bs, 1H), 4.00-4.13(m, 3H), 4.87(s, 2H), 6.47(d, J=7.8Hz, 1H), 6.50(d, J=7.8Hz, 1H), 7.11(t, J=7.8Hz, 1H), 7.16(s, 1H), 7.37(m, 3H), 7.64(s, 1H), 7.78(m, 3H)

By the same manner as in the above-mentioned Example, the following compounds can be synthesized.

Example 70

(S)-1-(2-(5-methyloxazol-2-yl)-1H-indol-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

Example 71

(S)-1-(2-(5-methyloxazol-2-yl)-1H-indol-4-yloxy)-3-(4-(4-chlorophenyl)piperidino)-2-propanol

Example 72

(S)-1-(2-(5-methyloxazol-2-yl)-1H-indol-4-yloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol

Example 73

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indol-4-yloxy)-3-(4-(4-chlorophenyl)piperidino)-2-propanol

Example 74

5 (S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indol-4-yloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol

Example 75

(S)-1-(2-(5-methyl-1H-imidazol-2-yl)-1H-indol-4-yloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol

10 **Example 76**

(S)-1-(2-(3-methyl-1H-pyrazol-5-yl)-1H-indol-4-yloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol

Example 77

15 (S)-1-(2-(3-methylisoxazol-5-yl)-1H-indol-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol


Example 78

(S)-1-(2-(5-methyloxazol-2-yl)-1H-indol-4-yloxy)-3-(4-(4-methylphenyl)piperidino)-2-propanol


The structural formulas of the compounds obtained in

20 Examples 69 to 78 are shown in the following.


69




70




71




72




73




74




75




76



77



78



Example 79

(R)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)-1H-indol-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

Example 80

5 (S)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

Example 81

(S)-3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol dihydrochloride

10 2-(4-Hydroxybenzo(b)furan)-5-methyloxazole (11.0 g) and (S)-glycidyl nosylate (13.0 g) were dissolved in dimethylformamide (100 ml) and potassium carbonate (15.0 g) was added. The mixture was stirred at room temperature for 10 hr. The reaction mixture was poured into ice water and extracted
15 with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oil (10.0 g). The oil and 4-(3,4-dichlorophenyl)piperidine were dissolved in methanol (100 ml)
20 and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol). The obtained yellow oil (10 g) was dissolved in acetone and hydrochloric acid was added to give a
25 hydrochloride. Recrystallization from ethanol gave the title compound (7.0 g) as pale-yellow crystals, melting point 190°C (decomposition).

¹H-NMR(DMSO-d₆)δ: 2.02-2.24(m, 4H), 2.43(s, 3H), 2.92(m, 1H), 3.20(m, 2H), 3.35-3.48(m, 2H), 3.71-3.81(m, 2H), 4.13-4.23(m,
30 2H), 4.57(m, 1H), 6.89(d, J=7.8, 1H), 7.08(s, 1H), 7.26-7.31(m, 2H), 7.37(t, J=7.8, 1H), 7.50(s, 1H), 7.56-7.67(m, 2H), 10.37(bs, 1H)

Example 82

(S)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(4-

chlorophenyl)piperidino)-2-propanol

Example 83

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(4-chlorophenyl)piperidino)-2-propanol

5 **Example 84**

(R)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

Example 85

10 (S)-1-(2-(3-methylisoxazol-5-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol










Example 86

(S)-1-(2-(5-methylthiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(4-chlorophenyl)piperidino)-2-propanol

Example 87

15 (S)-1-(2-(5-methylthiazol-2-yl)-1H-indol-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol

The structural formulas of the compounds obtained in Examples 79 to 87 are shown in the following.



Example 88

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride 1/4 hydrate

5 By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (23.0 g) obtained in the same manner as in Starting Material Synthesis Example 39 and 4-(3,4-dichlorophenyl)-piperidine (18.6 g), a brown oil (39.0 g) was obtained. This
10 was dissolved in ethanol. A solution of hydrochloric acid in ether was added and the mixture was allowed to stand. The precipitated crystals were collected by filtration and dried to give the title compound (23.5 g) as pale-yellow crystals, melting point 230-231°C.

15 **Example 89**

(S)-1-(4-(6-methoxynaphthalen-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-
20 oxadiazole (1.4 g) obtained in Starting Material Synthesis Example 39 and 4-(6-methoxynaphthalen-2-yl)piperidine (1.2 g), crude crystals were obtained. This was recrystallized from ethyl acetate to give the title compound (1.2 g) as white crystals, melting point 156-158°C.

25 **Example 90**

(S)-1-(4-(3,4-methylenedioxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride monohydrate

By the reactions in the same manner as in Example 1
30 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.50 g) obtained in Starting Material Synthesis Example 39 and 4-(3,4-methylenedioxyphenyl)piperidine (0.36 g), a brown oil (0.42 g) was obtained. This was dissolved in acetone and a solution of hydrochloric acid in ether was added.

The solvent was evaporated under reduced pressure and the resulting crude crystals were recrystallized from a mixed solvent of isopropanol - ethyl acetate (2:1) to give the title compound (0.27 g) as pale-yellow crystals, melting point 200-
5 202°C.

Example 91

(S)-1-(4-(3,4-dimethylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride monohydrate

10 By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.50 g) obtained in Starting Material Synthesis Example 39 and 4-(3,4-dimethylphenyl)piperidine (0.33 g), a brown oil (0.64 g) was obtained. This was dissolved in acetone
15 and a solution of hydrochloric acid in ether was added. The solvent was evaporated under reduced pressure and the resulting crude crystals were recrystallized from a mixed solvent of isopropanol - isopropyl ether (2:1) to give the title compound (0.33 g) as pale-yellow crystals, melting point 150-152°C.

20 **Example 92**

(S)-3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

The yellow oil (0.90 g) obtained by the reactions in the
25 same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole (0.50 g) obtained in Starting Material Synthesis Example 76 and 4-(3,4-dichlorophenyl)piperidine (0.40 g) was dissolved in acetone and a solution of hydrochloric acid in ether was added to give a
30 hydrochloride. Recrystallization from a mixed solvent of isopropanol - isopropyl ether gave the title compound (0.34 g) as white crystals, melting point 148-150°C.

Example 93

(S)-3-(4-(3,4-dimethylphenyl)piperidino)-1-(2-(5-ethyl-1,3,4-

oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride
1/2 hydrate

A yellow oil (5.0 g) obtained by the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo-
5 (b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole (3.0 g) obtained in Starting Material Synthesis Example 76 and 4-(3,4-dimethylphenyl)piperidine (2.0 g) was dissolved in a mixed solvent of acetone - ethyl acetate, and a solution of hydrochloric acid in ether was added to give a hydrochloride.
10 Recrystallization from a mixed solvent of acetone - ethyl acetate gave the title compound (2.0 g) as pale-yellow crystals, melting point 178-180°C.

Example 94

(S)-1-(2-(3-methylisoxazol-5-yl)benzo(b)furan-4-yloxy)-3-(4-
15 (naphthalen-2-yl)piperidino)-2-propanol hydrochloride 1/4
hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-3-methylisoxazole (0.50 g) obtained in Starting Material Synthesis Example 79 and
20 4-(naphthalen-2-yl)piperidine (0.37 g), a brown oil (0.69 g) was obtained. This was dissolved in ethyl acetate and a solution of hydrochloric acid in ether was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.36 g) as white crystals, melting
25 point 152-154°C.









Example 95

(S)-1-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-(2-(3-
methylisoxazol-5-yl)benzo(b)furan-4-yloxy)-2-propanol
hydrochloride 1/4 hydrate

30 By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-3-methylisoxazole (0.50 g) obtained in Starting Material Synthesis Example 79 and 4-(3,4-dichlorophenyl)piperazine (0.40 g), a brown oil (0.60 g) was obtained. This was dissolved in isopropanol and a solution

of hydrochloric acid in ether was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.36 g) as brown crystals, melting point 250°C or higher.

5 The structural formulas of the compounds obtained in Examples 88 to 95 are shown in the following.



Example 96

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-thiadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

5 monohydrate

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-thiadiazole (0.35 g) obtained in Starting Material Synthesis Example 82 and 4-(3,4-dichlorophenyl)piperidine (0.28 g), a
10 brown oil (0.60 g) was obtained. This was dissolved in isopropanol and a solution of hydrochloric acid in ether was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.19 g) as pale-yellow crystals, melting point 220-222°C.

15 **Example 97**

(S)-1-(4-(3,4-dimethylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-thiadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride
monohydrate

By the reactions in the same manner as in Example 1
20 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-thiadiazole (0.35 g) obtained in Starting Material Synthesis Example 82 and 4-(3,4-dimethylphenyl)piperidine (0.32 g), a brown oil (0.50 g) was obtained. This was dissolved in isopropanol and a solution of hydrochloric acid in ether was
25 added. The precipitated crystals were collected by filtration and dried to give the title compound (0.21 g) as pale-yellow crystals, melting point 191-194°C.

Example 98

(R)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride
30 monohydrate

By the reactions in the same manner as in Example 1 using (R)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.43 g), obtained by the reactions in the same

manner as in Starting Material Synthesis Example 1 using (R)-glycidyl nosylate and 2-(4-hydroxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole, and 4-(3,4-dichlorophenyl)piperidine (0.35 g), a brown oil (0.70 g) was obtained. This was dissolved in
5 ethanol and a solution of hydrochloric acid in ether was added. The mixture was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound (0.28 g) as pale-yellow crystals, melting point 230-231°C.

10 **Example 99**

(S)-1-(4-(3-chlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride
1/2 hydrate

By the reactions in the same manner as in Example 1
15 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.50 g) obtained in Starting Material Synthesis Example 39 and 4-(3-chlorophenyl)piperidine (0.32 g), a brown oil (0.70 g) was obtained. This was dissolved in ethanol and a solution of hydrochloric acid in ether was added. The mixture
20 was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound (0.09 g) as yellow crystals, melting point 170-172°C.

Example 100

(S)-1-(4-(4-chlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride
3/2 hydrate

By the reactions in the same manner as in Example 1
using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.50 g) obtained in Starting Material Synthesis
30 Example 39 and 4-(4-chlorophenyl)piperidine (0.32 g), a brown oil (0.62 g) was obtained. This was dissolved in ethanol and a solution of hydrochloric acid in ether was added. The mixture was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound

(0.32 g) as pale-yellow crystals, melting point 200-202°C.

Example 101

(S)-1-(4-(3,4-methylenedioxyphenyl)piperidino)-3-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

5 hydrochloride monohydrate

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole (0.80 g) obtained in Starting Material Synthesis Example 76 and 4-(3,4-methylenedioxyphenyl)piperidine (0.57 g),
10 a brown oil (1.02 g) was obtained. This was dissolved in isopropanol and a solution of hydrochloric acid in ether was added. The mixture was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound (0.36 g) as brown crystals, melting point 170-
15 173°C.

Example 102

(S)-1-(4-(2,4-dimethylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride monohydrate

20 By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.60 g) obtained in Starting Material Synthesis Example 39 and 4-(2,4-dimethylphenyl)piperidine (0.38 g), a brown oil (0.68 g) was obtained. This was dissolved in a mixed
25 solvent of isopropanol - isopropyl ether (1:1) and a solution of hydrochloric acid in ether was added. The mixture was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound (0.31 g) as white crystals, melting point 190-194°C.









30 **Example 103**

(S)-1-(4-phenylpiperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrobromide monohydrate

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-

oxadiazole (1.80 g) obtained in Starting Material Synthesis Example 39 and 4-phenylpiperidine (1.00 g), a brown oil (1.02 g) was obtained. This was dissolved in ethanol and hydrobromic acid was added. The mixture was allowed to stand and the
 5 precipitated crystals were collected by filtration and dried to give the title compound (1.84 g) as brown crystals, melting point 158-160°C.

The structural formulas of the compounds obtained in Examples 96 to 103 are shown in the following.



Example 104

(S)-1-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride 4/5 hydrate

5 By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-ethyl-1,3,4-oxadiazole (0.43 g) obtained in Starting Material Synthesis Example 76 and 4-(naphthalen-2-yl)piperidine (0.40 g), a brown oil (0.63 g) was obtained. This was dissolved in ethanol and a
10 solution of hydrogen chloride in ether was added. The mixture was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound (0.16 g) as brown crystals, melting point 113-115°C.

Example 105

15 (S)-1-(4-hydroxy-4-(naphthalen-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-
20 oxadiazole (1.0 g) obtained in Starting Material Synthesis Example 39 and 4-hydroxy-4-(naphthalen-2-yl)piperidine (0.77 g), a brown oil (1.6 g) was obtained. This was dissolved in ethanol and a solution of hydrogen chloride in ether was added. The mixture was allowed to stand and the precipitated crystals
25 were collected by filtration and dried to give the title compound (1.2 g) as white crystals, melting point 227-228°C.

Example 106

(S)-1-(4-(3,4-dichlorophenyl)piperazin-1-yl)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol
30 hydrochloride

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (0.80 g) obtained in Starting Material Synthesis Example 39 and 4-(3,4-dichlorophenyl)piperazine (0.65 g), a

brown oil (1.2 g) was obtained. This was dissolved in acetone and a solution of hydrogen chloride in ether was added. The mixture was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound
5 (0.4 g) as white crystals, melting point 232-233°C.

Example 107

(S)-1-(4-(3,4-dichlorophenyl)-3,6-dihydro-2H-pyridin-1-yl)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

10 By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole (1.7 g) obtained in Starting Material Synthesis Example 39 and 4-(3,4-dichlorophenyl)-3,6-dihydro-2H-pyridine (1.4 g), a brown oil (2.2 g) was obtained. This was dissolved
15 in ethanol and a solution of hydrogen chloride in ether was added. The mixture was allowed to stand and the precipitated crystals were collected by filtration and dried to give the title compound (1.5 g) as white crystals, melting point 204-207°C.

20 **Example 108**

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-ethylthiophen-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1
25 using (S)-2-(5-ethylthiophen-2-yl)-4-glycidyloxybenzo(b)furan (921 mg) obtained in Starting Material Synthesis Example 89 and 4-(3,4-dichlorophenyl)piperidine (777 mg), a colorless oil (1.51 g) was obtained. This was dissolved in methanol and 1 equivalent of hydrochloric acid was added. The mixture was
30 stirred for 15 min and the solvent was evaporated under reduced pressure. The residue was recrystallized from a mixed solvent of methanol - water and the precipitated crystals were collected by filtration and dried to give the title compound

(691 mg) as colorless crystals, melting point 183-185°C.

Example 109

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(1-methylimidazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol
5 dihydrochloride 5/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-4-glycidyloxy-2-(1-methylimidazol-2-yl)benzo(b)furan (649 mg) obtained in Starting Material Synthesis Example 92 and 4-(3,4-dichlorophenyl)piperidine (591 mg), a colorless
10 amorphous solid (521 mg) was obtained. This was dissolved in methanol and 2 equivalents of hydrochloric acid were added. The mixture was stirred for 15 min and the solvent was evaporated under reduced pressure. The residue was recrystallized twice from a mixed solvent of methanol - ethyl
15 acetate, and the precipitated crystals were collected by filtration and dried to give the title compound (397 mg) as colorless crystals, melting point >155°C.

Example 110

(S)-1-(2-(1-methylimidazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol
20 dihydrochloride 5/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-4-glycidyloxy-2-(1-methylimidazol-2-yl)benzo(b)furan (649 mg) obtained in Starting Material Synthesis Example 92 and
25 4-(naphthalen-2-yl)piperidine (542 mg), a colorless amorphous solid (547 mg) was obtained. This was dissolved in methanol and 2 equivalents of hydrochloric acid were added. The mixture was stirred for 15 min and the solvent was evaporated under reduced pressure. The residue was recrystallized from a mixed
30 solvent of methanol - ethyl acetate, and the precipitated crystals were collected by filtration and dried to give the title compound (295 mg) as colorless crystals, melting point >160°C.


Example 111

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-methyloxazol-2-yl)benzo(b)thiophen-4-yloxy)-2-propanol hydrochloride


By the reactions in the same manner as in Example 1
5 using (S)-4-glycidyloxy-2-(5-methyloxazol-2-yl)benzo(b)-
thiophene (735 mg) obtained in Starting Material Synthesis
Example 95 and 4-(3,4-dichlorophenyl)piperidine (589 mg), a
colorless amorphous solid (963 mg) was obtained. This was
dissolved in methanol and 2 equivalents of hydrochloric acid
10 were added. The mixture was stirred for 15 min and the solvent
was evaporated under reduced pressure. The residue was
recrystallized from methanol and the precipitated crystals were
collected by filtration and dried to give the title compound
(528 mg) as pale-yellow crystals, melting point >225°C
15 (decomposition).

The structural formulas of the compounds obtained in
Examples 104 to 111 are shown in the following.


104




105




106




107




108




109



110



111

**Example 112**

(S)-1-(2-(5-methyloxazol-2-yl)benzo(b)thiophen-4-yloxy)-3-(4-

(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-4-glycidyloxy-2-(5-methyloxazol-2-yl)benzo(b)-thiophene (735 mg) obtained in Starting Material Synthesis Example 95 and 4-(naphthalen-2-yl)piperidine (540 mg), a colorless amorphous solid (1.04 g) was obtained. This was recrystallized from ethyl acetate and the precipitated crystals were collected by filtration and dried to give the title compound (793 mg) as pale-yellow crystals, melting point 138-139°C.

Example 113

(S)-1-(4-(6-methoxynaphthalen-2-yl)piperidino)-3-(2-(5-methyloxazol-2-yl)benzo(b)thiophen-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-4-glycidyloxy-2-(5-methyloxazol-2-yl)benzo(b)-thiophene (735 mg) obtained in Starting Material Synthesis Example 95 and 4-(6-methoxynaphthalen-2-yl)piperidine (617 mg), a colorless amorphous solid (981 mg) was obtained. This was recrystallized from a mixed solvent of chloroform - hexane and the precipitated crystals were collected by filtration and dried to give the title compound (714 mg) as pale-yellow crystals, melting point 163-165.5°C

Example 114

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(4,4-dimethyloxazolin-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-4-glycidyloxy-2-(4,4-dimethyloxazolin-2-yl)benzo(b)furan (933 mg) obtained in Starting Material Synthesis Example 98 and 4-(3,4-dichlorophenyl)piperidine (748 mg), the title compound (1.12 g) was obtained as a pale yellow amorphous solid.

¹H-NMR(CDCl₃)δ: 1.41(s, 6H), 1.45-1.90(m, 5H), 2.12(br.t, J=12.0, 1H), 2.41(br.t, J=12.0, 1H), 2.45-2.65(m, 3H), 2.98(br.d, J=12.0, 1H), 3.14(br.d, J=12.0, 1H), 4.05-4.20(m,

3H), 4.14(s, 2H), 6.68(d, J=8.0, 1H), 7.05(dd, J=8.0, 1.0, 1H),
7.17(d, J=8.0, 1H), 7.25-7.40(m, 3H)

Example 115

(S)-1-(2-(4,4-dimethyloxazolin-2-yl)benzo(b)furan-4-yloxy)-3-
5 (4-(naphthalen-2-yl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1
using (S)-4-glycidyloxy-2-(4,4-dimethyloxazolin-2-
yl)benzo(b)furan (1.25 g) obtained in Starting Material
Synthesis Example 98 and 4-(naphthalen-2-yl)piperidine (914 mg),
10 the title compound (1.28 g) was obtained as a colorless
amorphous solid.

¹H-NMR(CDCl₃)δ: 1.41(s, 6H), 1.80-2.00(m, 4H), 2.19(br.t,
J=12.0, 1H), 2.40-2.55(m, 1H), 2.60-2.80(m, 3H), 3.00(br.d,
J=11.0, 1H), 3.19(br.d, J=11.0, 1H), 4.05-4.25(m, 3H), 4.13(s,
15 2H), 6.69(d, J=8.0, 1H), 7.17(d, J=8.0, 1H), 7.28(t, J=8.0,
1H) 7.35-7.45(m, 3H), 7.65(s, 1H), 7.75-7.85(m, 2H)

Example 116

(S)-1-(2-(4,4-dimethyloxazolin-2-yl)benzo(b)furan-4-yloxy)-3-
(4-(6-methoxynaphthalen-2-yl)piperidino)-2-propanol

20 By the reactions in the same manner as in Example 1
using (S)-4-glycidyloxy-2-(4,4-dimethyloxazolin-2-
yl)benzo(b)furan (933 mg) obtained in Starting Material
Synthesis Example 98 and 4-(6-methoxynaphthalen-2-yl)piperidine
(783 mg), a colorless amorphous solid was obtained. This was
25 recrystallized from ethyl acetate, and the precipitated
crystals were collected by filtration and dried to give the
title compound (871 mg) as colorless crystals, melting point
133-134°C.

Example 117

30 (S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-
ethylsulfonylbenzo(b)furan-4-yloxy)-2-propanol hydrochloride
1/4 hydrate

By the reactions in the same manner as in Example 1
using (S)-2-(ethylsulfonyl)-4-glycidyloxybenzo(b)furan (1.15 g)

obtained in Starting Material Synthesis Example 101 and 4-(3,4-dichlorophenyl)piperidine (936 mg), a colorless amorphous solid (1.11 g) was obtained. This was dissolved in methanol and 1 equivalent of hydrochloric acid was added. The mixture was stirred for 15 min and the solvent was evaporated under reduced pressure. The residue was recrystallized from a mixed solvent of acetone - ether, and the precipitated crystals were collected by filtration and dried to give the title compound (734 mg) as colorless crystals, melting point 135-140°C.

Example 118

(S)-1-(4-(3,4-dimethylphenyl)piperidino)-3-(2-ethylsulfonylbenzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-2-(ethylsulfonyl)-4-glycidyloxybenzo(b)furan (1.15 g) obtained in Starting Material Synthesis Example 101 and 4-(3,4-dimethylphenyl)piperidine (761 mg), the title compound (827 mg) was obtained as a yellowish brown oil.

¹H-NMR(CDCl₃)δ: 1.34(t, J=8.0, 3H), 1.40-1.95(m, 5H), 2.13(br.t, J=12.0, 1H), 2.22(s, 3H), 2.24(s, 3H), 2.40-2.70(m, 4H), 2.96(br.d, J=12.0, 1H), 3.14(br.d, J=12.0, 1H), 3.30(q, J=8.0, 2H), 4.05-4.20(m, 3H), 6.75(d, J=8.0, 1H), 6.96(d, J=8.0, 1H), 6.99(d, J=12.0, 1H), 7.17(d, J=8.0, 1H), 7.39(t, J=8.0, 1H), 7.66(s, 1H)

Example 119


(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(N,N-dimethylsulfamoyl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride 3/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-2-(N,N-dimethylsulfamoyl)-4-glycidyloxybenzo(b)furan (677 mg) obtained in Starting Material Synthesis Example 104 and 4-(3,4-dichlorophenyl)piperidine (524 mg), a colorless amorphous solid (1.01 g) was obtained. This was dissolved in methanol and 2 equivalents of hydrochloric acid were added. The mixture was stirred for 15 min and the solvent was


evaporated under reduced pressure. The residue was recrystallized from methanol, and the precipitated crystals were collected by filtration and dried to give the title compound (213 mg) as colorless crystals, melting point 222-
5 225°C.

The structural formulas of the compounds obtained in Examples 112 to 119 are shown in the following.


112




113




114




115




116




117



118



119



Example 120

(S)-1-(4-(3,4-dimethylphenyl)piperidino)-3-(2-(N,N-dimethylsulfamoyl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
5 using (S)-2-(N,N-dimethylsulfamoyl)-4-glycidyloxybenzo(b)furan
(677 mg) obtained in Starting Material Synthesis Example 104
and 4-(3,4-dimethylphenyl)piperidine (431 mg), a colorless
amorphous solid was obtained. This was recrystallized from a
mixed solvent of ethyl acetate - hexane, and the precipitated
10 crystals were collected by filtration and dried to give the
title compound (677 mg) as colorless crystals, melting point
146-148°C.

Example 121

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-ethyloxazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1
using (S)-2-(5-ethyloxazol-2-yl)-4-glycidyloxybenzo(b)furan
(500 mg) obtained in Starting Material Synthesis Example 108
and 4-(3,4-dichlorophenyl)piperidine (403 mg), a colorless
20 amorphous solid (854 mg) was obtained. This was dissolved in
methanol and 2 equivalents of hydrochloric acid were added.
The mixture was stirred for 15 min and the solvent was
evaporated under reduced pressure. The residue was
recrystallized from methanol, and the precipitated crystals
25 were collected by filtration and dried to give the title
compound (441 mg) as colorless crystals, melting point 232-
234°C.

Example 122

(S)-1-(4-(3,4-dimethylphenyl)piperidino)-3-(2-(5-ethyloxazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol dihydrochloride

By the reactions in the same manner as in Example 1
using (S)-2-(5-ethyloxazol-2-yl)-4-glycidyloxybenzo(b)furan
(500 mg) obtained in Starting Material Synthesis Example 108
and 4-(3,4-dimethylphenyl)piperidine (531 mg), a colorless

amorphous solid (1.03 g) was obtained. This was dissolved in methanol and 2 equivalents of hydrochloric acid were added. The mixture was stirred for 15 min and the solvent was evaporated under reduced pressure. The residue was

5 recrystallized from a mixed solvent of methanol-ethyl acetate-diisopropyl ether (2:1:1), and the precipitated crystals were collected by filtration and dried to give the title compound (468 mg) as pale-yellow crystals, melting point 86-89°C.

Example 123

10 (S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-isopropyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride monohydrate

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-isopropyl-1,3,4-oxadiazole (0.6 g) and 4-(3,4-dichlorophenyl)piperidine (0.46 g), an oil (0.75 g) was obtained. This was dissolved in ethanol and a solution of hydrogen chloride in dioxane was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.48 g) as white crystals,
20 melting point 142-144°C.

Example 124

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-phenyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-phenyl-1,3,4-oxadiazole (0.60 g) and 4-(3,4-dichlorophenyl)piperidine (0.50 g), an oil (0.85 g) was obtained. This was dissolved in ethanol and a solution of hydrogen chloride in dioxane was added. The precipitated crystals were collected by filtration
30 and dried to give the title compound (0.65 g) as white crystals, melting point 220-222°C.

Example 125

(S)-1-(4-(benzo(b)thiophen-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (0.4 g) and 4-(benzo(b)thiophen-2-yl)piperidine (0.35 g), an oil (0.65 g) was obtained. This was dissolved in ethanol and a solution of hydrogen chloride in dioxane was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.40 g) as white crystals, melting point 190-192°C.

Example 126

(S)-2-(4-(3,4-dimethylphenyl)piperidino)-1-(2-(2-methyloxazol-5-yl)benzo(b)furan-4-yloxy)-2-propanol dihydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyloxazole (0.60 g) and 4-(3,4-dimethylphenyl)piperidine (0.35 g), an oil (0.55 g) was obtained. This was dissolved in ethanol and a solution of hydrogen chloride in dioxane was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.19 g) as white crystals, melting point 79-81°C.

Example 127

(S)-1-(4-(2,3-dihydro-2-oxobenzimidazol-1-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-


propanol maleate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (1.0 g) and 4-(2,3-dihydro-2-oxobenzimidazol-1-yl)piperidine (1.0 g), an oil (1.4 g) was obtained. This was dissolved in acetone and maleic acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.68 g) as pale-yellow crystals, melting point 178-180°C.


The structural formulas of the compounds obtained in

Examples 120 to 127 are shown in the following.


120




121




122




123




124




125



126



127



Example 128

(S)-1-(4-(3,4-difluorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol p-toluenesulfonate 1/2 hydrate

5 By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (1.0 g) and 4-(3,4-difluoro)piperidine (1.0 g), an oil (1.7 g) was obtained. This was dissolved in acetone and p-toluenesulfonic acid was added. The precipitated crystals were
10 collected by filtration and dried to give the title compound (0.97 g) as pale-yellow crystals, melting point 80-82°C.

Example 129

(S)-1-(4-(3,4-dimethoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

15 By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (0.90 g) and 4-(3,4-dimethoxyphenyl)piperidine (0.90 g), the title compound (0.80 g) was obtained as a brown oil.
¹H-NMR(CDCl₃): 1.78-1.95(m, 4H), 2.16(t, 1H, J=1.8), 2.41-
20 2.60(m, 2H), 2.62-2.49(m, 5H), 3.00(d, 1H, J=11.2), 3.17(d, 1H, J=11.2), 3.86(s, 3H), 3.90(s, 3H), 4.15-4.30(m, 3H), 6.70-6.83(m, 4H), 7.22(d, 1H, J=8.3), 7.34(t, 1H, J=8.3), 7.61(s, 1H)

Example 130

25 (S)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-1-(4-(1,2,3,4-tetrahydronaphthalen-6-yl)piperidino)-2-propanol hydrochloride 1/4 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-
30 oxadiazole (1.0 g) and 4-(1,2,3,4-tetrahydronaphthalen-6-yl)piperidine (1.0 g), an oil (0.95 g) was obtained. This was dissolved in isopropanol and a solution of hydrogen chloride in dioxane was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.36 g) as

pale-yellow crystals, melting point 119-121°C.

Example 131

(S)-3-(4-(1,4-benzodioxan-6-yl)piperidino)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol
5 hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (1.0 g) and 4-(1,4-benzodioxan-6-yl)piperidine (1.0 g), an oil (1.45 g) was obtained. This was dissolved in
10 isopropanol and a solution of hydrogen chloride in dioxane was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.95 g) as white crystals, melting point 164-166°C.

Example 132

15 (S)-3-(4-(1,4-benzodioxan-6-yl)piperidino)-1-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride
1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-2-ethyl-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-
20 oxadiazole (1.0 g) and 4-(1,4-benzodioxan-6-yl)piperidine (1.0 g), an oil (1.55 g) was obtained. This was dissolved in a mixed solution of isopropanol - isopropyl ether (1:1) and a solution of hydrogen chloride in dioxane was added. The precipitated crystals were collected by filtration and dried to
25 give the title compound (0.71 g) as brown crystals, melting point 238-240°C.

Example 133

(S)-1-(4-(3,4-dichlorophenyl)piperidino)-3-(2-(5-dimethylamino-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

30 By the reactions in the same manner as in Example 1 using (S)-2-(dimethylamino)-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole (0.65 g) and 4-(3,4-dichlorophenyl)-piperidine (0.65 g), the title compound (0.38 g) was obtained

as brown crystals, melting point 209-211°C.

Example 134

(S)-3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(acetohydrazinocarbonyl)benzo(b)furan-4-yloxy)-2-propanol

5 N'-(4-Hydroxybenzo(b)furan-2-ylcarbonyl)acetohydrazide
(2.5 g) obtained in Starting Material Synthesis Example 109 and
(S)-glycidyl nosylate (2.3 g) were dissolved in
dimethylformamide (30 ml), and potassium carbonate (3.3 g) was
added. The mixture was stirred at room temperature for 15 hr.
10 The reaction mixture was poured into ice water and extracted
with ethyl acetate. The organic layer was washed with
saturated aqueous ammonium chloride solution, dried over
anhydrous sodium sulfate and concentrated under reduced
pressure to give a yellow oil (0.7 g). This oil and 4-(3,4-
15 dichlorophenyl)piperidine were dissolved in methanol (10 ml)
and the mixture was refluxed under heating for 3 hr. After
cooling, the solvent was evaporated under reduced pressure and
the residue was purified by silica gel column chromatography
(chloroform/methanol) to give the title compound (0.48 g) as
20 pale-yellow crystals, melting point 180°C (decomposition).
¹H-NMR(DMSO-d₆)δ: 1.72-1.88(m, 4H), 2.02(s, 3H), 2.14-2.17(m,
1H), 2.39-2.55(m, 2H), 2.64-2.66(m, 2H), 3.02(m, 1H), 3.18(m,
1H), 4.13-4.20(m, 2H), 4.54(m, 1H), 6.05(bs, 1H), 6.87(d, J=7.8,
1H), 7.06-7.09(m, 1H), 7.24(d, J=7.8, 1H), 7.32-7.39(m, 3H),
25 7.79(s, 1H), 9.92(s, 1H), 10.50(s, 1H)

Example 135









(S)-3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(5-methoxycarbonyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

30 5-Ethoxycarbonyl-2-(4-hydroxybenzo(b)furan-2-yl)-1,3,4-
oxadiazole (2.5 g) and (S)-glycidyl nosylate (2.3 g) were
dissolved in dimethylformamide (30 ml) and potassium carbonate
(2.45 g) was added. The mixture was stirred at room
temperature for 14 hr. The reaction mixture was poured into

ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give yellow crystals (1.7 g). This was
5 dissolved in methanol (20 ml) and 4-(3,4-dichlorophenyl)-piperidine (1.2 g) was added. The mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the
10 title compound (1.5 g) as pale-yellow crystals (ethyl ester was converted to methyl ester in methanol), melting point 160°C (decomposition).

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.71-1.89 (m, 4H), 2.14-2.20 (m, 1H), 2.45-2.57 (m, 2H), 2.64-2.66 (m, 2H), 3.02 (m, 1H), 3.18 (m, 1H), 4.11 (s, 3H),
15 5.05 (bs, 1H), 6.77 (d, $J=7.8$, 1H), 7.06-7.09 (m, 1H), 7.24 (d, $J=7.8$, 1H), 7.32-7.43 (m, 3H), 7.85 (s, 1H)

The structural formulas of the compounds obtained in Examples 128 to 135 are shown in the following.



Example 136

(S)-3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(5-hydroxymethyl-

1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

(S)-3-(4-(3,4-Dichlorophenyl)piperidino)-1-(2-(5-methoxycarbonyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol (1.3 g) obtained in Example 135 was dissolved in THF
5 (20 ml) and lithium borohydride (60 mg) was added with stirring under ice-cooling. The mixture was stirred for 1 hr at room temperature, then poured into ice-water, and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give yellow
10 crystals (1.2 g). The crystals were purified by silica gel column chromatography (chloroform/methanol) to give the title compound (100 mg) as pale-yellow crystals, melting point 158-160°C.

¹H-NMR (DMSO-d₆) δ: 1.62-1.72 (m, 4H), 2.09-2.15 (m, 2H), 2.50-
15 2.55 (m, 2H), 2.99-3.08 (m, 2H), 4.08 (m, 2H), 4.21 (m, 1H), 4.75 (d, J=8.0, 1H), 4.96 (bs, 1H), 6.01 (t, J=8.0, 1H), 6.92 (d, J=7.8, 1H), 7.25 (d, J=7.8, 1H), 7.33 (d, J=7.8, 1H), 7.42-7.53 (m, 3H), 7.80 (s, 1H)

Example 137

20 (S)-2-(4-(2-acetoxy-3-(4-(3,4-dichlorophenyl)piperidino)-propyloxy)benzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole hydrochloride

(S)-1-(4-(3,4-Dichlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol (1.0 g)
25 was dissolved in pyridine (20 ml) and acetic anhydride (10 ml), and the mixture was left standing at room temperature for 12 hr. The solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol). The obtained oil was dissolved in
30 ethanol and a solution of hydrogen chloride in ether was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.58 g) as pale-yellow crystals, melting point 163-166°C (decomposition).

Example 138

(S)-1-(2-(5-(1-methylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrochloride

5 By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1-methylethyl)-1,3,4-oxadiazole (0.8 g) and 4-(3,4-methylenedioxyphenyl)piperidine (0.7 g), an oil was obtained. This was dissolved in acetone and a hydrochloric acid-ethanol
10 solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.81 g) as white crystals, melting point 211-213°C.

Example 139

(R)-1-(2-(5-(1-methylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (R)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1-methylethyl)-1,3,4-oxadiazole (1.0 g) and 4-(3,4-methylenedioxyphenyl)piperidine (0.75 g), an oil was obtained.
20 This was dissolved in acetone and a hydrogen chloride-dioxane solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.61 g) as white crystals, melting point 209-211°C.

25 **Example 140**

(S)-1-(2-(5-(1-methylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride 3/4 hydrate

By the reactions in the same manner as in Example 1
30 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1-methylethyl)-1,3,4-oxadiazole (0.5 g) and 4-(naphthalen-2-yl)piperidine (0.5 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the

title compound (0.26 g) as white crystals, melting point 124-126°C

Example 141

(S)-1-(4-(3,4-dimethylphenyl)piperidino)-3-(2-(5-(1-methylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrobromide

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1-methylethyl)-1,3,4-oxadiazole (0.6 g) and 4-(3,4-dimethylphenyl)piperidine (0.6 g), an oil was obtained. This was dissolved in isopropanol, and 48% hydrobromic acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.48 g) as pale-yellow crystals, melting point 159-161°C.

Example 142

(S)-1-(2-(5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)-furan-4-yloxy)-3-(4-(1,4-benzodioxan-6-yl)piperidino)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1,1-dimethylethyl)-1,3,4-oxadiazole (0.8 g) and 4-(1,4-benzodioxan-6-yl)piperidine (0.70 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.55 g) as white crystals, melting point 197-199°C.









Example 143

(S)-1-(2-(5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1,1-dimethylethyl)-1,3,4-oxadiazole (1.0 g) and 4-(3,4-dichlorophenyl)piperidine (0.80 g), an oil was obtained. This

was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.55 g) as white crystals, melting point 207-209°C.

- 5 The structural formulas of the compounds obtained in Examples 136 to 143 are shown in the following.



Example 144

(S)-1-(2-(5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrochloride 1/2 hydrate

- 5 By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1,1-dimethylethyl)-1,3,4-oxadiazole (1.5 g) and 4-(3,4-methylenedioxyphenyl)piperidine (1.2 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added.
- 10 The precipitated crystals were collected by filtration and dried to give the title compound (1.33 g) as white crystals, melting point 204-206°C.

Example 145

- (S)-1-(2-(5-(2-methylpropyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrochloride
- 15

- By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(2-methylpropyl)-1,3,4-oxadiazole (0.62 g) and 4-(3,4-methylenedioxyphenyl)piperidine (0.4 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.51 g) as pale-yellow crystals, melting point 185-187°C.
- 20

Example 146

- (S)-1-(2-(5-(2-methylpropyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride
- 25

- By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxy)benzo(b)furan-2-yl)-2-(2-methylpropyl)-1,3,4-oxadiazole (0.62 g) and 4-(naphthalen-2-yl)piperidine (0.4 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the
- 30

title compound (0.35 g) as pale-yellow crystals, melting point 78-80°C.

Example 147

(S)-1-(4-(3,4-dimethylphenyl)piperidino)-3-(2-(5-(2-methylpropyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxy)benzo(b)furan-2-yl)-2-(2-methylpropyl)-1,3,4-oxadiazole (0.6 g) and 4-(3,4-dimethylphenyl)piperidine (0.4 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.45 g) as white crystals, melting point 156-158°C.

Example 148

(S)-1-(2-(5-(2-methylpropyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxy)benzo(b)furan-2-yl)-2-(2-methylpropyl)-1,3,4-oxadiazole (0.62 g) and 4-(3,4-dichlorophenyl)piperidine (0.46 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.23 g) as white crystals, melting point 98-100°C.

Example 149

(S)-1-((1-benzylpiperidin-4-yl)amino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol dihydrochloride monohydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (1.5 g) and 4-amino-1-benzylpiperidine (1.0 g), an oil was obtained. This was dissolved in acetone and

hydrochloric acid-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.2 g) as white crystals, melting point 260-262°C.

5 **Example 150**

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(4-methylphenyl)piperidino)-2-propanol hydrochloride 1/4 hydrate









By the reactions in the same manner as in Example 1
10 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (1.0 g) and 4-(4-methylphenyl)piperidine (0.75 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the
15 title compound (0.95 g) as white crystals, melting point 203-205°C.

Example 151

(S)-1-(4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol
20 hydrochloride

By the reactions in the same manner as in Example 1
using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (0.50 g) and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine (0.35 g), an oil was obtained. This was
25 dissolved in ethanol and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.55 g) as white crystals, melting point 198-200°C.

The structural formulas of the compounds obtained in
30 Examples 144 to 151 are shown in the following.



Example 152

(S)-1-(4-(2,3-dihydrobenzo(b) furan-5-yl)piperidino)-3-(2-(5-

hydroxymethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-hydroxymethyl-1,3,4-oxadiazole (0.58 g) and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine (0.40 g), the title compound (0.58 g) was obtained as white crystals, melting point 156-158°C.

Example 153

(R)-1-(4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrobromide

By the reactions in the same manner as in Example 1 using (R)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (0.25 g) and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine (0.18 g), an oil was obtained. This was dissolved in isopropanol and hydrogen chloride-ether solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.32 g) as white crystals, melting point 176-179°C.

Example 154

(S)-1-(4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-2-ethyl-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole (0.50 g) and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine (0.35 g), an oil was obtained. This was dissolved in a mixed solution of isopropanol-ethyl acetate (1:1) and a hydrogen chloride-ether solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.25 g) as white crystals, melting point 180-182°C.

Example 155

(S)-1-(4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-(1-

methylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol dihydrobromide 3/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1-methylethyl)-1,3,4-oxadiazole (0.50 g) and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine (0.35 g), an oil was obtained. This was dissolved in isopropanol and 48% hydrobromic acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.38 g) as yellow crystals, melting point 131-136°C (decomposition).

Example 156

(S)-1-(4-(chroman-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (1.0 g) and 4-(chroman-6-yl)piperidine (0.90 g), an oil was obtained. This was dissolved in isopropanol and hydrogen chloride-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.3 g) as white crystals, melting point 202-204°C.

Example 157

(S)-1-(4-(chroman-6-yl)piperidino)-3-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol (L)-tartrate

By the reactions in the same manner as in Example 1 using (S)-2-ethyl-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole (1.0 g) and 4-(chroman-6-yl)piperidine (0.90 g), an oil was obtained. This was dissolved in ethanol and (L)-tartaric acid-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.4 g) as white crystals, melting point 176-178°C.

Example 158

(S)-1-(4-(chroman-6-yl)piperidino)-3-(2-(5-(1-methylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

5 By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-(1-methylethyl)-1,3,4-oxadiazole (1.0 g) and 4-(chroman-6-yl)piperidine (0.90 g), an oil was obtained. This was dissolved in isopropanol and hydrogen chloride-ethanol solution
10 was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.3 g) as white crystals, melting point 183-185°C.









Example 159

ethyl (S)-4-(4-fluorophenyl)-1-(2-hydroxy-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)propyl)piperidine-4-carboxylate hydrobromide monohydrate

15

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (1.0 g) and ethyl 4-(4-fluorophenyl)piperidine-4-
20 carboxylate (0.82 g), an oil was obtained. This was dissolved in acetone, hydrobromic acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.66 g) as white crystals, melting point 132-135°C.

25 The structural formulas of the compounds obtained in Examples 152 to 159 are shown in the following.



Example 160

(S)-4-(3,4-dichlorophenyl)-1-(2-hydroxy-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)propyl)piperidine-4-carbonitrile hydrobromide monohydrate

5 By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (0.22 g) and 4-(3,4-dichlorophenyl)piperidine-4-carbonitrile (0.20 g), an oil was obtained. This was dissolved in acetone and hydrobromic acid was added. The precipitated
10 crystals were collected by filtration and dried to give the title compound (0.12 g) as brown crystals, melting point 265-268°C (decomposition).

Example 161

(S)-1-(4-hydroxy-4-(3,4-methylenedioxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

15

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole (0.51 g) and 4-hydroxy-4-(3,4-methylenedioxyphenyl)-
20 piperidine (0.42 g), an oil was obtained. This was dissolved in isopropanol and hydrochloric acid-ether solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.51 g) as white crystals, melting point 184-186°C.

25 **Example 162**

(S)-1-(2-(5-methylthio-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride 1/4 hydrate

By the reactions in the same manner as in Example 1
30 using 5-(4-hydroxybenzo(b)furan-2-yl)-2-methylthio-1,3,4-oxadiazole (0.62 g) obtained in Starting Material Synthesis Example 115, (S)-glycidyl nosylate (0.71 g) and potassium carbonate (0.41 g), (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methylthio-1,3,4-oxadiazole (0.54 g) was obtained. This was

dissolved in methanol and 4-(naphthalen-2-yl)piperidine (0.5 g) was added. The mixture was refluxed under heating for 5 hr. The reaction solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give an oil. This was dissolved in ethyl acetate and a hydrochloric acid-ether solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.34 g) as brown crystals, melting point 121-126°C.

Example 163

(S)-1-(2-(5-methoxy-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride 1/4 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methoxy-1,3,4-oxadiazole (0.18 g) obtained in Starting Material Synthesis Example 118 and 4-(naphthalen-2-yl)piperidine (0.12 g), an oil was obtained. This was dissolved in acetone and a hydrochloric acid-ether solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.11 g) as white crystals, melting point 250-251°C.

Example 164

(S)-1-(2-(5-methoxy-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrochloride 1/4 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methoxy-1,3,4-oxadiazole (0.30 g) and 4-(3,4-methylenedioxyphenyl)piperidine (0.20 g), an oil was obtained. This was dissolved in acetone and a hydrochloric acid-ether solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.17 g) as white crystals, melting point 242-246°C (decomposition).

Example 165

(S)-1-(2-(5-ethoxy-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1
5 using (S)-2-ethoxy-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole (1.0 g) and 4-(3,4-methylenedioxyphenyl)piperidine (0.65 g), crude crystals were obtained. The crystals were recrystallized from ethanol and purified to give the title compound (1.2 g) as white crystals, melting point 117-118°C.

10 **Example 166**

(S)-1-(2-(5-(1-methylethyloxy)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrobromide 1/2 hydrate

By the reactions in the same manner as in Example 1
15 using (S)-2-(1-methylethyloxy)-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole (0.50 g) and 4-(naphthalen-2-yl)piperidine (0.35 g), an oil was obtained. This was dissolved in isopropanol and 48% hydrobromic acid was added. The precipitated crystals were collected by filtration and dried to
20 give the title compound (0.52 g) as yellow crystals, melting point 123-135°C.









Example 167

(S)-1-(2-(5-(1-methylethyloxy)-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrobromide
25

By the reactions in the same manner as in Example 1
using (S)-2-(1-methylethyloxy)-5-(4-glycidyloxybenzo(b)furan-2-yl)-1,3,4-oxadiazole (0.50 g) and 4-(3,4-methylenedioxyphenyl)-piperidine (0.35 g), an oil was obtained. This was dissolved
30 in isopropanol and 48% hydrobromic acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.58 g) as yellow crystals, melting point 196-198°C.

The structural formulas of the compounds obtained in

Examples 160 to 167 are shown in the following.



Example 168

(S)-1-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1
5 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyloxazole
(2.0 g) and 4-(3,4-methylenedioxyphenyl)piperidine (1.4 g), an
oil was obtained. This was dissolved in ethanol and
hydrochloric acid was added. The precipitated crystals were
collected by filtration and dried to give the title compound
10 (1.2 g) as pale-yellow crystals, melting point 239-241°C.

¹H-NMR(DMSO-d₆) δ: 1.93-2.00 (m, 4H), 2.42 (s, 3H), 2.78 (m, 1H),
3.20 (m, 2H), 3.35-3.42 (m, 2H), 3.69-3.73 (m, 2H), 4.12-4.20 (m,
2H), 4.53 (m, 1H), 5.98 (s, 2H), 6.05 (m, 1H), 6.72 (d, J=7.8, 1H),
6.79-6.90 (m, 3H), 7.08 (s, 1H), 7.31-7.38 (m, 2H), 7.61 (s, 1H),
15 10.05 (bs, 1H)

Example 169

(S)-1-(4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyloxazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

20 By the reactions in the same manner as in Example 1
using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyloxazole
(1.4 g) and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine (1.0 g),
an oil was obtained. This was dissolved in ethanol and
hydrochloric acid was added. The precipitated crystals were
25 collected by filtration and dried to give the title compound
(0.81 g) as white crystals, melting point 230-233°C.

¹H-NMR(CDCl₃) δ: 2.01-2.08 (m, 2H), 2.45 (s, 3H), 2.59-2.74 (m, 3H),
2.90-3.02 (m, 2H), 3.18 (t, J=8.0, 2H), 3.32 (m, 2H), 3.80-3.92 (m,
2H), 4.02 (m, 1H), 4.35 (m, 1H), 4.56 (t, J=8.0, 2H), 4.79 (m, 1H),
30 5.68 (bs, 1H), 6.72 (m, 2H), 6.93 (s, 1H), 6.98 (d, J=7.8, 1H),
7.16 (s, 1H), 7.23-7.35 (m, 2H), 7.42 (s, 1H), 11.85 (bs, 1H)

Example 170

(S)-1-(2-(5-ethyloxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol dihydrochloride 1/2

hydrate

By the reactions in the same manner as in Example 1 using (S)-5-ethyl-2-(4-glycidyloxybenzo(b)furan-2-yl)oxazole (1.0 g) and 4-(3,4-methylenedioxyphenyl)piperidine (0.8 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.05 g) as white crystals, melting point 145-147°C.

Example 171

10 (S)-1-(2-(5-ethyloxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(1,4-benzodioxan-6-yl)piperidino)-2-propanol hydrochloride 5/4
hydrate

By the reactions in the same manner as in Example 1 using (S)-5-ethyl-2-(4-glycidyloxybenzo(b)furan-2-yl)oxazole (0.6 g) and 4-(1,4-benzodioxan-6-yl)piperidine (0.6 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.5 g) as white crystals, melting point 89-91°C.

20 **Example 172**

(S)-1-(2-(5-ethyloxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(naphthalen-2-yl)piperidino)-2-propanol hydrochloride 1/2
hydrate

By the reactions in the same manner as in Example 1 using (S)-5-ethyl-2-(4-glycidyloxybenzo(b)furan-2-yl)oxazole (0.7 g) and 4-(naphthalen-2-yl)piperidine (0.6 g), an oil was obtained. This was dissolved in acetone, and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.6 g) as white crystals, melting point 114-116°C.

Example 173

(S)-1-(2-(5-(1-methylethyl)oxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol dihydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-(1-methylethyl)oxazole (1.0 g) and 4-(3,4-methylenedioxyphenyl)-piperidine (0.8 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.97 g) as white crystals, melting point 139-141°C.

Example 174

(S)-1-(2-(5-(1-methylethyl)oxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-dimethylphenyl)piperidino)-2-propanol hydrochloride 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-(1-methylethyl)oxazole (0.8 g) and 4-(3,4-dimethylphenyl)-piperidine (0.6 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.59 g) as white crystals, melting point 119-121°C.


Example 175

(S)-1-(2-(5-(1-methylethyl)oxazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol hydrochloride 1/2 hydrate


By the reactions in the same manner as in Example 1 using (S)-2-(4-glycidyloxybenzo(b)furan-2-yl)-5-(1-methylethyl)oxazole (0.95 g) and 4-(3,4-dichlorophenyl)-piperidine (0.85 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.98 g) as white crystals, melting point 202-204°C.

The structural formulas of the compounds obtained in Examples 168 to 175 are shown in the following.


168




169




170




171




172




173



174



175



Example 176

5-methyl-2-(4-(3-(4-(3,4-methylenedioxyphenyl)piperidino)-propyloxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole

5-Methyl-2-(4-(3-chloropropyloxy)benzo(b)furan-2-yl)-
5 1,3,4-oxadiazole (1.0 g), 4-(3,4-methylenedioxyphenyl)-
piperidine (0.7 g), potassium carbonate (1.5 g) and potassium
iodide (1.0 g) were dissolved in a mixed solvent (50 ml) of
DMF-toluene (2:1) and the solution was stirred with refluxing
under heating at 80°C for 4 hr. The reaction mixture was
10 poured into water, extracted with ethyl acetate and dried. The
solvent was evaporated under reduced pressure and the residue
was purified by silica gel column chromatography
(chloroform/methanol) to give the title compound (0.99 g) as
white crystals, melting point 125-127°C.

15 **Example 177**

5-methyl-2-(4-(3-(4-(naphthalen-2-yl)piperidino)propyloxy)-benzo(b)furan-2-yl)-1,3,4-oxadiazole hydrochloride

By the reactions in the same manner as in Example 176
using 5-methyl-2-(4-(3-chloropropyloxy)benzo(b)furan-2-yl)-
20 1,3,4-oxadiazole (1.0 g), 4-(naphthalen-2-yl)piperidine (0.7 g),
potassium carbonate (1.5 g) and potassium iodide (1.0 g), an
oil was obtained. This was dissolved in acetone and a
hydrochloric acid-ethanol solution was added. The precipitated
crystals were collected by filtration and dried to give the
25 title compound (1.2 g) as white crystals, melting point 205-
207°C.

Example 178

2-(4-(3-((1-benzylpiperidin-4-yl)amino)propyloxy)benzo(b)furan-2-yl)-5-methyl-1,3,4-oxadiazole dihydrochloride 1/2 hydrate

30 By the reactions in the same manner as in Example 176
using 2-(4-(3-bromopropyloxy)benzo(b)-2-yl)furan-5-methyl-
1,3,4-oxadiazole (0.8 g), 4-amino-1-benzylpiperidine (0.7 g),
potassium carbonate (1.5 g) and potassium iodide (1.0 g), an
oil was obtained. This was dissolved in acetone and a

hydrochloric acid-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.51 g) as white crystals, melting point 265°C or higher.

5 **Example 179**

5-ethyl-2-(4-(3-(4-(naphthalen-2-yl)piperidino)propyloxy)-benzo(b)furan-2-yl)-1,3,4-oxadiazole hydrochloride

By the reactions in the same manner as in Example 176 using 5-ethyl-2-(4-(3-chloropropyloxy)benzo(b)furan-2-yl)-
10 1,3,4-oxadiazole (0.7 g), 4-(naphthalen-2-yl)piperidine (0.5 g), potassium carbonate (1.5 g) and potassium iodide (1.0 g), an oil was obtained. This was dissolved in acetone and a hydrochloric acid-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the
15 title compound (0.16 g) as white crystals, melting point 198-200°C.

Example 180

5-ethyl-2-(4-(3-(4-(3,4-methylenedioxyphenyl)piperidino)-propyloxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole hydrochloride

20 1/4 hydrate

By the reactions in the same manner as in Example 176 using 5-ethyl-2-(4-(3-chloropropyloxy)benzo(b)furan-2-yl)-
1,3,4-oxadiazole (1.0 g), 4-(3,4-methylenedioxyphenyl)-
piperidine (0.8 g), potassium carbonate (0.68 g) and potassium
25 iodide (0.57 g), an oil was obtained. This was dissolved in ethanol and a hydrochloric acid-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.92 g) as white crystals, melting point 192-193°C.

30 **Example 181**

5-ethyl-2-(4-(3-(4-(3,4-dichlorophenyl)piperidino)propyloxy)-benzo(b)furan-2-yl)-1,3,4-oxadiazole hydrochloride 1/4 hydrate

By the reactions in the same manner as in Example 176 using 5-ethyl-2-(4-(3-chloropropyloxy)benzo(b)furan-2-yl)-

1,3,4-oxadiazole (1.0 g), 4-(3,4-dichlorophenyl)piperidine (0.79 g), potassium carbonate (0.68 g) and potassium iodide (0.57 g), an oil was obtained. This was dissolved in ethanol and a hydrochloric acid-ethanol solution was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.1 g) as white crystals, melting point 225-227°C.

Example 182

2-(4-(3-(4-(1,4-benzodioxan-6-yl)piperidino)propyloxy)-benzo(b)furan-2-yl)-5-(1-methylethyl)-1,3,4-oxadiazole hydrochloride

By the reactions in the same manner as in Example 176 using 5-(1-methylethyl)-2-(4-(3-chloropropoxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole (0.9 g), 4-(1,4-benzodioxan-6-yl)piperidine (0.7 g), potassium carbonate (1.5 g) and potassium iodide (1.0 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.68 g) as white crystals, melting point 203-205°C.

Example 183


2-(4-(3-(4-(3,4-dimethylphenyl)piperidino)propyloxy)-benzo(b)furan-2-yl)-5-(1-methylethyl)-1,3,4-oxadiazole hydrochloride

By the reactions in the same manner as in Example 176 using 5-(1-methylethyl)-2-(4-(3-chloropropoxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole (1.0 g), 4-(3,4-dimethylphenyl)-piperidine (0.8 g), potassium carbonate (1.5 g) and potassium iodide (1.0 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.9 g) as white crystals, melting point 217-219°C.


The structural formulas of the compounds obtained in

Examples 176 to 183 are shown in the following.


176




177




178




179




180




181



182



183



Example 184

2-(4-(3-(4-(3,4-dichlorophenyl)piperidino)propyloxy)-
benzo(b)furan-2-yl)-5-(1-methylethyl)-1,3,4-oxadiazole
hydrochloride

5 By the reactions in the same manner as in Example 176
using 5-(1-methylethyl)-2-(4-(3-chloropropoxy)benzo(b)furan-
2-yl)-1,3,4-oxadiazole (1.0 g), 4-(3,4-dichlorophenyl)-
piperidine (0.8 g), potassium carbonate (1.5 g) and potassium
iodide (1.0 g), an oil was obtained. This was dissolved in
10 acetone and hydrochloric acid was added. The precipitated
crystals were collected by filtration and dried to give the
title compound (0.9 g) as white crystals, melting point 247-
249°C.

Example 185

15 2-(4-(3-(4-(naphthalen-2-yl)piperidino)propyloxy)benzo(b)furan-
2-yl)-5-(1-methylethyl)-1,3,4-oxadiazole hydrochloride

By the reactions in the same manner as in Example 176
using 5-(1-methylethyl)-2-(4-(3-chloropropoxy)benzo(b)furan-
2-yl)-1,3,4-oxadiazole (1.0 g), 4-(naphthalen-2-yl)piperidine
20 (0.9 g), potassium carbonate (1.5 g) and potassium iodide (1.0
g), an oil was obtained. This was dissolved in acetone and
hydrochloric acid was added. The precipitated crystals were
collected by filtration and dried to give the title compound
(1.15 g) as white crystals, melting point 215-217°C.

25 **Example 186**

2-(4-(3-(4-(3,4-methylenedioxyphenyl)piperidino)-
propyloxy)benzo(b)furan-2-yl)-5-(1-methylethyl)-1,3,4-
oxadiazole hydrochloride

By the reactions in the same manner as in Example 176
30 using 5-(1-methylethyl)-2-(4-(3-chloropropoxy)benzo(b)furan-
2-yl)-1,3,4-oxadiazole (1.0 g), 4-(3,4-methylenedioxyphenyl)-
piperidine (1.0 g), potassium carbonate (1.5 g) and potassium
iodide (1.0 g), an oil was obtained. This was dissolved in
acetone and hydrochloric acid was added. The precipitated

crystals were collected by filtration and dried to give the title compound (1.07 g) as white crystals, melting point 193-195°C.

Example 187

5 2-(4-(3-(4-(3,4-dimethylphenyl)piperidino)propyloxy)-
benzo(b)furan-2-yl)-5-(1,1-dimethylethyl)-1,3,4-oxadiazole
hydrochloride

By the reactions in the same manner as in Example 176 using 5-(1,1-dimethylethyl)-2-(4-(3-chloropropyloxy)-
10 benzo(b)furan-2-yl)-1,3,4-oxadiazole (0.8 g), 4-(3,4-dimethylphenyl)piperidine (0.7 g), potassium carbonate (1.5 g) and potassium iodide (1.0 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to
15 give the title compound (0.68 g) as white crystals, melting point 226-228°C.

Example 188

2-(4-(3-(4-(3,4-dichlorophenyl)piperidino)propyloxy)-
benzo(b)furan-2-yl)-5-(1,1-dimethylethyl)-1,3,4-oxadiazole
20 hydrochloride

By the reactions in the same manner as in Example 176 using 5-(1,1-dimethylethyl)-2-(4-(3-chloropropyloxy)-
benzo(b)furan-2-yl)-1,3,4-oxadiazole (1.0 g), 4-(3,4-dichlorophenyl)piperidine (0.8 g), potassium carbonate (1.5 g)
25 and potassium iodide (1.0 g), an oil was obtained. This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.75 g) as white crystals, melting point 249-251°C.

30 **Example 189**

2-(4-(3-(4-(3,4-methylenedioxyphenyl)piperidino)propyloxy)-
benzo(b)furan-2-yl)-5-(1,1-dimethylethyl)-1,3,4-oxadiazole
hydrochloride

By the reactions in the same manner as in Example 176

using 5-(1,1-dimethylethyl)-2-(4-(3-chloropropoxy)-benzo(b)furan-2-yl)-1,3,4-oxadiazole (1.0 g), 4-(3,4-methylenedioxyphenyl)piperidine (0.8 g), potassium carbonate (1.5 g) and potassium iodide (1.0 g), an oil was obtained.

5 This was dissolved in acetone and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (1.1 g) as white crystals, melting point 211-213°C.

Example 190

10 5-(2-methylpropyl)-2-(4-(3-(4-(3,4-methylenedioxyphenyl)-piperidin-1-yl)propoxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole

By the reactions in the same manner as in Example 176 using 5-(2-methylpropyl)-2-(4-(3-chloropropoxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole (0.5 g), 4-(3,4-methylenedioxyphenyl)-
15 piperidine (0.37 g), potassium carbonate (0.2 g) and potassium iodide (0.25 g), the title compound (0.32 g) was obtained as brown crystals, melting point 103-105°C.

Example 191


2-(4-(3-(4-(3,4-dimethylphenyl)piperidin-1-yl)propoxy)-benzo(b)furan-2-yl)-5-(2-methylpropyl)-1,3,4-oxadiazole
20 hydrochloride

By the reactions in the same manner as in Example 176 using 5-(2-methylpropyl)-2-(4-(3-chloropropoxy)benzo(b)furan-2-yl)-1,3,4-oxadiazole (0.67 g), 4-(3,4-dimethylphenyl)-
25 piperidine (0.4 g), potassium carbonate (0.33 g) and potassium iodide (0.33 g), an oil was obtained. This was dissolved in ethanol and hydrochloric acid was added. The precipitated crystals were collected by filtration and dried to give the title compound (0.28 g) as white crystals, melting point 200-
30 203°C.


The structural formulas of the compounds obtained in Examples 184 to 191 are shown in the following.

Publ. 630660


184




185




186




187




188




189



190



191



Example 192

2-(4-(3-(4-(3,4-dichlorophenyl)piperidin-1-yl)propyloxy)-
benzo(b)furan-2-yl)-5-(2-methylpropyl)-1,3,4-oxadiazole
hydrochloride

5 By the reactions in the same manner as in Example 176
using 5-(2-methylpropyl)-2-(4-(3-chloropropoxy)benzo(b)furan-
2-yl)-1,3,4-oxadiazole (0.43 g), 4-(3,4-dichlorophenyl)-
piperidine (0.3 g), potassium carbonate (0.2 g) and potassium
iodide (0.25 g), an oil was obtained. This was dissolved in
10 ethanol and hydrochloric acid was added. The precipitated
crystals were collected by filtration and dried to give the
title compound (0.3 g) as white crystals, melting point 212-
215°C.

Example 193

15 2-(4-(3-(4-(naphthalen-2-yl)piperidin-1-yl)propyloxy)-
benzo(b)furan-2-yl)-5-(2-methylpropyl)-1,3,4-oxadiazole
hydrochloride

By the reactions in the same manner as in Example 176
using 5-(2-methylpropyl)-2-(4-(3-chloropropoxy)benzo(b)furan-
20 2-yl)-1,3,4-oxadiazole (0.67 g), 4-(naphthalen-2-yl)piperidine
(0.4 g), potassium carbonate (0.33 g) and potassium iodide
(0.33 g), an oil was obtained. This was dissolved in ethanol
and hydrochloric acid was added. The precipitated crystals
were collected by filtration and dried to give the title
25 compound (0.48 g) as pale-yellow crystals, melting point 183-
185°C.

Example 194

(S)-3-(4-(3,4-dichlorophenyl)piperidino)-1-(2-(5-methyl-1,3,4-
oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanone

30 To a solution of (S)-3-(4-(3,4-dichlorophenyl)-
piperidino)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-
4-yloxy)-2-propanol (1.0 g) in a mixed solvent (20 ml) of DMSO-
toluene (1:1) were added WSC (1.9 g) and trifluoroacetic acid
(0.60 ml), and the mixture was further stirred under ice-

cooling for 1 hr and then at room temperature for 1 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium hydrogencarbonate solution and dried over
5 anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the obtained crude crystals were washed with ethanol and ethyl acetate to give the title compound (0.28 g) as white crystals, melting point 128°C (decomposition).

¹H-NMR(CDCl₃) δ: 1.83 (m, 4H), 2.05 (m, 2H), 2.31 (m, 1H), 2.67 (s, 3H), 3.05 (m, 2H), 3.54 (s, 2H), 4.90 (s, 2H), 6.62 (d, J=7.8, 1H),
10 7.07 (m, 1H), 7.26-7.38 (m, 4H), 7.64 (s, 1H)

The following compounds can be also synthesized by similar reactions.

Example 195

15 (S)-1-(2-(2-(5-methyl-1,3,4-oxadiazol-2-yl)vinyl)phenyloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-5-(2'-glycidyloxystyryl)-2-methyl-1,3,4-oxadiazole and 4-(2,3-dihydrobenzo(b)furan-6-yl)piperidine, the title
20 compound can be obtained.

Example 196

(S)-1-(2-(2-(3-methyl-1,2,4-oxadiazol-5-yl)vinyl)phenyloxy)-3-(4-(3,4-dichlorophenyl)piperidino)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-5-(2'-glycidyloxystyryl)-3-methyl-1,2,4-oxadiazole
25 and 4-(3,4-dichlorophenyl)piperidine, the title compound can be obtained.

Example 197

30 (S)-4-(2-hydroxy-3-(4-(chroman-6-yl)piperidino)propyloxy)-N,N-dimethylbenzo(b)thiophene-2-carboxamide

By the reactions in the same manner as in Example 1 using (S)-4-(glycidyloxy)benzo(b)thiophen-2-yl-N,N-dimethylcarboxamide and 4-(chroman-6-yl)piperidine, the title compound can be obtained.

Example 198

(S)-1-(4-(2,3-dihydrobenzo(b) furan-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol









By the reactions in the same manner as in Example 1
5 using (S)-5-(4-glycidyloxybenzo(b) furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(2,3-dihydrobenzo(b) furan-6-yl)piperidine, the title compound was obtained as white crystals, melting point 133-135°C.

Example 199

10 (S)-1-(4-(2,3-dihydrobenzo(b) furan-6-yl)piperidino)-3-(2-(5-methyloxazol-2-yl)benzo(b) thiophen-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
using (S)-2-(4-glycidyloxybenzo(b) furan-2-yl)-5-methyloxazole
and 4-(2,3-dihydrobenzo(b) furan-6-yl)piperidine, the title
15 compound can be obtained.

The structural formulas of the compounds obtained in Examples 192 to 199 are shown in the following.



Example 200

(S)-1-(4-(4-methoxy-3-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
5 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(4-methoxy-3-methylphenyl)piperidine, the title compound was obtained as white crystals, melting point 128-130°C.

Example 201

10 (S)-1-(4-(2,3-dihydrobenzo(b)thiophen-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(2,3-dihydrobenzo(b)thiophen-5-yl)piperidine,
15 the title compound can be obtained.

Example 202

(S)-1-(4-(indan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

By the reactions in the same manner as in Example 1
20 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(indan-5-yl)piperidine, the title compound was obtained as white crystals, melting point 202-205°C.

Example 203

25 (S)-1-(4-(inden-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(inden-5-yl)piperidine, the title compound can be obtained.

30 **Example 204**

(S)-1-(4-(1-methylindolin-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-

oxadiazole and 4-(1-methylindolin-5-yl)piperidine, the title compound can be obtained.

Example 205

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-
5 3-(1,3-dihydrobenzo(c)furan-1-spiro-4'-piperidin-1'-yl)-2-
propanol 1/4 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyoxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 1,3-dihydrobenzo(c)furan-1-spiro-4'-piperidine,
10 the title compound was obtained as white crystals, melting point 198-199°C.

Example 206

(S)-1-(2-(5-ethyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-
3-(1,3-dihydrobenzo(c)furan-1-spiro-4'-piperidin-1'-yl)-2-
15 propanol

By the reactions in the same manner as in Example 1 using (S)-2-ethyl-5-(4-glycidyoxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 1,3-dihydrobenzo(c)furan-1-spiro-4'-piperidine, the title compound can be obtained.


20 **Example 207**

(S)-1-(2-(5-(1-methylethyl)-1,3,4-oxadiazol-2-yl)benzo(b)furan-
4-yloxy)-3-(1,3-dihydrobenzo(c)furan-1-spiro-4'-piperidin-1'-
yl)-2-propanol


By the reactions in the same manner as in Example 1
25 using (S)-2-(1-methylethyl)-5-(4-glycidyoxybenzo(b)furan-2-yl)-1,3,4-oxadiazole and 1,3-dihydrobenzo(c)furan-1-spiro-4'-piperidine, the title compound can be obtained.

The structural formulas of the compounds obtained in Examples 200 to 207 are shown in the following.


200




201




202




203




204




205



206



207



Example 208

(S)-1-(4-benzylpiperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-benzylpiperidine, the title compound can be obtained.

5 **Example 209**

(S)-1-(4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)thiophen-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
10 using (S)-5-(4-glycidyloxybenzo(b)thiophen-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine, the title compound can be obtained.

Example 210

(S)-1-(4-(benzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride
15 1/2 hydrate

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(benzo(b)furan-5-yl)piperidine, the title
20 compound could be obtained, melting point 182-184°C.

Example 211

(S)-1-(4-(2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,2,4-oxadiazol-3-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
25 using (S)-3-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole and 4-(2,3-dihydrobenzo(b)furan-5-yl)piperidine, the title compound can be obtained.

Example 212

(S)-1-(3-(3,4-dichlorophenyl)propylamino)-3-(2-(5-methyl-1,2,4-oxadiazol-3-yl)benzo(b)furan-4-yloxy)-2-propanol
30

By the reactions in the same manner as in Example 1 using (S)-3-(4-glycidyloxybenzo(b)furan-2-yl)-5-methyl-1,2,4-oxadiazole and 3-(3,4-dichlorophenyl)propylamine, the title compound can be obtained.

Example 213

(S)-1-(4-(naphthalen-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-thiadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
5 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-thiadiazole and 4-(naphthalen-2-yl)piperidine, the title compound can be obtained.

Example 214

(S)-1-(4-(2,3-dihydrobenzo(b)thiophen-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1
using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(2,3-dihydrobenzo(b)thiophen-6-yl)piperidine, the title compound can be obtained.


15 **Example 215**

(S)-1-(4-(chroman-7-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride 1/2 hydrate


By the reactions in the same manner as in Example 1
20 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(chroman-7-yl)piperidine, the title compound was obtained as white crystals, melting point 210-212°C.

The structural formulas of the compounds obtained in Examples 208 to 215 are shown in the following.


208




209




210




211




212




213



214



215

**Example 216**

(S)-1-(4-(benzo(b)thiophen-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(benzo(b)thiophen-5-yl)piperidine, the title compound can be obtained.

5 **Example 217**

(S)-1-(4-(2H-chromen-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

By the reactions in the same manner as in Example 1 using (S)-5-(4-glycidyloxybenzo(b)furan-2-yl)-2-methyl-1,3,4-oxadiazole and 4-(2H-chromen-6-yl)piperidine, the title compound can be obtained.

Example 218

(S)-1-(4-(3,4-dihydro-2H-benzo(b)thiain-7-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 219

(S)-1-(4-(3-chloro-4-methoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol
hydrochloride
20 white crystals, melting point 218-220°C

Example 220

(S)-1-(4-(4-chloro-3-methoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 221

25 (S)-1-(4-(benzo(d)isoxazol-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 222

(S)-1-(4-(4-chloro-3-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol
30 hydrochloride
melting point 184-186°C

Example 223

(S)-1-(4-(3-chloro-4-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol


hydrochloride

white crystals, melting point 188-190°C


The structural formulas of the compounds obtained in Examples 216 to 223 are shown in the following.

5


216




217




218




219




220




221



222



223



Example 224

(S)-1-(4-(benzo(b) furan-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

Example 225

5 (S)-1-(4-(2H-chromen-7-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

Example 226

(S)-1-(4-(1H-indol-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

10 **Example 227**

(S)-1-(4-(1-methylindol-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

Example 228

15 (S)-1-(4-(1,3-dihydrobenzo(c) furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

Example 229

(S)-1-(4-(isochroman-7-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

Example 230

20 (S)-1-(4-(isochroman-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

Example 231

(S)-1-(4-(2,3-dihydrobenzo(b) furan-4-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol

25 **Example 232**

(S)-1-(4-(2,2-dimethyl-2,3-dihydrobenzo(b) furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol 1/2 terephthalate

melting point 159-162°C


30 **Example 233**

(S)-1-(4-(3,4-dihydro-2H-benzo(b) thiin-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b) furan-4-yloxy)-2-propanol hydrochloride


white crystals, melting point 223-225°C

The structural formulas of the compounds obtained in Examples 224 to 233 are shown in the following.


224




225




226




227




228




229




230




231



232



233



Example 234

(S)-1-(4-(4-chloro-3-trifluoromethylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

5 white crystals, melting point 222-224°C

Example 235

(S)-1-(4-(4-chloro-3-fluorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

10 white crystals, melting point 139-141°C

Example 236

(S)-1-(4-(benzo(b)furan-6-yl)-5,6-dihydro-2H-pyridin-1-yl)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol hydrochloride

15 white crystals, melting point 135-138°C

Example 237

(S)-1-(2-(5-isopropenyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-3-(4-(3,4-methylenedioxyphenyl)piperidino)-2-propanol hydrochloride

20 white crystals, melting point 128-130°C


Example 238

(S)-1-(4-(benzo(b)furan-5-yl)-5,6-dihydro-2H-pyridin-1-yl)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol


25 white crystals, melting point 168-170°C

The structural formulas of the compounds obtained in Examples 234 to 238 are shown in the following.


234




235




236



237



238

**Example 301**

(1) 3,5-dihydro-5-(2'-methoxybenzylidene)-2-methylimidazol-4-

5 one

4-(2'-Methoxybenzylidene)-2-methyl-4H-oxazol-5-one (2.4 g) was dissolved in ethanol (50 ml) and aqueous ammonia (20 ml) and potassium carbonate (3 g) were added. The mixture was heated at 80°C for 10 hr. After cooling, the solvent was evaporated under reduced pressure and water was added to the residue. The mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The precipitated yellow

crystals were collected by filtration to give the title compound (2 g).

¹H-NMR(DMSO-d₆)δ:2.23(s, 3H), 3.87(s, 3H), 7.00(t, J=7.8, 1H), 7.02(d, J=8.3, 1H), 7.18(s, 1H), 7.35(t, J=7.8, 1H), 8.71(d, J=8.3, 1H)

(2) (S)-3,5-dihydro-5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidin-1-yl)propyloxy)benzylidene)-2-methylimidazol-4-one

3,5-Dihydro-5-(2'-methoxybenzylidene)-2-methylimidazol-4-one (1.28 g) was dissolved in dichloromethane (20 ml) and boron tribromide (4.5 g) was added dropwise while stirring at -40°C. Thereafter, the mixture was stirred for 2 hr under ice-cooling and the reaction mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 3,5-dihydro-5-(2'-hydroxybenzylidene)-2-methylimidazol-4-one (1 g) as red crystals. The crystals and (S)-glycidyl nosylate (1.3 g) were dissolved in dimethylformamide (20 ml) and potassium carbonate (1.38 g) was added. The mixture was heated at 50°C for 2 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (1.12 g). This oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (15 ml) and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.23 g) as an oil.

¹H-NMR(DMSO-d₆)δ:1.91-2.20(m, 4H), 2.45(s, 3H), 2.90-3.05(m, 4H), 3.16(m, 1H), 3.54(m, 2H), 4.17(m, 2H), 4.31(m, 1H), 5.01(bs, 1H), 6.87(t, J=7.8, 1H), 6.91(d, J=8.3, 1H), 7.30(s, 1H), 7.42-7.54(m, 5H), 7.72(s, 1H), 7.84(m, 2H), 8.78(d, J=8.3, 1H)

Example 302

(1) 3,5-dihydro-2,3-dimethyl-5-(2'-methoxybenzylidene)imidazol-4-one

4-(2'-Methoxybenzylidene)-2-methyl-4H-oxazol-5-one (5 g)
5 was dissolved in ethanol (100 ml) and aqueous methylamine solution (20 ml) and potassium carbonate (7 g) were added. The mixture was heated at 80°C for 7 hr. After cooling, the solvent was evaporated under reduced pressure and the precipitated yellow crystals were collected by filtration to
10 give the title compound (2 g).

¹H-NMR(DMSO-d₆)δ:2.35(s, 3H), 3.10(s, 3H), 3.89(s, 3H), 7.00(t, J=7.8, 1H), 7.05(d, J=8.3, 1H), 7.33(s, 1H), 7.39(t, J=7.8, 1H), 8.73(d, J=8.3, 1H)

(2) (S)-3,5-dihydro-2,3-dimethyl-5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidin-1-yl)propyloxy)benzylidene)imidazol-4-one

3,5-Dihydro-2,3-dimethyl-5-(2'-methoxybenzylidene)-imidazol-4-one (2 g) was dissolved in dichloromethane (30 ml) and boron tribromide (2 ml) was added dropwise while stirring
20 the mixture at -40°C. Thereafter, the reaction mixture was stirred under ice-cooling for 1 hr, and the reaction mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 3,5-dihydro-2,3-dimethyl-5-(2'-hydroxybenzylidene)imidazol-4-one (1.72 g) as
25 red crystals. The crystals and (S)-glycidyl nosylate (2 g) were dissolved in dimethylformamide (20 ml) and potassium carbonate (2.2 g) was added. The mixture was heated at 50°C for 3 hr, and the reaction mixture was poured into ice water
30 and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (2.27 g). This oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol

(20 ml) and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound
5 (1.25 g) as an oil.

¹H-NMR(DMSO-d₆) δ: 2.03-2.20(m, 4H), 2.37(s, 3H), 2.99(m, 1H), 3.11(s, 3H), 3.10-3.23(m, 3H), 3.31(m, 1H), 3.63(m, 2H), 4.13(m, 2H), 4.34(m, 1H), 5.05(bs, 1H), 7.06(t, J=7.8, 1H), 7.12(d, J=8.3, 1H), 7.38-7.52(m, 6H), 7.74(s, 1H), 7.88(m, 2H), 8.78(d, J=8.3, 1H)
10

Example 303

(1) 3,5-dihydro-5-(2'-methoxybenzylidene)-2-methyl-3-phenylimidazol-4-one

4-(2'-Methoxybenzylidene)-2-methyl-4H-oxazol-5-one (5 g)
15 was dissolved in acetic acid (100 ml), and aniline (2.33 g) and sodium acetate (1.8 g) were added. The mixture was heated at 100°C for 5 hr. The reaction mixture was poured into ice-water, neutralized with potassium carbonate and extracted with ethyl acetate. The organic layer was washed with water, dried over
20 anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.85 g) as yellow crystals.

¹H-NMR(CDCl₃) δ: 2.27(s, 3H), 3.90(s, 3H), 6.92(d, J=8.3, 1H),
25 7.06(t, J=7.8, 1H), 7.26(m, 2H), 7.35-7.53(m, 4H), 7.78(s, 1H), 8.78(d, J=8.3, 1H)

(2) (S)-3,5-dihydro-5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidin-1-yl)propyloxy)benzylidene)-2-methyl-3-phenylimidazol-4-one

30 3,5-Dihydro-5-(2'-methoxybenzylidene)-2-methyl-3-phenylimidazol-4-one (1.85 g) was dissolved in dichloromethane (30 ml) and boron tribromide (2 ml) was added dropwise while stirring the mixture at -40°C. Thereafter, the mixture was stirred under ice-cooling for 1 hr, poured into ice water and

extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 3,5-dihydro-5-(2'-hydroxybenzylidene)-2-methyl-3-phenylimidazol-4-one (1.76 g) as red crystals. The crystals and (S)-glycidyl nosylate (1.63 g) were dissolved in dimethylformamide (20 ml) and potassium carbonate (1.74 g) was added. The mixture was heated at 50°C for 2 hr. The reaction mixture was poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (2.13 g). This oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (20 ml) and refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.45 g) as an oil.

¹H-NMR(DMSO-d₆) δ: 2.09-2.20 (m, 4H), 2.23 (s, 3H), 3.00 (m, 1H), 3.15-3.29 (m, 3H), 3.37 (m, 1H), 3.66 (m, 2H), 4.14 (m, 2H), 4.39 (m, 1H), 5.02 (bs, 1H), 7.10 (t, J=7.8, 1H), 7.14 (d, J=8.3, 1H), 7.38-7.56 (m, 10H), 7.73 (s, 1H), 7.88 (m, 3H), 8.84 (d, J=8.3, 1H)

Example 304

(1) 3-ethyl-3,5-dihydro-5-(2'-methoxybenzylidene)-2-methylimidazol-4-one

4-(2'-Methoxybenzylidene)-2-methyl-4H-oxazol-5-one (5 g) was dissolved in ethanol (100 ml) and an aqueous ethylamine solution (15 ml) and potassium carbonate (7 g) were added thereto. The mixture was heated at 80°C for 5 hr. After cooling, the solvent was evaporated under reduced pressure and the precipitated yellow crystals were collected by filtration to give the title compound (1.2 g).

¹H-NMR(CDCl₃) δ: 1.25 (t, J=7.3, 3H), 2.39 (s, 3H), 3.67 (q, J=7.3, 2H), 3.88 (s, 3H), 6.89 (d, J=8.3, 1H), 7.02 (t, J=7.8, 1H), 7.34 (t, J=7.8, 1H), 7.67 (s, 1H), 8.72 (d, J=8.3, 1H)

(2) (S)-3-ethyl-3,5-dihydro-5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidin-1-yl)propyloxy)benzylidene)-2-methylimidazol-4-one

3-Ethyl-3,5-dihydro-5-(2'-methoxybenzylidene)-2-methylimidazol-4-one (1.2 g) was dissolved in dichloromethane (20 ml) and boron tribromide (1.5 ml) was added dropwise while stirring the mixture at -40°C. Thereafter, the mixture was stirred under ice-cooling for 1 hr, poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 3-ethyl-3,5-dihydro-5-(2'-hydroxybenzylidene)-2-methylimidazol-4-one (1 g) as red crystals. The crystals and (S)-glycidyl nosylate (1.3 g) were dissolved in dimethylformamide (20 ml) and potassium carbonate (1.38 g) was added. The mixture was heated at 50°C for 2 hr, poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (1.02 g). This oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (10 ml) and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.52 g) as an oil.

¹H-NMR (DMSO-d₆) δ: 1.27 (t, J=7.3, 3H), 2.03-2.20 (m, 4H), 2.42 (s, 3H), 2.99 (m, 1H), 3.68 (q, J=7.3, 2H), 3.10-3.23 (m, 3H), 3.31 (m, 1H), 3.63 (m, 2H), 4.13 (m, 2H), 4.34 (m, 1H), 5.05 (bs, 1H), 7.06 (t, J=7.8, 1H), 7.12 (d, J=8.3, 1H), 7.38-7.52 (m, 6H), 7.74 (s, 1H), 7.88 (m, 2H), 8.78 (d, J=8.3, 1H)

Example 305

(1) 3-benzyl-3,5-dihydro-5-(2'-methoxybenzylidene)-2-methylimidazol-4-one

4-(2'-Methoxybenzylidene)-2-methyl-4H-oxazol-5-one (5 g)

was dissolved in acetic acid (50 ml) and benzylamine (2.68 g) and sodium acetate (2 g) were added. The mixture was heated at 100°C for 2 hr, poured into ice-water, neutralized with potassium carbonate and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.38 g) as yellow crystals.

¹H-NMR(CDCl₃) δ: 2.25(s, 3H), 3.89(s, 3H), 4.83(s, 2H), 6.90(d, J=8.3, 1H), 7.02(t, J=7.8, 1H), 7.22-7.35(m, 6H), 7.77(s, 1H), 8.73(d, J=8.3, 1H)

(2) (S)-3-benzyl-3,5-dihydro-5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidin-1-yl)propyloxy)benzylidene)-2-methylimidazol-4-

one

3-Benzyl-3,5-dihydro-5-(2'-methoxybenzylidene)-2-methylimidazol-4-one (1.38 g) was dissolved in dichloromethane (20 ml) and boron tribromide (1.3 ml) was added dropwise while stirring the mixture at -40°C. Thereafter, the mixture was stirred under ice-cooling for 1 hr, poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 3-benzyl-3,5-dihydro-5-(2'-hydroxybenzylidene)-2-methylimidazol-4-one (1.14 g) as red crystals. The crystals and (S)-glycidyl nosylate (1 g) were dissolved in dimethylformamide (20 ml) and potassium carbonate (1.1 g) was added. The mixture was heated at 50°C for 2 hr, poured into ice water and extracted with ethyl acetate. The organic layer was washed with saturated aqueous ammonium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (1.05 g). This oily product and 4-(naphthalen-2-yl)piperidine were dissolved in methanol (10 ml) and refluxed under heating for 2 hr. After cooling, the solvent was evaporated under

reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.25 g) as an oil.

¹H-NMR (DMSO-d₆) δ: 2.09-2.20 (m, 4H), 2.23 (s, 3H), 3.00 (m, 1H),
5 3.15-3.29 (m, 3H), 3.37 (m, 1H), 3.66 (m, 2H), 4.14 (m, 2H), 4.39 (m, 1H), 4.89 (s, 2H), 5.02 (bs, 1H), 7.12 (t, J=7.8, 1H), 7.15 (d, J=8.3, 1H), 7.40-7.58 (m, 10H), 7.75 (s, 1H), 7.85 (m, 3H), 8.85 (d, J=8.3, 1H)

Example 306

10 (S)-5-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-1,3-dimethylimidazolidine-2,4-dione

2-(Methoxymethyloxy)benzaldehyde (10 g) and hydantoin (8.2 g) were dissolved in piperidine (20 ml). Benzylamine (2.68 g) and sodium acetate (2 g) were added thereto and the
15 mixture was heated at 130°C for 5 hr. The reaction mixture was poured into ice water, neutrized with hydrochloric acid and extracted with ethyl acetate to give a reaction concentrate (10 g). Thereto were added dimethylformamide (100 ml) and potassium carbonate (13.5 g), and then methyl iodide (3.7 ml)
20 was added. The mixture was stirred while refluxing under heating at 40°C for 2 hr. The reaction mixture was concentrated under reduced pressure and water was added. The mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under
25 reduced pressure. The residue was purified by silica gel column chromatography (hexane/acetone) to give an oily product (6.2 g). Acetic acid (50 ml) and water (50 ml) were poured into the oily product and the mixture was refluxed under heating for 1 hr. After completion of the reaction, the
30 reaction mixture was poured into water, and extracted with chloroform/methanol. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily product (6.0 g). Thereto were added dimethylformamide (60 ml) and potassium carbonate (10.7 g), and

(S)-glycidyl nosylate (6.7 g) was further added. The mixture was stirred with heating at 50°C for 2 hr. The reaction mixture was concentrated under reduced pressure and water was added. The mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give an oily product (2.4 g). This oil (1.0 g) and 4-(naphthalen-2-yl)piperidine (0.9 g) were dissolved in methanol (30 ml) and the mixture was refluxed under heating for 3 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give an oily product.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.85-1.95 (m, 4H), 2.16 (m, 1H), 2.47-2.70 (m, 3H), 2.94 (s, 3H), 2.99 (bs, 1H), 3.13 (s, 3H), 3.20 (bs, 1H), 4.01-4.13 (m, 3H), 6.63-6.99 (m, 2H), 7.15 (d, 1H, $J=7.4$), 7.28-7.44 (m, 6H), 7.64 (s, 1H), 7.77 (m, 2H)

Example 307

(1) α -(2'-hydroxybenzylidene)- γ -butyrolactone

Salicylaldehyde (293 g) and γ -butyrolactone (413 g) were dissolved in toluene (2.4 L) and the solution was cooled to not more than 3°C in an ice-salt bath. Thereto was added sodium methoxide (324 g) over 20 min. The temperature of the reaction mixture rose to 24°C. After stirring at room temperature for 3 hr, the mixture was stirred for 45 min under heating at 60-65°C. The reaction mixture was cooled again in an ice-bath and 10% sulfuric acid (2.51 ml) was added dropwise. The obtained white suspension was filtrated, washed with water and dried to give the title compound (324 g), melting point 162-164°C.

(2) (S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone

To α -(2'-hydroxybenzylidene)- γ -butyrolactone (60 g) were added dimethylformamide (600 ml) and potassium carbonate (87 g) and (S)-glycidyl nosylate (82 g) was added. The reaction mixture was stirred for 2 days at room temperature. The

reaction mixture was concentrated under reduced pressure and water was added. The mixture was extracted with ethyl acetate.

The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The obtained crystals
5 were recrystallized from ethyl acetate to give the title compound (55 g), melting point 68-70°C.

(3) (S)- α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)benzylidene)- γ -butyrolactone

(S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -
10 butyrolactone (50 g) and 4-(naphthalen-2-yl)piperidine (43 g) were dissolved in ethanol (500 ml) and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the obtained crystals were collected by filtration. The crystals were recrystallized
15 once from acetonitrile and once from a mixed solvent of ethyl acetate and ethanol to give the title compound (50 g), melting point 138-140°C.

Example 308

(1) (R)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -
20 butyrolactone

To α -(2'-hydroxybenzylidene)- γ -butyrolactone (11 g) were added dimethylformamide (100 ml) and potassium carbonate (20 g) and (R)-glycidyl nosylate (15 g) was added. The mixture was stirred at 50°C for 3 hr. Water was added and the mixture was
25 extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (10 g) as crystals, melting point 135-137°C.

(2) (R)- α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-
30 propyloxy)benzylidene)- γ -butyrolactone 1/4 hydrate

By the reactions in the same manner as in Example 307 using (R)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (2.7 g) and 4-(naphthalen-2-yl)piperidine (2.5 g) in methanol (50 ml), the title compound (2.4 g) was obtained,

melting point 136-138°C.

Example 309

α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)-benzylidene)- γ -butyrolactone

5 Dimethylformamide (30 ml) and potassium carbonate (4.4 g) were added to α -(2'-hydroxybenzylidene)- γ -butyrolactone (3 g), and a racemic compound of glycidyl nosylate (3.4 g) was added. The mixture was stirred at 50°C for 3 hr. Water was added and the mixture was extracted with ethyl acetate. The
10 organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give oily α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (3.2 g). This compound (0.4 g) was reacted in methanol with 4-(naphthalen-2-yl)piperidine (0.5 g) in the same manner as in Example 307 to
15 give the title compound, melting point 127-128°C.


Example 310

(S)- α -(2'-(3-(4-(3-fluoro-4-methylphenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone 1/10 hydrate


By the reactions in the same manner as in Example 307
20 using (S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (1.0 g) and 4-(3-fluoro-4-methylphenyl)piperidine (0.8 g) in methanol (50 ml), the title compound (1.24 g) was obtained, melting point 131-133°C.

The structural formulas of the compounds obtained in
25 Examples 301 to 310 are shown in the following.


301-1




301-2




302-1




302-2




303-1




303-2




304-1




304-2




305-1




305-2




306




307-1




307-2




307-3




308-1




308-2



309



310



Example 311

(S)- α -(2'-(3-(4-(3,4-dimethylphenyl)piperidino)-2-
 5 hydroxypropyloxy)benzylidene)- γ -butyrolactone 1/5 hydrate

By the reactions in the same manner as in Example 307

using (S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (1.0 g) and 4-(3,4-dimethylphenyl)piperidine (0.8 g) in methanol (50 ml), the title compound (0.61 g) was obtained, melting point 125-126°C.

5 **Example 312**

(S)- α -(2'-(3-(4-(4-chloro-3-fluorophenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone 1/2 hydrate

By the reaction in same manner as in Example 307 using (S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (1.0 g) and 4-(4-chloro-3-fluorophenyl)piperidine (0.8 g) in methanol (50 ml), the title compound (0.58 g), melting point 114-115°C.

Example 313

15 (S)- α -(2'-(3-(4-(4-chloro-3-trifluoromethylphenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone hydrochloride monohydrate

The reaction was performed in same manner as in Example 307 using (S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (1.0 g) and 4-(4-chloro-3-trifluoromethylphenyl)-piperidine (0.8 g) in methanol (50 ml), and the obtained oil was treated with methanol/hydrochloric acid to give the title compound (0.45 g), melting point 172-174°C.

Example 314

25 (S)- α -(2'-(2-hydroxy-3-(4-(naphthalen-1-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone p-toluenesulfonate

The reaction was performed in same manner as in Example 307 using (S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (5.8 g) and 4-(naphthalen-1-yl)piperidine (5.0 g) in methanol (100 ml) and the obtained oil was treated with ethyl acetate/p-toluenesulfonic acid to give the title compound (7.8 g), melting point 119-121°C.

Example 315

(S)- α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-

propyloxy)benzylidene)- δ -valerolactone 1/5 hydrate

Salicylaldehyde (15.2 g) and δ -valerolactone (25 g) were dissolved in toluene (150 ml). By the reaction in the same manner as in Example 307, oily α -(2'-hydroxybenzylidene)- δ -valerolactone (7.0 g) was obtained. Dimethylformamide (70 ml) and potassium carbonate (9.5 g) were added hereto and (S)-glycidyl nosylate (8.9 g) was added. The mixture was stirred at 40°C for 2 hr. The reaction mixture was concentrated under reduced pressure. Water was added and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the oily title compound (7.8 g). The reaction in the same manner as in Example 307 using this product (1.0 g) and 4-(naphthalen-2-yl)piperidine (1.0 g) in methanol (50 ml) was performed and the obtained oil was recrystallized from ethyl acetate to give the title compound (0.38 g), melting point 142-144°C.

Example 316

(1) α -(2'-hydroxybenzylidene)- γ -valerolactone

Salicylaldehyde (15.2 g) and γ -valerolactone (25 g) were dissolved in toluene (150 ml). By the reactions as in the same manner as in Example 307, the title compound (17 g) was obtained, melting point 112-114°C.

(2) (S)- α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino))-propyloxy)benzylidene)- γ -valerolactone 1/5 hydrate

Dimethylformamide (70 ml) and potassium carbonate (9.5 g) were added to α -(2'-hydroxybenzylidene)- γ -valerolactone (7 g) and (S)-glycidyl nosylate (8.9 g) was added. The mixture was stirred at 40°C for 2 hr. The reaction mixture was concentrated under reduced pressure. Water was added and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oil (7.5 g). By the reaction using this product (1.0 g) and 4-(naphthalen-2-yl)piperidine (1.0 g)

in methanol (50 ml) in the same manner as in Example 307, an oily title compound (0.8 g) was obtained.

$^1\text{H-NMR}(\text{CDCl}_3)\delta$: 1.44 (d, 3H, $J=6.3$), 1.83-2.00 (m, 4H), 2.24 (t, 1H, $J=11.2$), 2.48-2.60 (m, 1H), 2.62-2.82 (m, 4H), 3.02 (d, 1H, $J=12.2$), 3.20 (d, 1H, $J=12.2$), 3.30 (dd, 1H, $J=5.4, 11.7$), 4.00-4.23 (m, 3H), 4.68-4.73 (m, 1H), 7.00 (m, 2H), 7.33-7.45 (m, 6H), 7.64 (s, 1H), 7.77 (d, 2H, $J=8.3$), 8.00 (s, 1H)

Example 317

(S)-3-(2'-(2-hydroxy-3-(4-(5,6,7,8-tetrahydronaphthalen-2-yl)piperidino)propyloxy)benzylidene)-2-pyrrolidone 1/4 hydrate

Sodium hydride (16 g) was suspended in tetrahydrofuran (100 ml) and a solution of N-acetylpyrrolidone (25 g) and o-anisaldehyde (26.8 g) in tetrahydrofuran (100 ml) was added dropwise under ice-cooling. After completion of the reaction, the reaction mixture was poured into water. The mixture was acidified with hydrochloric acid and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate. After filtration, the solvent was evaporated under reduced pressure to give oily 3-(2'-methoxybenzylidene)-2-pyrrolidone (4.7 g). The oil was dissolved in methylene chloride (60 ml). By demethylation using boron tribromide under ice-cooling, 3-(2'-hydroxybenzylidene)-2-pyrrolidone (4.0 g) was obtained as yellow crystals. To the crystals (1.5 g) were added dimethylformamide (50 ml) and potassium carbonate (2.2 g). (S)-Glycidyl nosylate (2.3 g) was added and the mixture was stirred at room temperature for 2 days. The reaction mixture was concentrated under reduced pressure and water was added. The mixture was extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate to give an oily compound (1.8 g). By the reaction of the oil (0.5 g) and 4-(5,6,7,8-tetrahydronaphthalen-2-yl)piperidine (0.5 g) in methanol in the same manner as in Example 307, the title compound (0.4 g) was obtained, melting point 157-159°C.

Example 318

(S)-3-(2'-(3-(4-(3,4-dimethylphenyl)piperidino)-2-hydroxypropyloxy)benzylidene)-2-pyrrolidone 1/4 hydrate

By the method in the same manner as in Example 317 using
5 4-(3,4-dimethylphenyl)piperidine (0.5 g), the title compound
(0.52 g) was obtained, melting point 156-158°C.

Example 319

(S)-3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-2-pyrrolidone

10 By the method in the same manner as in Example 317 using
4-(naphthalen-2-yl)piperidine (0.5 g), the title compound (0.40
g) was obtained, melting point 172-174°C.


Example 320

(R)-3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)-2-pyrrolidone


By the method in the same manner as in Example 317 using
(R)-glycidyl nosylate and 4-(naphthalen-2-yl)piperidine, the
title compound (0.55 g) was obtained, melting point 188-190°C

The structural formulas of the compounds obtained in
20 Examples 311 to 320 are shown in the following.


311




312




313




314




315




316-1




316-2




317




318



319



320



Example 321

(1) 3-(2'-methoxybenzylidene)-1-methyl-2-pyrrolidone

The intermediate 3-(2'-methoxybenzylidene)-2-pyrrolidone (5.0 g) obtained in Example 317 was dissolved in dimethylformamide (50 ml) and sodium hydride (0.99 g) was added thereto under ice-cooling. The mixture was stirred at room temperature for 30 min. Methyl iodide (1.72 ml) was added under ice-cooling and the mixture was stirred for 8 hr at room temperature. After completing of the reaction, the reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (3.6 g).

¹H-NMR(CDCl₃): 3.00 (m, 5H), 3.46 (t, 2H), 3.85 (s, 3H), 6.91 (d, J=8.2, 1H), 6.96 (t, J=8.2, 1H), 7.20-7.30 (m, 1H), 7.41 (d, J=8.2, 1H), 7.70 (m, 1H)

(2) (S)-3-(2'-(2-hydroxy-3-(4-(5,6,7,8-tetrahydronaphthalen-2-yl)piperidino)propyloxy)benzylidene)-1-methyl-2-pyrrolidone dihydrochloride

3-(2'-Methoxybenzylidene)-1-methyl-2-pyrrolidone (3.6 g) was dissolved in methylene chloride (50 ml). Demethylation was performed using boron tribromide (12.7 g) under ice-cooling to give white crystals (3.0 g). To the crystals (1.6 g) were added dimethylformamide (50 ml) and potassium carbonate (2.2 g), and (S)-glycidyl nosylate (2.3 g) was added. The reaction mixture was stirred at room temperature for 2 days. The reaction mixture was concentrated under reduced pressure. Water was added to the residue and the mixture was extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate to give an oily compound (2.1 g). By the reaction of the oil (1.0 g) and 4-(5,6,7,8-tetrahydronaphthalen-2-yl)piperidine (0.5 g) in methanol in the same manner as in Example 317, the oily title compound was obtained. By the treatment of the oily compound in a mixed

solvent of hydrochloric acid and methanol, the title compound (0.026 g) was obtained, melting point 227-230°C.

Example 322

(S)-3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino))-propyloxy)benzylidene)-1-methyl-2-pyrrolidone dihydrochloride

By the methods in the same manner as in Example 317 using 4-(naphthalen-2-yl)piperidine (1.0 g), the title compound (0.33 g) was obtained, melting point 136-139°C.

Example 323

(R)-3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino))-propyloxy)benzylidene)-1-methyl-2-pyrrolidone monohydrochloride monohydrate

By the methods in the same manner as in Example 317 using (R)-glycidyl nosylate and 4-(naphthalen-2-yl)piperidine (0.6 g), the title compound (0.40 g) was obtained, melting point 127-129°C.

Example 324

(1) 3-(2'-methoxybenzylidene)-1-(2-methoxyethyl)-2-pyrrolidone

The intermediate 3-(2'-methoxybenzylidene)-2-pyrrolidone (5.0 g) obtained in Example 317 was dissolved in dimethylformamide (50 ml) and sodium hydride (0.99 g) was added under ice-cooling. The mixture was stirred at room temperature for 30 min. To the reaction mixture was again added 2-methoxyethyl chloride (2.3 ml) under ice-cooling, and the mixture was stirred at room temperature for 1 hr and at 70°C for 1 hr. After cooling, the reaction mixture was poured into water and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give an oily title compound (2.8 g).

¹H-NMR(CDCl₃): 2.98-3.02 (m, 2H), 3.37-3.39 (m, 2H), 3.42-3.65 (m, 4H), 3.81 (s, 3H), 3.85 (s, 3H), 6.87-6.98 (m, 2H), 7.22 (t, J=8.2, 1H), 7.29 (d, J=8.2, 1H), 7.71 (bs, 1H)

(2) (S)-3-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino))-propyloxy)benzylidene)-1-(2-methoxyethyl)-2-pyrrolidone

hydrochloride

3-(2'-Methoxybenzylidene)-1-(2-methoxyethyl)-2-pyrrolidone (2.8 g) was dissolved in methylene chloride (30 ml) and demethylated with boron tribromide (8.7 g) under ice-cooling to give an oil (1.68 g). To the oil were added dimethylformamide (50 ml) and potassium carbonate (2.2 g). (S)-Glycidyl nosylate (2.3 g) was added and the mixture was stirred at room temperature for one day. The reaction mixture was concentrated under reduced pressure, and water was added. The reaction mixture was extracted with chloroform and the organic layer was dried over anhydrous sodium sulfate to give an oily compound (0.8 g). The oil (1.0 g) and 4-(naphthalen-2-yl)piperidine (0.8 g) were reacted in methanol in the same manner as in Example 317 to give an oily title compound. The oily compound was treated with a mixed solvent of hydrochloric acid and methanol to give the title compound (0.075 g), melting point 254-257°C.

Example 325

(S)- α -(2'-(2-hydroxy-3-(4-(6-methoxynaphthalen-2-yl)piperidino)propyloxy)benzylidene)- γ -butyrolactone

(S)- α -(2'-(2,3-Epoxypropan-1-yloxy)benzylidene)- γ -butyrolactone (1.2 g) and 4-(6-methoxynaphthalen-2-yl)piperidine (1.2 g) were dissolved in methanol (50 ml), and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the residue was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (1.0 g), melting point 148-150°C.

Example 326

(Z)-(S)- α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone

(S)- α -(2'-(2-Hydroxy-3-(4-(naphthalen-2-yl)piperidino)propyloxy)benzylidene)- γ -butyrolactone (2.0 g) was dissolved in ethanol (300 ml) and exposed to the sunlight

for 6 hr. The solvent was concentrated under reduced pressure and the obtained oil was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (0.3 g).

5 $^1\text{H-NMR}(\text{CDCl}_3)$: 1.81-2.05(m, 4H), 2.17-2.25(m, 1H), 2.50-2.80(m, 4H), 3.06(d, 1H, $J=10.7$), 3.15(dt, 2H, $J=2.0, 5.4$), 3.24(d, 1H, $J=10.7$), 4.00-4.05(m, 2H), 4.18-4.28(m, 1H), 4.40(t, 2H, $J=7.3$), 6.90(d, 1H, $J=8.3$), 6.97(t, 1H, $J=7.5$), 7.30-7.558(m, 4H), 7.66(s, 1H), 7.70-7.88(m, 3H), 7.90(d, 1H, $J=7.8$)

10 **Example 327**

(1) α -(2'-hydroxybenzyl)- γ -butyrolactone

α -(2'-Hydroxybenzylidene)- γ -butyrolactone (5.0 g) was dissolved in ethanol (400 ml) and 10% palladium carbon (0.5 g) was added. The reaction mixture was reduced for 6 hr at 50 atm.

15 The reaction mixture was filtered and the organic solvent was concentrated to give the title compound (5.0 g).

$^1\text{H-NMR}(\text{CDCl}_3)$: 2.05-2.20(m, 1H), 2.12-2.39(m, 1H), 2.92-2.97(m, 1H), 2.97-3.01(m, 1H), 3.05-3.20(m, 1H), 4.18-4.23(m, 1H), 4.28-4.38(m, 1H), 6.80-6.92(m, 2H), 7.05-7.20(m, 2H)

20 (2) (2S)- α -(2'-(2,3-epoxypropan-1-yloxy)benzyl)- γ -butyrolactone

Dimethylformamide (100 ml) and potassium carbonate (10 g) were added to α -(2'-hydroxybenzyl)- γ -butyrolactone (5.0 g), and (S)-glycidyl nosylate (6 g) was added. The mixture was stirred at room temperature for 2 days. The reaction mixture
25 was concentrated under reduced pressure and water was added. The reaction mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (6.1 g).

30 $^1\text{H-NMR}(\text{CDCl}_3)$: 1.90-2.05(m, 1H), 2.06-2.30(m, 1H), 2.62-3.00(m, 5H), 3.28-3.40(m, 2H), 3.92-4.00(m, 1H), 4.06-4.20(m, 1H), 4.22-4.38(m, 1H), 6.82-6.90(m, 1H), 6.91-6.98(m, 1H), 7.03-7.18(m, 2H)

(3) (2S)- α -(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzyl)- γ -butyrolactone

(2S)- α -(2'-(2,3-Epoxypropan-1-yloxy)benzyl)- γ -butyrolactone (1.5 g) and 4-(naphthalen-2-yl)piperidine (1.5 g) were dissolved in methanol (50 ml) and the reaction mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the obtained oil was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (1.3 g).

$^1\text{H-NMR}(\text{CDCl}_3)$: 1.80-2.00 (m, 5H), 2.10-2.22 (m, 1H), 2.48-2.58 (bs, 1H), 2.60-2.78 (m, 4H), 2.98-3.05 (m, 1H), 3.07-3.09 (bs, 1H), 3.37-3.40 (m, 1H), 4.00-4.04 (m, 2H), 4.06-4.23 (m, 2H), 4.25-4.32 (m, 1H), 6.80-6.92 (m, 2H), 7.15-7.20 (m, 1H), 7.22-7.25 (m, 1H), 7.37-7.43 (m, 3H), 7.66 (s, 1H), 7.77-7.82 (m, 3H)

Example 328

(1) N-(2'-hydroxybenzyl)-2-oxazolidone

2-Oxazolidone (3.0 g) was dissolved in dimethylformamide (120 ml) and sodium hydride (2.5 g) was added under ice-cooling. The mixture was stirred at room temperature for 1 hr and 2-methoxybenzyl chloride (6.5 g) was again added under ice-cooling. The mixture was heated to 60°C and stirred for 1 hr. The mixture was allowed to reach room temperature and poured into ice-water. The aqueous layer was extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The organic solvent was concentrated under reduced pressure to give an oil (8.0 g). Methylene chloride (100) was added thereto, and boron tribromide (9 ml) was added dropwise under stirring at -78°C. Then, the reaction mixture was stirred for 2 hr under ice-cooling. The reaction mixture was poured into ice water and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give the title compound (5.5 g).

$^1\text{H-NMR}(\text{CDCl}_3)$: 3.43 (t, 2H, J=8.3), 3.82 (s, 3H), 4.28 (t, 2H,

J=8.3), 4.42(s, 2H), 6.81-6.97(m, 2H), 7.24-7.30(m, 2H)


(2) (S)-N-(2'-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzyl)-2-oxazolidone

Dimethylformamide (100 ml) and potassium carbonate (10 g) were added to N-(2'-hydroxybenzyl)-2-oxazolidone (5.0 g) and (S)-glycidyl nosylate (6 g) was added, and then the mixture was stirred at room temperature for 2 days. The reaction mixture was concentrated under reduced pressure and water was added. The reaction mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was washed with hexane to give an oil (6.0 g). The oil (1.5 g) and 4-(naphthalen-2-yl)piperidine (1.5 g) were dissolved in methanol (50 ml), and the mixture was refluxed under heating for 2 hr. After cooling, the solvent was evaporated under reduced pressure and the obtained oil was purified by silica gel column chromatography (chloroform/methanol) to give the title compound (2.7 g).


¹H-NMR(CDCl₃): 1.80-2.00(m, 4H), 2.20-2.25(m, 1H), 2.32-2.40(m, 1H), 2.58-2.75(m, 3H), 3.10(t, 2H, J=9.7), 3.39(t, 2H, J=8.3), 3.91-3.98(m, 1H), 4.04-4.10(m, 1H), 4.17-4.15(m, 3H), 4.42-4.60(m, 2H), 6.82-6.95(m, 2H), 7.19-7.30(m, 1H), 7.31-7.38(m, 1H), 7.39-7.50(m, 3H), 7.65(s, 1H), 7.77-7.82(m, 3H)

The structural formulas of the compounds obtained in Examples 321 to 328 are shown in the following.


321-1




321-2




322




323




324-1




324-2




325




326




327-1




327-2




327-3



328-1



328-2



By the reactions in the same manner, the following

compounds can be synthesized.

Example 401

(S)-1-(4-(1-methyl-2-oxoindolin-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

5 **Example 402**

(S)-1-(4-(1H-indol-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 403

10 (S)-1-(4-(1-methylindol-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 404

(S)-1-(4-(2,2-dimethyl-2,3-dihydrobenzo(b)furan-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

15 **Example 405**

(S)-1-(4-(4-chloro-2,2-dimethyl-2,3-dihydrobenzo(b)thiophen-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 406

20 (S)-1-(4-(2-methyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 407

25 (S)-1-(4-(2,4,6-trimethylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 408

(S)-1-(4-(3,5-dichlorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 409

30 (S)-1-(4-(2-methylbenzo(b)furan-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 410

(S)-1-(4-(3-methylbenzo(d)isoxazol-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 411


(S)-1-(4-(2H-1-oxoisoindolin-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 412

- 5 (S)-1-(4-(2-methyl-1-oxoisoindolin-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

The structural formulas of the compounds obtained in Examples 401 to 412 are shown in the following Table 1.

Table 1



Ex. No.	Substituent R ¹	Ex. No.	Substituent R ¹
401		407	
402		408	
403		409	
404		410	
405		411	
406		412	

Example 413

- 5 (S)-1-(4-(quinolin-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 414

(S)-1-(4-(isoquinolin-6-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 415

5 (S)-1-(4-(quinolin-3-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 416

(S)-1-(4-(7-methyl-2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-
10 propanol

Example 417

(S)-1-(4-(7-chloro-2,3-dihydrobenzo(b)furan-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-
propanol

15 **Example 418**

(S)-1-(4-(1H-2-oxoindolin-5-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 419

(S)-1-(4-(3-chloro-4-fluorophenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 420

(S)-1-(4-(4,5-dimethylthiophen-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 421

25 (S)-1-(4-(4,5-dichlorothiophen-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

Example 422

(S)-1-(4-(2-methylpyridin-4-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

30 **Example 423**

(S)-1-(4-(4,5-dimethylfuran-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol


Example 424

(S)-1-(4-(4,5-dichlorofuran-2-yl)piperidino)-3-(2-(5-methyl-

1,3,4-oxadiazol-2-yl)benzo(b)furan-4-yloxy)-2-propanol

The structural formulas of the compounds obtained in Examples 413 to 424 are shown in the following Table 2.

5 Table 2



Ex. No.	Substituent R ¹	Ex. No.	Substituent R ¹
413		419	
414		420	
415		421	
416		422	
417		423	
418		424	

Example 425

(S)-1-(4-(4-chloro-2-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

5 (S)-2-(4-Glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole (7.0 g) and 4-(4-chloro-2-methylphenyl)piperidine (6 g) were heated in methanol for 3 h with stirring. The solvent was evaporated and an oil was purified by silica gel column chromatography (chloroform/methanol). The obtained oil was
10 dissolved in acetone and hydrochloric acid - ethanol was added to give the title compound (6.8 g), melting point 201 - 203°C.

Example 426

(S)-1-(4-(2,6-dimethoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol 1/2

15 terephthalate 1/2 hydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(2,6-dimethoxyphenyl)piperidine, melting point 170 - 171°C.

20 **Example 427**

(S)-1-(4-(3-fluoro-4-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride

In the same manner as in Example 425, the title compound
25 was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(3-fluoro-4-methylphenyl)piperidine, melting point 214 - 216°C.

Example 428

(S)-1-(4-(2,4,6-trimethoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol 1/2
30 terephthalate 1/2 hydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(2,4,6-

trimethoxyphenyl)piperidine, melting point 209 - 211°C.

Example 429

(S)-1-(4-(4-chloro-2,6-dimethoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol
5 1/2 terephthalate dihydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(4-chloro-2,6-dimethoxyphenyl)piperidine, melting point 231 - 232°C.

10 **Example 430**

(S)-1-(4-(3-chloro-4-ethoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol
hydrochloride

In the same manner as in Example 425, the title compound
15 was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(3-chloro-4-ethoxyphenyl)piperidine, melting point 212 - 214°C.

Example 431

(S)-1-(4-(3-chloro-4-isopropoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol
20 hydrochloride

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(3-chloro-4-isopropoxyphenyl)piperidine, melting point 204 - 206°C.

Example 432

(S)-1-(4-(4-methoxy-2-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol
hydrochloride monohydrate

30 In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(4-methoxy-2-methylphenyl)piperidine, melting point 190 - 192°C.

Example 433

(S)-1-(4-(5-chloro-4-methoxy-2-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(5-chloro-4-methoxy-2-methylphenyl)piperidine, melting point 240 - 242°C.

Example 434

(S)-1-(4-(2,4-dimethoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol terephthalate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(2,4-dimethoxyphenyl)piperidine, melting point 192 - 194°C.

Example 435

(S)-1-(4-(4-chloro-2-fluoro-3-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(4-chloro-2-fluoro-3-methylphenyl)piperidine, melting point 218 - 220°C.

Example 436

(S)-1-(4-(4-fluoro-2-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(4-fluoro-2-methylphenyl)piperidine, melting point 245 - 247°C.

Example 437

(S)-1-(4-(3-chloro-4-methoxy-5-methylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(3-chloro-4-methoxy-5-methylphenyl)piperidine, melting point 222 - 224°C.

5 **Example 438**

(S)-1-(4-(1-methoxynaphthalen-2-yl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride 1/2 hydrate

In the same manner as in Example 425, the title compound
10 was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(1-methoxynaphthalen-2-yl)piperidine, melting point 149 - 151°C.

Example 439

(S)-1-(4-(3,4-dimethyl-2-methoxyphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride monohydrate
15

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(3,4-dimethyl-2-methoxyphenyl)-
20 piperidine, melting point 214 - 216°C.

Example 440

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-3-(4-(2,4,6-trimethylphenyl)piperidino)-2-propanol hydrochloride

25 In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(2,4,6-trimethylphenyl)piperidine, melting point 158 - 160°C.

Example 441

30 (S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-3-(4-(3-methylthiophenyl)piperidino)-2-propanol hydrochloride

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(3-methylthiophenyl)piperidine,

melting point 176 - 178°C.

Example 442

(S)-1-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-
3-(4-(4-methylthiophenyl)piperidino)-2-propanol hydrochloride
5 1/2 hydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(4-methylthiophenyl)piperidine, melting point 180 - 182°C.

10 **Example 443**

(S)-1-(4-(indolin-1-yl)piperidino)-3-(2-(5-methyl-1,3,4-
oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol 3/2
terephthalate 1/2 hydrate

In the same manner as in Example 425, the title compound
15 was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(indolin-1-yl)piperidine, melting point 182 - 184°C.

Example 444

(S)-1-(4-(indol-1-yl)piperidino)-3-(2-(5-methyl-1,3,4-
20 oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol
monoterephthalate 1/2 hydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(indol-1-yl)piperidine, melting
25 point 145 - 147°C.

Example 445

(S)-1-(4-(4-chloro-3-ethylphenyl)piperidino)-3-(2-(5-methyl-
1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol
hydrochloride

30 In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(4-chloro-3-ethylphenyl)-piperidine, melting point 177 - 178°C.

Example 446

(S)-1-(4-(4-chloro-3-isopropylphenyl)piperidino)-3-(2-(5-methyl-1,3,4-oxadiazol-2-yl)benzo(b)fran-4-yloxy)-2-propanol hydrochloride

In the same manner as in Example 425, the title compound was synthesized from (S)-2-(4-glycidyloxybenzo(b)fran-2-yl)-5-methyl-1,3,4-oxadiazole and 4-(4-chloro-3-isopropylphenyl)-piperidine, melting point 154 - 156°C.

Example 447

(S)-3-(3-(3-(4-(3,4-dichlorophenyl)piperidino)-2-hydroxypropyloxy)benzyl)oxazolidin-2-one terephthalate

In the same manner as in Example 425, the title compound was synthesized from (S)-3-(3-glycidyloxybenzyl)oxazolidin-2-one and 4-(3,4-dichlorophenyl)piperidine, melting point 157 - 159°C.

Example 448

(S)-1-(3-(3-(4-(3,4-dichlorophenyl)piperidino)-2-hydroxypropyloxy)benzyl)pyrrolidin-2-one terephthalate 1/2 hydrate

In the same manner as in Example 425, the title compound was synthesized from (S)-1-(3-glycidyloxybenzyl)pyrrolidin-2-one and 4-(3,4-dichlorophenyl)piperidine, melting point 132 - 134°C.

Example 449

(S)-1-(3-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzyl)pyrrolidin-2-one terephthalate

In the same manner as in Example 425, the title compound was synthesized from (S)-1-(3-glycidyloxybenzyl)pyrrolidin-2-one and 4-(naphthalen-2-yl)piperidine, melting point 153 - 155°C.

Example 450

(S)-1-(2-(2-hydroxy-3-(4-(3,4-methylenedioxyphenyl)piperidino)-propyloxy)benzyl)pyrrolidin-2-one terephthalate

In the same manner as in Example 425, the title compound was synthesized from (S)-1-(2-glycidyloxybenzyl)pyrrolidin-2-

one and 4-(3,4-methylenedioxyphenyl)piperidine, melting point 169 - 171°C.

Example 451

(S)- α -(3-(3-(4-(3,4-dichlorophenyl)piperidino)-2-hydroxypropyloxy)benzylidene)- γ -butyrolactone hydrochloride

In the same manner as in Example 425, the title compound was synthesized from (S)- α -(3-glycidyloxybenzylidene)- γ -butyrolactone and 4-(3,4-dichlorophenyl)piperidine, melting point 147 - 149°C.


Example 452

(S)- α -(3-(2-hydroxy-3-(4-(naphthalen-2-yl)piperidino)-propyloxy)benzylidene)- γ -butyrolactone hydrochloride


In the same manner as in Example 425, the title compound was synthesized from (S)- α -(3-glycidyloxybenzylidene)- γ -butyrolactone and 4-(naphthalen-2-yl)piperidine, melting point 82 - 84°C.

The structural formulas of the compounds obtained in Examples 425 - 452 are shown in the following.


425




430




426




431




427




432




428




433




429




434




435




436




437




438




439




440




441




442




443




444




445




449




446




450




447




451



448



452



Formulation Example

Of the compounds of the present invention, a compound of the formula (I) (50 mg) is thoroughly kneaded with lactose (98 mg), cornstarch (45 mg) and hydroxypropylcellulose (3 mg) in a kneader. The kneaded product is passed through a 200 mesh sieve, dried at 50°C and passed through a 24 mesh sieve. The resulting product is mixed with talc (3 mg) and magnesium stearate (1 mg) and compressed with a 9 mm diameter poulder to give a tablet weighing 200 mg. The tablets may be sugar coated or film coated as necessary.

Experimental Examples 1 - 5 are shown in the following. In Experimental Examples 1, 2 and 5, (S)-1-(4-indolyloxy)-3-[4-hydroxy-4-(2-naphthyl)piperidino]propan-2-ol described in WO97/48698 was used as a compound for comparison.

15 Experimental Example 1: 5-HT_{1A} receptor binding test

The experiment was conducted according to the method of M.D. Hall et al (J. Neurochem. 44, 1685-1696 (1985)).

Cryopreserved rat hippocampus was homogenized in a 20-fold wet weight amount of 50 mM Tris-HCl buffer (pH 7.4), and the homogenate was centrifuged at 500xg for 10 min. The supernatant was centrifuged at 40000xg for 10 min and the sediment was incubated at 37°C for 10 min, which was followed by centrifugation at 40000xg for 10 min. To the sediment was added a 20-fold amount of 50 mM Tris-HCl buffer (pH 7.4) and the mixture was homogenized, which was followed by centrifugation again at 40000xg for 10 min. 50 mM Tris-HCl buffer (pH 7.4, 100-fold volume) containing 1 mM MnCl₂ was added to the sediment and the mixture was homogenized, which was used as a membrane solution. To a 96 well plate were successively added a test solution (25 ml), (³H)-8-OH-DPAT ((±)-8-hydroxy-2-(DL-N-propylamino)tetraline, Sigma, lot No. 57H4131) solution (final concentration 2 nM, 25 ml) and the membrane solution (0.45 ml) preincubated at 37°C, and incubated at 37°C for 12 min. After completion, the reaction mixture was

filtered through a GF/B glass filter and the filter was washed 5 times with 50 mM Tris-HCl buffer (pH 7.4). The radioactivity left on the filter was measured with a Top Count. For total binding measurement, 0.005N hydrochloric acid (25 ml) was used, and for the measurement of nonspecific binding, a test solution containing WAY-100635 (final concentration 1M, 25 ml) instead of the test substance was used. The total binding and nonspecific binding were measured in quadruplicate, and the test substance was measured in duplicate.

The IC_{50} value was calculated by two-point interpolation and K_i value was calculated according to the following equation using K_d value obtained from each measurement.

$$K_i = IC_{50} / (1 + C / K_d)$$

IC_{50} : concentration of 50% binding inhibition

C: concentration of ligand

The results are shown in Table 3 below.

Experimental Example 2: 5-HT transporter binding test

The experiment was conducted according to the method of Habert, E. et al (Eur. J. Pharmacol., 118; 107-114 (1985)).

Rat brain cortex was homogenized using Polytron in ice-cooled 50 mmol/L Tris-HCl buffer (pH 7.4). After centrifugation at 1000×g and 4°C for 10 min, the supernatant was transferred to a different centrifugation tube. This was centrifuged at 40000×g and 4°C for 20 min, and 50 mmol/L Tris-HCl buffer (pH 7.4) was added to the sediment to give a suspension. This was incubated at 37°C for 10 min, centrifuged at 40000×g and 4°C for 20 min, and suspended in 50 mmol/L Tris-HCl buffer (pH 7.4) (diluted 100-fold of brain wet weight) containing 120 mmol/L NaCl and 5 mmol/L KCl, which was used as a membrane solution. For binding inhibition test, it was reacted with (³H) paroxetine prepared to the final concentration of 0.2 nmol/L in a plastic test tube at 25°C for 90 min. For total binding, a solvent was used and for nonspecific binding, fluvoxamine having a final concentration

of 10 $\mu\text{mol/L}$ was used.

Using a cell harvester, the reaction mixture was filtered through a GF/B glass filter treated with 0.1% polyethyleneimine to stop the reaction and washed 3 times with 3 mL of ice-cooled 50 mmol/L Tris-HCl buffer (pH 7.4). The radioactivity was measured using a β plate.

The results are shown in Table 3.

Table 3 Test results of Experimental Examples 1 and 2

	5-HT _{1A} receptor binding Ki value (nM)	5-HT transporter binding Ki value (nM)
compound for reference	0.16	55
compound of Example 6	2.3	1.10
compound of Example 88	0.75	0.32
compound of Example 136	0.37	0.18
compound of Example 138	0.68	1.60

As is evident from Table 3, the compound of the present invention showed strong affinity for both 5-HT_{1A} receptor and 5-HT transporter.

Experimental Example 3: antagonistic action against lowering of body temperature

From the antagonistic action of the test substance against decrease in the body temperature due to 8-OH-DPAT, transfer of the test substance into the brain was established. At the same time, it was clarified if the test substance acts as an agonist or as an antagonist on the 5-HT_{1A} receptor.

The rectal temperature of male ddY mice was measured with a digital thermostat (KN-91, Natsume) (pre-value). Thereafter, the test substance was administered orally or parenterally, and after a certain time, 8-OH-DPAT (1 mg/kg) was subcutaneously administered. The rectal temperature was

measured 30 min later (post-value).

The pre-value and post-value obtained by the measurement were compared, and the action of the test substance on the decrease of body temperature due to 8-OH-DPAT was observed.

5 The results of Experimental Example 3 establish that the compound of the present invention acts as an antagonist on 5-HT_{1A} receptor, because the compound given orally in 0.1 - 100 mg/kg antagonizes the lowering of the body temperature due to 8-OH-DPAT. From the results, it is suggested that the compound
10 of the present invention is superior in the bioavailability and transfer into the brain.

Experimental Example 4: forced swimming test

The test substance was administered orally or parenterally to male ddY mice, and after a certain time, the
15 mice were placed in a water tank (material: vinyl chloride, color: black, inner diameter: 10 cm, height: 25 cm, water depth: 15 cm, water temperature: 25°C), and subjected to 6 min test trial. The movement of the animal was videotaped with a CCD camera set right above the water tank, and analyzed against
20 immobility time during 4 minutes from 2 to 6 min after the start of swimming, using an image analysis system/forced swimming analysis program [Neuroscience Inc.: Videoimage motion analyzer (AXIS series)/(TARGET/7M)].

The results of Experimental Example 4 reveal that, while
25 the conventional SSRI requires several days for expression of an action, the compound of the present invention significantly shortened the immobility time by the single oral administration of 0.1 - 100 mg/kg thereof. From this, it is suggested that the compound of the present invention can be a so-called rapid
30 onset antidepressant that shows quick expression of the anti-depressive effect, as compared to conventional SSRI.

Experimental Example 5: test for electrophysical evaluation of agonist/antagonist of 5-HT_{1A} receptor

The experiment followed the method of Katayama et al

(Brain Res.; 745, 283-292, 1997).

The brain of 2-week-old male Wistar rat was extracted, and a brain thin section (thickness 350 μ m) containing dorsal raphe nuclei was prepared using a microslicer. The brain thin section was treated in Ringer's solution containing pronase (0.4 mg/mL) and protease type X (0.25 mg/mL) at 30°C for 25 min and 15 min, respectively, and the dorsal raphe nuclei region was micropunched out. The brain thin section punched out was subjected to pipetting in a culture dish filled with the Ringer's solution to isolate the nerve cell. The isolated nerve cell (dorsal raphe nuclei cell) was subjected to nystatin perforated patch clamp method (Akaike & Harata, Jpn. J. Physiol.; 44, 433-437, 1994) and the inward K^+ current induced by the test substance and the like was measured under membrane voltage-clamp condition ($V_H = -60$ mV). For the measurement of the inward K^+ current via the 5-HT_{1A} receptor, an extracellular solution and a patch pipette solution having the following compositions were used simultaneously.

extracellular solution (mmol/L): NaCl, 135; KCl, 20; MgCl₂, 1; CaCl₂, 2; D-glucose, 10; HEPES, 10; tetrodotoxin, 3×10^{-4} ; LaCl₃, 10

patch pipette solution (mmol/L): KCl, 150; HEPES, 10

To the above-mentioned patch pipette solution was added nystatin (Sigma, Lot No. 33H0762) to the final concentration of 75 μ g/mL before the electric measurement.

The current response was measured with a voltage clamp amplifier (List Medical, L/M-EPC7), and the obtained results were recorded on a chart of a recticorder (Nihondenki Sanei, RECTIHORIZ-8K), digitized by a PCM recording device (NF electric Instruments, RP-882) and videotaped by a VCR (Panasonic, NV-G40).

The administration of the test substance and the like followed a Y-tube method (Murase et al., Brain Res.; 525, 84-91, 1990).

First, 8-OH-DPAT (10^{-7} mol/L) was administered to the isolated nerve cell (dorsal raphe nuclei cell) obtained above, and the level of the inward K^{+} current response was measured. The cells that showed an inward K^{+} current response of not less
5 than 15 pA were subjected to the following test.

8-OH-DPAT was washed out for 2 min from the above-mentioned cells and the test substance and pindolol (reference substance which is an antagonist) were administered for 1 min, and the level of the inward K^{+} current response by each of them
10 was measured. From immediately after administration of the test substance or the reference substance, a mixture of 8-OH-DPAT (10^{-7} mol/L) and the test substance or reference substance was administered, and the level of the induced inward K^{+} current response was measured. The measurement results and the
15 level of the current response in the same cell by 8-OH-DPAT alone were compared, based on which the antagonistic action on the 8-OH-DPAT induction current was considered. Every current response was expressed upon standardizing according to the following calculation equation based on the level of the 8-OH-
20 DPAT (10^{-7} mol/L) induction current in the same cell.

The inward K^{+} current response inducing action (5-HT_{1A} agonistic action) by the test substance alone was determined from the following equation.

inward K^{+} current response inducing action (%) by test
25 substance alone = $(ITD \div IDPAT) \times 100$

ITD: level of inward K^{+} current response by test substance alone

IDPAT: level of 8-OH-DPAT (10^{-7} mol/L) induced inward K^{+} current in the same cell

30

The suppressive effect (5-HT_{1A} antagonistic action) on the 8-OH-DPAT induced inward K^{+} current by the test substance was determined according to the following equation.

antagonistic action of test substance on 8-OH-DPAT
induced inward K^+ current (% of control) = $(IMIX \div IDPAT) \times 100$

IMIX: level of 8-OH-DPAT (10^{-7} mol/L) induced inward K^+
current in the presence of test substance

5 IDPAT: as defined above

According to the results of Experimental Example 5, the
respective values of inward K^+ current inducing action (%) by
the test substance alone at a concentration of 10^{-7} mol/L and
10 the antagonistic action (% of control) of the test substance on
the 8-OH-DPAT induced inward K^+ current were almost nil,
showing that the compounds of the present invention (compounds
of Examples 6, 88, 136, 138 etc.) are silent antagonists of 5-
HT_{1A} receptor.

15 In contrast, the compound for reference showed an action
of a partial agonist at a high dose (10^{-7} mol/L).

Effect of the Invention

The compound of the present invention shows selective
affinity for as well as simultaneous antagonistic activity
20 against 5-HT_{1A} receptor, and also shows a 5-HT reuptake
inhibitory activity. Thus, the compound of the present
invention is useful as a so-called rapid onset antidepressant
that shows quick expression of an anti-depressive effect. It
is also useful for the treatment of 5-HT mediated diseases of
25 the central nervous system, such as schizophrenia, anxiety
neurosis, obsessive-compulsive disorder (OCD), panic disorder,
social anxiety disorder (social phobia), seasonal emotional
disorder (seasonal affective disorder), Anorexia Nervosa,
Bulimia Nervosa, nocturnal enuresis, children's hyperlocomotion,
30 post-traumatic stress disorder (PTSD), senile dementia,
hemicrania, stroke, Alzheimer's disease, recognition disorder,
hypertension, gastrointestinal injury, feeding disorders,
premenstrual syndrome (PMS), abnormal body temperature

regulation and sexual disorder, pain, abnormality in the cardiovascular system, drug abuse and the like.

This application is based on patent application Nos.
5 142750/1999, 166160/1999, 277384/1999 and 018080/2000 filed in Japan, the contents of which are hereby incorporated by reference.